

# Calculus II

## Polar coordinates

Todor Milev

2019

# Outline

## 1 Polar Coordinates

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called  $O$  (the origin).
- Draw a ray starting at  $O$ . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- Let  $P$  be a point in the plane.
- Let  $\theta$  denote the angle between the polar axis and the line  $OP$ .
- Let  $r$  denote the length of the segment  $OP$ .
- Then  $P$  is represented by the ordered pair  $(r, \theta)$ .



# Polar Coordinates

- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called  $O$  (the origin).
- Draw a ray starting at  $O$ . The ray is called the polar axis. This ray is usually drawn horizontally to the right.







- The letters  $(x, y)$  imply Cartesian coordinates and the letters  $(r, \theta)$ - polar. When we use other letters, it should be clear from context whether we mean Cartesian or polar coordinates. If not, one must request clarification.

- 1 What if  $\theta$  is negative?
- 2 What if  $r$  is negative?
- 3 What if  $r$  is 0?



- 1 Positive angles  $\theta$  are measured in the counterclockwise direction from  $O$ . Negative angles are measured in the clockwise direction.
- 2 Points with polar coordinates  $(-r, \theta)$  and  $(r, \theta)$  lie on the same line through  $O$  and at the same distance from  $O$ , but on opposite sides.
- 3 If  $r = 0$ , then  $(0, \theta)$  represents  $O$  for all values of  $\theta$ .





- There are many ways to represent the same point.
- We could use a negative  $\theta$ .
- We could go around more than once.
- We could use a negative  $r$ .

- Let  $P_1$  be point with polar coordinates  $(r_1, \theta_1)$ .
- Let  $P_2$  be point with polar coordinates  $(r_2, \theta_2)$ .

## Observation

$P_1$  coincides with  $P_2$  if one of the three mutually exclusive possibilities holds:

- $r_1 = r_2 \neq 0$  and  $\theta_2 = \theta_1 + 2k\pi, k \in \mathbb{Z}$ ,
- $r_1 = -r_2 \neq 0$  and  $\theta_2 = \theta_1 + (2k + 1)\pi, k \in \mathbb{Z}$ ,
- $r_1 = r_2 = 0$  and  $\theta$  is arbitrary.






- Let  $P_1$  be point with polar coordinates  $(r_1, \theta_1)$ .
- Let  $P_2$  be point with polar coordinates  $(r_2, \theta_2)$ .


## Observation

$P_1$  coincides with  $P_2$  if one of the three mutually exclusive possibilities holds:

- $r_1 = r_2 \neq 0$  and  $\theta_2 = \theta_1 + 2k\pi, k \in \mathbb{Z}$ ,
- $r_1 = -r_2 \neq 0$  and  $\theta_2 = \theta_1 + (2k + 1)\pi, k \in \mathbb{Z}$ ,
- $r_1 = r_2 = 0$  and  $\theta$  is arbitrary.



- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates  $(r, \theta)$  and Cartesian coordinates  $(x, y)$ .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

---


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

$$r^2 = x^2 + y^2$$


---

$$r = \sqrt{x^2 + y^2}$$

$$\theta = \arcsin\left(\frac{y}{r}\right) \quad \text{if } x > 0$$

$$= \arccos\left(\frac{x}{r}\right) \quad \text{if } y > 0$$

$$= \arctan\left(\frac{y}{x}\right) \quad \text{if } x > 0$$

## Example


Convert the point  $(2, \frac{\pi}{3})$  from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left( \frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3} = 2 \left( \frac{\sqrt{3}}{2} \right) = \sqrt{3}$$

Therefore the point with polar coordinates  $(2, \frac{\pi}{3})$  has Cartesian coordinates  $(1, \sqrt{3})$ .

## Example



Represent the point with Cartesian coordinates  $(1, -1)$  in terms of polar coordinates.

- Suppose  $r$  is positive.
- $\tan \theta = -1$  for  $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$ , and many other angles.
- $(1, -1)$  is in the fourth quadrant.
- Of the two values above, only  $\theta = \frac{7\pi}{4}$  gives a point in the fourth quadrant.
- $\Rightarrow$  one representation of  $(1, -1)$  in polar coordinates is  $(\sqrt{2}, \frac{7\pi}{4})$ .
- $(\sqrt{2}, -\frac{\pi}{4})$  is another.

$$\begin{aligned} r &= \pm \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\begin{aligned} \tan \theta &= \frac{y}{x} \\ &= -1 \end{aligned}$$