

# Calculus I

## Definite integrals and areas between curves

Todor Milev

2019

# Outline

## 1 Integration and symmetry

# Outline

1 Integration and symmetry

2 More About Areas

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_{-a}^0 f(x) dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is **even** (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is **odd** (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_{-a}^0 f(x) dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) \, dx = \int_0^a f(x) \, dx + \int_{-a}^0 f(x) \, dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_{-a}^0 f(x) dx + \int_0^a f(x) dx = 0.$$





# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is **odd** (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_{-a}^0 f(x) dx + \int_0^a f(x) dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_{-a}^0 f(x) dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_{-a}^0 f(x) dx = 0.$$



# Symmetry



## Theorem (Integrals of Symmetric Functions)

Suppose  $f$  is continuous on  $[-a, a]$ .

① If  $f$  is even (that is,  $f(-x) = f(x)$ ), then  $\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx$ .

② If  $f$  is odd (that is,  $f(-x) = -f(x)$ ), then

$$\int_{-a}^a f(x) \, dx = \int_{-a}^0 f(x) \, dx + \int_0^a f(x) \, dx = 0.$$



## Example

Since  $f(x) = x^6 + 1$  satisfies  $f(-x) = f(x)$ , it is even, and so

$$\int_{-2}^2 (x^6 + 1) \, dx =$$

## Example

Since  $f(x) = x^6 + 1$  satisfies  $f(-x) = f(x)$ , it is even, and so

$$\int_{-2}^2 (x^6 + 1) \, dx = 2 \int_0^2 (x^6 + 1) \, dx$$

## Example

Since  $f(x) = x^6 + 1$  satisfies  $f(-x) = f(x)$ , it is even, and so

$$\begin{aligned}\int_{-2}^2 (x^6 + 1) \, dx &= 2 \int_0^2 (x^6 + 1) \, dx \\ &= 2 \left[ \frac{1}{7} x^7 + x \right]_0^2\end{aligned}$$

## Example

Since  $f(x) = x^6 + 1$  satisfies  $f(-x) = f(x)$ , it is even, and so

$$\begin{aligned}\int_{-2}^2 (x^6 + 1) \, dx &= 2 \int_0^2 (x^6 + 1) \, dx \\ &= 2 \left[ \frac{1}{7} x^7 + x \right]_0^2 \\ &= 2 \left( \frac{128}{7} + 2 \right)\end{aligned}$$




## Example

Since  $f(x) = x^6 + 1$  satisfies  $f(-x) = f(x)$ , it is even, and so

$$\begin{aligned}\int_{-2}^2 (x^6 + 1) \, dx &= 2 \int_0^2 (x^6 + 1) \, dx \\ &= 2 \left[ \frac{1}{7} x^7 + x \right]_0^2 \\ &= 2 \left( \frac{128}{7} + 2 \right) \\ &= \frac{284}{7}.\end{aligned}$$


## Example



Since  $f(x) = \frac{\tan x - x}{1 - 2x^2 + 2x^4}$  satisfies  $f(-x) = -f(x)$ , it is odd, and so

$$\int_{-1}^1 \frac{\tan x - x}{1 - 2x^2 + 2x^4} dx =$$

## Example




Since  $f(x) = \frac{\tan x - x}{1 - 2x^2 + 2x^4}$  satisfies  $f(-x) = -f(x)$ , it is odd, and so


$$\int_{-1}^1 \frac{\tan x - x}{1 - 2x^2 + 2x^4} dx = 0.$$

# More About Areas


Suppose two curves,  $y = f(x)$  and  $y = g(x)$ , are given. How do we find the area bounded by those curves between the endpoints  $x = a$  and  $x = b$ ?




## The Area Under a Curve



## The Area Between Two Curves




## The Area Under a Curve




rectangle area = height · width

## The Area Between Two Curves




## The Area Under a Curve




rectangle area = height · width

## The Area Between Two Curves




## The Area Under a Curve




rectangle area = height  $\cdot \Delta x$

## The Area Between Two Curves






## The Area Under a Curve




rectangle area = height  $\cdot \Delta x$

## The Area Between Two Curves




## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 4$$

$$A = \sum_{i=1}^4 f(x_i) \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 8$$

$$A = \sum_{i=1}^8 f(x_i) \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 16$$

$$A = \sum_{i=1}^{16} f(x_i) \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




# rectangles =  $n \rightarrow \infty$

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




# rectangles =  $n \rightarrow \infty$

$$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$


## The Area Between Two Curves



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves






## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = \text{height} \cdot \text{width}$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = \text{height} \cdot \text{width}$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = \text{height} \cdot \Delta x$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = \text{height} \cdot \Delta x$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$


$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 4$$

$$A = \sum_{i=1}^4 (f(x_i) - g(x_i)) \Delta x$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves




$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 8$$

$$A = \sum_{i=1}^8 (f(x_i) - g(x_i)) \Delta x$$

## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves



$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$




$$\# \text{ rectangles} = n = 16$$


$$A = \sum_{i=1}^{16} (f(x_i) - g(x_i)) \Delta x$$



## The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b f(x) dx$$

## The Area Between Two Curves



$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b [f(x) - g(x)] dx$$

## Definition (The Area Between Two Curves)


The area between two curves  $y = f(x)$  and  $y = g(x)$  bounded by the endpoints  $x = a$  and  $x = b$  is

$$\int_a^b |f(x) - g(x)| dx.$$

Note that we use the absolute value, because in general we don't know which curve is above the other.


## Example

Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



1 Graph the functions.

## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



1 Graph the functions.

## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



1 Graph the functions.

## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



- 1 Graph the functions.
- 2 Identify the region.

## Example

Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .




- 1 Graph the functions.
- 2 Identify the region.



## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.

## Example

Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .




- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.

$$\begin{aligned} A &= \int_0^1 |(x^2 + 1) - x| dx \\ &= \int_0^1 (x^2 - x + 1) dx \end{aligned}$$

## Example

Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .




- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.

$$\begin{aligned} A &= \int_0^1 |(x^2 + 1) - x| dx \\ &= \int_0^1 (x^2 - x + 1) dx \\ &= \left[ \frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1 \end{aligned}$$

## Example

Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .




- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.

$$\begin{aligned} A &= \int_0^1 |(x^2 + 1) - x| dx \\ &= \int_0^1 (x^2 - x + 1) dx \\ &= \left[ \frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1 \\ &= \frac{1}{3} - \frac{1}{2} + 1 \end{aligned}$$

## Example


Find the area of the region bounded above by  $y = x^2 + 1$ , bounded below by  $y = x$ , and bounded on its sides by  $x = 0$  and  $x = 1$ .



- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.


$$\begin{aligned} A &= \int_0^1 |(x^2 + 1) - x| dx \\ &= \int_0^1 (x^2 - x + 1) dx \\ &= \left[ \frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1 \\ &= \frac{1}{3} - \frac{1}{2} + 1 = \frac{5}{6}. \end{aligned}$$

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

- 1 Find the point of intersection.

## Example




Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$

- 1 Find the point of intersection.



## Example




Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2$$

- 1 Find the point of intersection.

## Example




Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

- 1 Find the point of intersection.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.



## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .


$$x^2 = 2x - x^2$$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$A = \int_0^1 (2x - 2x^2) dx$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$A = \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$\begin{aligned} A &= \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx \\ &= 2 \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 \end{aligned}$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$\begin{aligned} A &= \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx \\ &= 2 \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \left( \frac{1}{2} - \frac{1}{3} \right) \end{aligned}$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the parabolas  $y = x^2$  and  $y = 2x - x^2$ .

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$\begin{aligned} A &= \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx \\ &= 2 \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \left( \frac{1}{2} - \frac{1}{3} \right) = \frac{1}{3}. \end{aligned}$$


- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

- 1 Find the point of intersection.



## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

- 1 Find the point of intersection.

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

- 1 Find the point of intersection.
- 2 Graph the functions.

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.

## Example




Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

$$A = A_1 + A_2$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.

## Example



Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .


The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx + \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

## Example



Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$


$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.



## Example



Find the area of the region enclosed by the curves  $y = \sin x$ ,  $y = \cos x$ ,  $x = 0$  and  $x = \pi/2$ .

The only point of intersection in the interval  $[0, \pi/2]$  is  $(\pi/4, 1/\sqrt{2})$ .

$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

$$= 2\sqrt{2} - 2.$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.