

## Calculus II


Definite integrals of the form  $\int_p^q \sqrt{ax^2 + c} dx,$   
 $a, c > 0$

Todor Milev

2019


## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



## Example


Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



why do we call  
 $y = \sqrt{x^2 + 1}$  hyperbola?

## Example


Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola?

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2\end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1\end{aligned}$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$(y - x)(y + x) = 1$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{1}{2} (y - x) (y + x) = \frac{1}{2}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) = \frac{1}{2}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}.$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:


$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}
 \sqrt{x^2 + 1} &= y \\
 x^2 + 1 &= y^2 \\
 y^2 - x^2 &= 1 \\
 \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\
 uv &= \frac{1}{2} \\
 v &= \frac{1}{2u},
 \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2}$$

We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$

$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) = \frac{1}{2}$$


$$uv = \frac{1}{2}$$

$$v = \frac{1}{2u}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} \end{aligned}$$


We studied  $v = \frac{1}{2}u$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2}u, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} = \pm \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



**Signed** distance b-n  $(x, y)$  and line  $u = 0$  equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) = \\ & \quad \textcolor{red}{u}. \end{aligned}$$

We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:


$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ .


We studied  $v = \frac{1}{2} \frac{y+x}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that **signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .**


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .  
 $\Rightarrow y^2 - x^2 = 1$  is the **hyperbola**  
 $v = \frac{1/2}{u}$  in the  $(u, v)$ -plane.


We studied  $v = \frac{1}{2u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{?}^{?} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned} & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\ &= 2 \left[ x\sqrt{x^2 + 1} \right. \\ & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right)
 \end{aligned}$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right)
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2})
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- **Geometric interpretation of  $\theta$ ?**

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- **Geometric interpretation of  $\theta$ ?**

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- **Geometric interpretation of  $\theta$ ?**

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



- Recall: integral can be solved via  $x = \tan \theta$ .
- **Geometric interpretation of  $\theta$ ?**

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$