

# Calculus I

## Continuity

Todor Milev

2019

# Outline

## 1 Continuity

# Outline

- 1 Continuity
- 2 Intermediate Value Theorem

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.


# Continuity

- Let  $f$  be a function and  $a$  be a point in its domain.



# Continuity

- Let  $f$  be a function and  $a$  be a point in its domain.
- Suppose  $\lim_{x \rightarrow a} f(x)$  exists.




# Continuity

- Let  $f$  be a function and  $a$  be a point in its domain.
- Suppose  $\lim_{x \rightarrow a} f(x)$  exists.

## Definition (Continuous at a Number)

We say that  $f$  is continuous at  $a$  if

$$\lim_{x \rightarrow a} f(x) = f(a).$$



## Definition (Discontinuous at a Number)

Suppose  $f$  is defined at  $a$ . We say  $f$  is discontinuous at  $a$  if it is not continuous at  $a$ .



## Definition (Discontinuous at a Number)

Suppose  $f$  is defined at  $a$ . We say  $f$  is discontinuous at  $a$  if it is not continuous at  $a$ .

Physical phenomena are often continuous. The majority of the physical phenomena that are understood are continuous. Examples:

Discontinuous phenomena examples:

## Definition (Discontinuous at a Number)

Suppose  $f$  is defined at  $a$ . We say  $f$  is discontinuous at  $a$  if it is not continuous at  $a$ .

Physical phenomena are often continuous. The majority of the physical phenomena that are understood are continuous. Examples:

- Motion of a vehicle with respect to time without sudden brakes.
- Orbits of planets and celestial bodies with respect to time.
- A person's height with respect to time.
- And many more.

Discontinuous phenomena examples:

## Definition (Discontinuous at a Number)

Suppose  $f$  is defined at  $a$ . We say  $f$  is discontinuous at  $a$  if it is not continuous at  $a$ .

Physical phenomena are often continuous. The majority of the physical phenomena that are understood are continuous. Examples:


- Motion of a vehicle with respect to time without sudden brakes.
- Orbits of planets and celestial bodies with respect to time.
- A person's height with respect to time.
- And many more.

Discontinuous phenomena examples:

- Particle velocities during collisions and explosions.
- Electric current phenomena, gating events in porins (the event of a molecule passing in and out of a cell).
- Particle physics phenomena.
- And many more.


## Example

The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? Why?




## Example

The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? Why?



## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? Why?



- Not defined at 1:
- Discontinuous at 2:
- Discontinuous at 4:

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$
- $f(1)$
- Discontinuous at 2:
- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example

The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**




- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  ?
- $f(1)$
- Discontinuous at 2:
- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$



## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$
- Discontinuous at 2:
- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  ?
- Discontinuous at 2:
- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- **$f(2)$  ?**
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x) ?$
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- $f(4)$
- $\lim_{x \rightarrow 4} f(x)$

## Example

The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**




- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- **$f(4)$  ?**
- $\lim_{x \rightarrow 4} f(x)$



## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- **$f(4)$  is defined.**
- $\lim_{x \rightarrow 4} f(x)$

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- $f(4)$  is defined.
- $\lim_{x \rightarrow 4} f(x)$  ?

## Example


The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**



- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- $f(4)$  is defined.
- $\lim_{x \rightarrow 4} f(x)$  exists.

## Example

The picture below shows a graph of a function  $f$ . At which numbers is  $f$  either discontinuous or not defined? **Why?**




- Not defined at 1:
- $\lim_{x \rightarrow 1} f(x)$  exists.
- $f(1)$  is not defined.
- Discontinuous at 2:
- $f(2)$  is defined.
- $\lim_{x \rightarrow 2} f(x)$  doesn't exist.
- Discontinuous at 4:
- $f(4)$  is defined.
- $\lim_{x \rightarrow 4} f(x)$  exists.
- $\lim_{x \rightarrow 4} f(x) \neq f(4)$ .

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .


In computer science this function is called the *floor* function.



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = ?$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = ?$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$

$$\left\lfloor -\frac{1}{2} \right\rfloor =$$


$$\lfloor -\pi \rfloor =$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = ?$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = ?$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor = ?$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$

$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$


$$\lfloor -\pi \rfloor = ?$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$

$$\lfloor -\pi \rfloor = -4$$

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

- $f(2)$
- $\lim_{x \rightarrow 2} f(x)$




## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

●  $f(2)$  ?

●  $\lim_{x \rightarrow 2} f(x)$




## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

●  $f(2)$  is defined ( $f(2) = 1$ ).

●  $\lim_{x \rightarrow 2} f(x)$




## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

●  $f(2)$  is defined ( $f(2) = 1$ ).

●  $\lim_{x \rightarrow 2} f(x) ?$




## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$


- $f(2)$  is defined ( $f(2) = 1$ ).
- $\lim_{x \rightarrow 2} f(x)$  exists ( $= 3$ ).



## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




- $f(2)$  is defined ( $f(2) = 1$ ).
- $\lim_{x \rightarrow 2} f(x)$  exists ( $= 3$ ).
- $\lim_{x \rightarrow 2} f(x) \neq f(2)$ .



## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




- $f(2)$  is defined ( $f(2) = 1$ ).
- $\lim_{x \rightarrow 2} f(x)$  exists ( $= 3$ ).
- $\lim_{x \rightarrow 2} f(x) \neq f(2)$ .
- Discontinuous at 2.

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{x^2-x-2}{x-2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




- $f(2)$  is defined ( $f(2) = 1$ ).
- $\lim_{x \rightarrow 2} f(x)$  exists ( $= 3$ ).
- $\lim_{x \rightarrow 2} f(x) \neq f(2)$ .
- Discontinuous at 2.
- This is called a removable discontinuity because we can redefine  $f$  at one point to make  $f$  continuous.

## Example

Where is this function discontinuous?


$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$



## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$
- $\lim_{x \rightarrow 0} f(x)$

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




●  $f(0) ?$

●  $\lim_{x \rightarrow 0} f(x)$

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$  is defined ( $f(0) = 1$ ).
- $\lim_{x \rightarrow 0} f(x)$

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$  is defined ( $f(0) = 1$ ).
- $\lim_{x \rightarrow 0} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$  is defined ( $f(0) = 1$ ).
- $\lim_{x \rightarrow 0} f(x)$  doesn't exist ( $\infty$ ).



## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$  is defined ( $f(0) = 1$ ).
- $\lim_{x \rightarrow 0} f(x)$  doesn't exist ( $\infty$ ).
- Discontinuous at 0.

## Example

Where is this function discontinuous?

$$f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




- $f(0)$  is defined ( $f(0) = 1$ ).
- $\lim_{x \rightarrow 0} f(x)$  doesn't exist ( $\infty$ ).
- Discontinuous at 0.
- This is called an infinite discontinuity.

## Example

Where is this function discontinuous?


$$f(x) = \lfloor x \rfloor$$



## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



•  $f(1)$  ?

•  $\lim_{x \rightarrow 1^+} f(x) = ?$


•  $\lim_{x \rightarrow 1^-} f(x) = ?$

•  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



●  $f(1)$  ?

●  $\lim_{x \rightarrow 1^+} f(x) = ?$ .


●  $\lim_{x \rightarrow 1^-} f(x) = ?$ .

●  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



●  $f(1)$  exists ( $f(1) = 1$ ).

●  $\lim_{x \rightarrow 1^+} f(x) = ?$ .


●  $\lim_{x \rightarrow 1^-} f(x) = ?$ .

●  $\lim_{x \rightarrow 1} f(x) = ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



●  $f(1)$  exists ( $f(1) = 1$ ).

●  $\lim_{x \rightarrow 1^+} f(x) = ?$ .


●  $\lim_{x \rightarrow 1^-} f(x) = ?$ .

●  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



●  $f(1)$  exists ( $f(1) = 1$ ).

●  $\lim_{x \rightarrow 1^+} f(x) = 1$ .

●  $\lim_{x \rightarrow 1^-} f(x) = ?$ .


●  $\lim_{x \rightarrow 1} f(x) ?$



## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



•  $f(1)$  exists ( $f(1) = 1$ ).

•  $\lim_{x \rightarrow 1^+} f(x) = 1$ .


•  $\lim_{x \rightarrow 1^-} f(x) = ?$ .

•  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



•  $f(1)$  exists ( $f(1) = 1$ ).

•  $\lim_{x \rightarrow 1^+} f(x) = 1$ .


•  $\lim_{x \rightarrow 1^-} f(x) = 0$ .

•  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



•  $f(1)$  exists ( $f(1) = 1$ ).

•  $\lim_{x \rightarrow 1^+} f(x) = 1$ .


•  $\lim_{x \rightarrow 1^-} f(x) = 0$ .

•  $\lim_{x \rightarrow 1} f(x) ?$

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$




- $f(1)$  exists ( $f(1) = 1$ ).
- $\lim_{x \rightarrow 1^+} f(x) = 1$ .
- $\lim_{x \rightarrow 1^-} f(x) = 0$ .
- $\lim_{x \rightarrow 1} f(x)$  doesn't exist.

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$




- $f(1)$  exists ( $f(1) = 1$ ).
- $\lim_{x \rightarrow 1^+} f(x) = 1$ .
- $\lim_{x \rightarrow 1^-} f(x) = 0$ .
- $\lim_{x \rightarrow 1} f(x)$  doesn't exist.
- Discontinuous at 1.

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$




- $f(1)$  exists ( $f(1) = 1$ ).
- $\lim_{x \rightarrow 1^+} f(x) = 1$ .
- $\lim_{x \rightarrow 1^-} f(x) = 0$ .
- $\lim_{x \rightarrow 1} f(x)$  doesn't exist.
- Discontinuous at 1.
- Discontinuous at every integer  $n$ .

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$




- $f(1)$  exists ( $f(1) = 1$ ).
- $\lim_{x \rightarrow 1^+} f(x) = 1$ .
- $\lim_{x \rightarrow 1^-} f(x) = 0$ .
- $\lim_{x \rightarrow 1} f(x)$  doesn't exist.
- Discontinuous at 1.
- Discontinuous at every integer  $n$ .
- The left and right limits both exist but are not equal.

## Example

Where is this function discontinuous?

$$f(x) = \lfloor x \rfloor$$



- $f(1)$  exists ( $f(1) = 1$ ).
- $\lim_{x \rightarrow 1^+} f(x) = 1$ .
- $\lim_{x \rightarrow 1^-} f(x) = 0$ .
- $\lim_{x \rightarrow 1} f(x)$  doesn't exist.
- Discontinuous at 1.
- Discontinuous at every integer  $n$ .
- The left and right limits both exist but are not equal.
- Such discontinuities are called jump discontinuities (the function appears to “jump”).



## Definition (Continuous from the Right or Left)

A function  $f$  is continuous from the right at a number  $a$  if


$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

and  $f$  is continuous from the left at  $a$  if

$$\lim_{x \rightarrow a^-} f(x) = f(a).$$


## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



## Example


Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



- $f(n) = \quad$ .
- $\lim_{x \rightarrow n^+} f(x) = ?$ .
- $\lim_{x \rightarrow n^-} f(x) = ?$ .

## Example


Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



- $f(n) = ?$ .
- $\lim_{x \rightarrow n^+} f(x) = ?$ .
- $\lim_{x \rightarrow n^-} f(x) = ?$ .

## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .




●  $f(n) = n.$

●  $\lim_{x \rightarrow n^+} f(x) = ?.$

●  $\lim_{x \rightarrow n^-} f(x) = ?.$

## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .




●  $f(n) = n.$

●  $\lim_{x \rightarrow n^+} f(x) = ?.$

●  $\lim_{x \rightarrow n^-} f(x) = ?.$

## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .




- $f(n) = n.$

- $\lim_{x \rightarrow n^+} f(x) = n.$

- $\lim_{x \rightarrow n^-} f(x) = ?$

## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .




- $f(n) = n$ .
- $\lim_{x \rightarrow n^+} f(x) = n$ .
- Continuous from the right at  $n$ .
- $\lim_{x \rightarrow n^-} f(x) = ?$ .



## Example


Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



- $f(n) = n$ .
- $\lim_{x \rightarrow n^+} f(x) = n$ .
- Continuous from the right at  $n$ .
- $\lim_{x \rightarrow n^-} f(x) = ?$ .

## Example


Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



- $f(n) = n$ .
- $\lim_{x \rightarrow n^+} f(x) = n$ .
- Continuous from the right at  $n$ .
- $\lim_{x \rightarrow n^-} f(x) = n - 1$ .

## Example

Consider  $f(x) = \lfloor x \rfloor$ , and pick any integer  $n$ .



- $f(n) = n$ .
- $\lim_{x \rightarrow n^+} f(x) = n$ .
- Continuous from the right at  $n$ .
- $\lim_{x \rightarrow n^-} f(x) = n - 1$ .
- Discontinuous from the left at  $n$ .

## Definition (Continuous on an Interval)

A function  $f$  is continuous on an interval if it is continuous at every number in the interval.

## Definition (Continuous on an Interval)

A function  $f$  is continuous on an interval if it is continuous at every number in the interval.

- If  $f$  is defined at the **right** endpoint of an interval, continuous means continuous from the **left**.

## Definition (Continuous on an Interval)

A function  $f$  is continuous on an interval if it is continuous at every number in the interval.

- If  $f$  is defined at the **left** endpoint of an interval, continuous means continuous from the **right**.

## Definition (Continuous on an Interval)

A function  $f$  is continuous on an interval if it is continuous at every number in the interval.

- If  $f$  is defined at the left endpoint of an interval, continuous means continuous from the right.
- Think of a function that is continuous on an interval as a function that has no breaks in its graph, and so can be drawn “without lifting your pen”.

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

①  $f + g$

③  $cf$

⑤  $\frac{f}{g}$  if  $g(a) \neq 0$ .

②  $f - g$

④  $fg$

Proof.



## Theorem (Algebra of Continuous Functions)

If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :

1  $f + g$

3  $cf$

5  $\frac{f}{g}$  if  $g(a) \neq 0$ .

2  $f - g$

4  $fg$

## Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

1  $f + g$

3  $cf$

5  $\frac{f}{g}$  if  $g(a) \neq 0$ .

2  $f - g$

4  $fg$

### Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\lim_{x \rightarrow a} (f + g)(x)$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

①  $f + g$

③  $cf$

⑤  $\frac{f}{g}$  if  $g(a) \neq 0$ .

②  $f - g$

④  $fg$

## Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\lim_{x \rightarrow a} (f + g)(x) = \lim_{x \rightarrow a} [f(x) + g(x)]$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

1  $f + g$

3  $cf$

5  $\frac{f}{g}$  if  $g(a) \neq 0$ .

2  $f - g$

4  $fg$

### Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \end{aligned}$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

①  $f + g$

③  $cf$

⑤  $\frac{f}{g}$  if  $g(a) \neq 0$ .

②  $f - g$

④  $fg$

## Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \\ &= f(a) + g(a) \end{aligned}$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

$$① \quad f + g$$

$$③ \quad cf$$

$$⑤ \quad \frac{f}{g} \text{ if } g(a) \neq 0.$$

$$② \quad f - g$$

$$④ \quad fg$$

## Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \\ &= f(a) + g(a) \end{aligned}$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

1  $f + g$

3  $cf$

5  $\frac{f}{g}$  if  $g(a) \neq 0$ .

2  $f - g$

4  $fg$

## Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \\ &= f(a) + g(a) = (f + g)(a) \end{aligned}$$

## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

1  $f + g$

3  $cf$

5  $\frac{f}{g}$  if  $g(a) \neq 0$ .

2  $f - g$

4  $fg$

### Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \\ &= f(a) + g(a) = (f + g)(a) \end{aligned}$$

This shows  $f + g$  is continuous at  $a$ .



## Theorem (Algebra of Continuous Functions)

*If  $f$  and  $g$  are continuous at  $a$  and  $c$  is a constant, then the following functions are also continuous at  $a$ :*

①  $f + g$

③  $cf$

⑤  $\frac{f}{g}$  if  $g(a) \neq 0$ .

②  $f - g$

④  $fg$

### Proof.

$$\lim_{x \rightarrow a} f(x) = f(a) \text{ and } \lim_{x \rightarrow a} g(x) = g(a).$$

$$\begin{aligned} \lim_{x \rightarrow a} (f + g)(x) &= \lim_{x \rightarrow a} [f(x) + g(x)] \\ &= \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) \quad (\text{by Law 1}) \\ &= f(a) + g(a) = (f + g)(a) \end{aligned}$$

This shows  $f + g$  is continuous at  $a$ . The other parts are similar.  $\square$

## Theorem (Classes of Continuous Functions)

*The following types of functions are continuous at every number in their domains:*

*polynomials      rational functions*  
*root functions      trigonometric functions*

## Theorem (Compositions of Continuous Functions)

*If  $g$  is continuous at  $a$  and  $f$  is continuous at  $g(a)$ , then the composition function  $f \circ g$  given by  $(f \circ g)(x) = f(g(x))$  is continuous at  $a$ .*

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain.

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq ?$ .

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$$

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} = \lim_{x \rightarrow -2} f(x)$$



## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\begin{aligned} \lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} &= \lim_{x \rightarrow -2} f(x) \\ &= f(-2) \end{aligned}$$

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\begin{aligned} \lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} &= \lim_{x \rightarrow -2} f(x) \\ &= f(-2) \\ &= \frac{(-2)^3 + 2(-2)^2 - 1}{5 - 3(-2)} \end{aligned}$$

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\begin{aligned} \lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} &= \lim_{x \rightarrow -2} f(x) \\ &= f(-2) \\ &= \frac{(-2)^3 + 2(-2)^2 - 1}{5 - 3(-2)} \\ &= ? \end{aligned}$$

## Example

Find  $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ .

The function  $f(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$  is rational, so is continuous on its domain. Its domain is given by  $x \neq \frac{5}{3}$ .

$$\begin{aligned} \lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} &= \lim_{x \rightarrow -2} f(x) \\ &= f(-2) \\ &= \frac{(-2)^3 + 2(-2)^2 - 1}{5 - 3(-2)} \\ &= -\frac{1}{11} \end{aligned}$$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) =$
- $h(u) =$
- $g(v) =$
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = ?$
- $h(u) =$
- $g(v) =$
- $f(w) =$



## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) =$
- $g(v) =$
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = ?$
- $g(v) =$
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) =$
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = ?$
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) =$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = ?$

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = \frac{1}{w}$ .

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = \frac{1}{w}$ .
- These functions are continuous on their domains, so  $F$  is continuous on its domain.



## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = \frac{1}{w}$ .
- These functions are continuous on their domains, so  $F$  is continuous on its domain.
- Its domain is ?

## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = \frac{1}{w}$ .
- These functions are continuous on their domains, so  $F$  is continuous on its domain.
- Its domain is **given by  $x \neq 3$  and  $x \neq -3$** .


## Example

Where is the function  $F(x) = \frac{1}{\sqrt{x^2+7}-4}$  continuous?

- We can write  $F$  as the composition of 4 functions:
- $F = f \circ g \circ h \circ k$ , or  $F(x) = f(g(h(k(x))))$ .
- $k(x) = x^2 + 7$ .
- $h(u) = \sqrt{u}$ .
- $g(v) = v - 4$ .
- $f(w) = \frac{1}{w}$ .
- These functions are continuous on their domains, so  $F$  is continuous on its domain.
- Its domain is given by  $x \neq 3$  and  $x \neq -3$ .
- Therefore  $F$  is continuous on  $(-\infty, -3) \cup (-3, 3) \cup (3, \infty)$ .


## Theorem (The Intermediate Value Theorem)

*Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ , where  $f(a) \neq f(b)$ . Then there exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$ .*




## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and *let  $N$  be any number between  $f(a)$  and  $f(b)$* , where  $f(a) \neq f(b)$ . Then there exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$ .




## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ , where  $f(a) \neq f(b)$ . Then there **exists a number  $c$**  in  $(a, b)$  **such that  $f(c) = N$** .




## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and *let  $N$  be any number between  $f(a)$  and  $f(b)$* , where  $f(a) \neq f(b)$ . Then there exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$ .



## Theorem (The Intermediate Value Theorem)


Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ , where  $f(a) \neq f(b)$ . Then there **exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$** .






## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and *let  $N$  be any number between  $f(a)$  and  $f(b)$* , where  $f(a) \neq f(b)$ . Then there exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$ .




## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ , where  $f(a) \neq f(b)$ . Then there **exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$** .




## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and *let  $N$  be any number between  $f(a)$  and  $f(b)$* , where  $f(a) \neq f(b)$ . Then there exists a number  $c$  in  $(a, b)$  such that  $f(c) = N$ .



## Theorem (The Intermediate Value Theorem)

Suppose  $f$  is continuous on the closed interval  $[a, b]$  and let  $N$  be any number between  $f(a)$  and  $f(b)$ , where  $f(a) \neq f(b)$ . Then there **exists a number  $c$**  in  $(a, b)$  **such that  $f(c) = N$** .




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .






## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .
- $f(1) = -1$ .




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .
- $f(1) = -1$ .
- $f(2) = 12$ .




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .
- $f(1) = -1$ .
- $f(2) = 12$ .
- $f(1) < 0 < f(2)$ .




## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .
- $f(1) = -1$ .
- $f(2) = 12$ .
- $f(1) < 0 < f(2)$ .
- Therefore there is a  $c$  between 1 and 2 such that  $f(c) = 0$ .



## Example

Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.

- Let  $f(x) = 4x^3 - 6x^2 + 3x - 2$ .
- $f$  is continuous.
- Use the IVT with  $a = 1$ ,  $b = 2$ , and  $N = 0$ .
- $f(1) = -1$ .
- $f(2) = 12$ .
- $f(1) < 0 < f(2)$ .
- Therefore there is a  $c$  between 1 and 2 such that  $f(c) = 0$ .

