

Calculus II

Tangents and curve length

Todor Milev

2019

Outline

- 1 Tangents to Curves
 - Tangents to Polar Curves

- 2 Arc Length
 - Arc Length in Polar Coordinates

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Tangents

Let C be the curve $C : \begin{cases} x = f(t) \\ y = g(t) \end{cases}, t \in [a, b]$.

Definition

Suppose $f'(t)$ and $g'(t)$ are not simultaneously equal to 0.


- We define $(f'(t), g'(t))$ to be the *tangent vector* to C at t .
- We define the line passing through $(f(t), g(t))$ with direction vector equal to the tangent vector to be *tangent line* to C at t . In other words, the tangent line has equation

$$(x - f(t))g'(t) = (y - g(t))f'(t) \quad .$$

- We say that the tangent to C at t is vertical if $f'(t) = 0$ (and therefore $g'(t) \neq 0$).

Note. When $f'(t) = g'(t) = 0$, for curves C with additional properties, natural definition(s) of tangent(s) do exist but are beyond Calc II.

Example



Find the tangent to the curve

$$\gamma : \begin{cases} x = \cos t \\ y = \sin t \end{cases}, t \in [0, 2\pi) \text{ at } t = \frac{\pi}{4}, t = \frac{2\pi}{3}, t = \pi.$$

Recall C : $\begin{cases} x = x(t) \\ y = y(t) \end{cases}, t \in [a, b]$, tangent vector at t is $(x'(t), y'(t))$.

We write informally $x = x(t), y = y(t)$ to simplify notation.


- Suppose we could eliminate the parameter t and write $y = F(x)$ for some function F near the point $(x, y) = (x(t), y(t))$.
- Suppose in $x'(t) \neq 0$ for some t .

$$\begin{array}{lcl}
 y & = & F(x) \\
 \frac{dy}{dt} & = & \frac{d}{dt}(F(x)) \\
 & = & \frac{dF}{dx} \frac{dx}{dt} = \frac{dy}{dx} \frac{dx}{dt} \\
 \frac{dy}{dx} & = & \frac{\frac{dy}{dt}}{\frac{dx}{dt}}
 \end{array}
 \quad \left| \begin{array}{l} \text{apply } \frac{d}{dt} \\ \text{use chain rule} \\ \text{divide by } x'(t) \end{array} \right.$$

Observation

If $\frac{dx}{dt} \neq 0$, we have $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$.


Example



A curve C is defined by $x = t^2$, $y = t^3 - 3t$.

- 1 Show C traverses $(x, y) = (3, 0)$ for two values of t ; find the tangent slopes for both of these values.
- 2 Find the points on C where the tangents are horizontal or vertical.
- 3 Find two intervals where we can write y as a function of x .
- 4 Determine concavity intervals of the functions found in item 3.


Example



A curve C is defined by $x = t^2$, $y = t^3 - 3t$.

- ① Show C traverses $(x, y) = (3, 0)$ for two values of t ; find the tangent slopes for both of these values.
 - $3 = x = t^2$ if $t = \pm\sqrt{3}$.
 - $0 = y = t^3 - 3t = t(t^2 - 3)$ if $t = 0$ or $\pm\sqrt{3}$.
 - Therefore the point $(3, 0)$ is traversed when t equals $\sqrt{3}$ or $-\sqrt{3}$.
 - $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3t^2 - 3}{2t}$.
 - Plug in $t = \pm\sqrt{3}$: $\frac{dy}{dx} \Big|_{t=\pm\sqrt{3}} = \frac{3(\pm\sqrt{3})^2 - 3}{2(\pm\sqrt{3})} = \pm \frac{6}{2\sqrt{3}} = \pm\sqrt{3}$
- Therefore the tangents at $(3, 0)$ have slopes $\pm\sqrt{3}$.

Example



A curve C is defined by $x = t^2, y = t^3 - 3t$.

- ② Find the points on C where the tangents are horizontal or vertical.

Horizontal tangent:

$$\frac{dy}{dt} = 0$$

$$3t^2 - 3 = 0$$

$$3(t^2 - 1) = 0$$

$$t = \pm 1$$

$\frac{dx}{dt} \neq 0$ when $t = \pm 1$, so there are horizontal tangents when $t = \pm 1$.

The points are $(1, 2)$ and $(1, -2)$.

Vertical tangent:

$$\frac{dx}{dt} = 0$$


$$2t = 0$$

$$t = 0$$

$\frac{dy}{dt} \neq 0$ when $t = 0$, so there is a vertical tangent when $t = 0$.

The point is $(0, 0)$.

Example




A curve C is defined by $x = t^2$, $y = t^3 - 3t$.

③ Find two intervals where we can write y as a function of x .

From $x = t^2$ we have that $t = \pm\sqrt{x}$. Therefore, when $t > 0$, we have that $t = \sqrt{x}$. Since that determines uniquely t via x , this means that for $t > 0$ y is a function of x . In other words, for $t > 0$, the curve satisfies the vertical line test. Similarly we conclude that when $t < 0$, y is a function of x .

Example



A curve C is defined by $x = t^2, y = t^3 - 3t$.

- 4 Determine the concavity intervals of the functions found in item 3.


Find the second derivative:

$$\begin{aligned}
 \frac{d^2y}{dx^2} &= \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}} = \frac{\frac{d}{dt} \left(\frac{3t^2-3}{2t} \right)}{2t} \\
 &= \frac{\frac{d}{dt} \left(\frac{3}{2} \left(t - \frac{1}{t} \right) \right)}{2t} = \frac{\frac{3}{2} + \frac{3}{2t^2}}{2t} \\
 &= \frac{\frac{3t^2+3}{2t^2}}{2t} = \frac{3(t^2+1)}{4t^3}
 \end{aligned}$$

Therefore y as a function of x (which is a function of t) is concave up when $t > 0$ and concave down when $t < 0$.

Example


Consider the cycloid $x = r(\theta - \sin \theta)$, $y = r(1 - \cos \theta)$.



- 1 At what points is the tangent horizontal?
- 2 At what points is the tangent vertical?

Example

Consider the cycloid $x = r(\theta - \sin \theta)$, $y = r(1 - \cos \theta)$.




1 At what points is the tangent horizontal?

- The slope of the tangent is $\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{r \sin \theta}{r(1 - \cos \theta)} = \frac{\sin \theta}{1 - \cos \theta}$
- The tangent is horizontal when $dy/dx = 0$, that is, when $dy/d\theta = 0$ and $dx/d\theta \neq 0$.
- $r \sin \theta = dy/d\theta = 0$ if $\theta = n\pi$, where n is any integer.
- $r(1 - \cos \theta) = dx/d\theta = 0$ if $\theta = 2n\pi$, where n is any integer.
- Therefore there is a horizontal tangent when $\theta = (2n + 1)\pi$.

Example

Consider the cycloid $x = r(\theta - \sin \theta)$, $y = r(1 - \cos \theta)$.



2 At what points is the tangent vertical?

- When $\theta = 2n\pi$ both $dy/d\theta$ and $dx/d\theta$ are 0.
- To see if there is a vertical tangent, use L'Hospital's Rule.

$$\lim_{\theta \rightarrow 2n\pi^+} \frac{dy}{dx} = \lim_{\theta \rightarrow 2n\pi^+} \frac{\sin \theta}{1 - \cos \theta} = \lim_{\theta \rightarrow 2n\pi^+} \frac{\cos \theta}{\sin \theta} \rightarrow \frac{1}{0^+}$$

- Therefore $\lim_{\theta \rightarrow 2n\pi^+} (dy/dx) = \infty$.
- A similar argument shows $\lim_{\theta \rightarrow 2n\pi^-} (dy/dx) = -\infty$.
- Therefore there is a vertical tangent when $\theta = 2n\pi$.

Tangents to Polar Curves

To find the tangent line to a polar curve $r = f(\theta)$, regard θ as a parameter and write the parametric equations as


$$x = r \cos \theta = f(\theta) \cos \theta \quad y = r \sin \theta = f(\theta) \sin \theta$$

Then use the formula for the slope of a parametric curve:

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} \\ &= \frac{\frac{d}{d\theta} (f(\theta) \sin \theta)}{\frac{d}{d\theta} (f(\theta) \cos \theta)} \\ &= \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta + f(\theta)(-\sin \theta)} \\ &= \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} \end{aligned}$$

Example

Find the points on $r = 1 + \sin \theta$ where the tangent is horizontal or vertical.




$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta} = \frac{\cos \theta \sin \theta + (1 + \sin \theta) \cos \theta}{\cos \theta \cos \theta - (1 + \sin \theta) \sin \theta} \\ &= \frac{\cos \theta (1 + 2 \sin \theta)}{1 - 2 \sin^2 \theta - \sin \theta} = \frac{\cos \theta (1 + 2 \sin \theta)}{(1 + \sin \theta)(1 - 2 \sin \theta)}\end{aligned}$$

- $\cos \theta (1 + 2 \sin \theta) = 0$
when $\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{6}, \frac{11\pi}{6}$.
- $(1 + \sin \theta)(1 - 2 \sin \theta) = 0$
when $\theta = \frac{3\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}$.

- Horizontal tangents at $(2, \pi/2)$, $(1/2, 7\pi/6)$, and $(1/2, 11\pi/6)$.
- Vertical tangents at $(3/2, \pi/6)$, and $(3/2, 5\pi/6)$.
- If $\theta = 3\pi/2$, top and bottom are both 0, so use L'Hospital's Rule.

$$\lim_{\theta \rightarrow 3\pi/2^-} \frac{dy}{dx} = \lim_{\theta \rightarrow 3\pi/2^-} \frac{1 + 2 \sin \theta}{1 - 2 \sin \theta} \cdot \lim_{\theta \rightarrow 3\pi/2^-} \frac{\cos \theta}{1 + \sin \theta} = -\frac{1}{3} \lim_{\theta \rightarrow 3\pi/2^-} \frac{-\sin \theta}{\cos \theta} = \infty$$


Arc Length



- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to ∞ .

Let γ be the curve $\gamma : \begin{cases} x = x(t) \\ y = y(t) \end{cases}, t \in [a, b]$

- Divide $[a, b]$ into n subintervals with endpoints t_0, t_1, \dots, t_n and equal width Δt .
- The points $P_i = (x(t_i), y(t_i))$ lie on the curve γ . The lengths of the segments with endpoints with consecutive indices from P_0, P_1, \dots, P_n approximate the length of the curve γ .
- The length L of the curve γ is the limit of the lengths of these segments as $n \rightarrow \infty$.



$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1} P_i|$$

Let γ be the curve $\gamma : \begin{cases} x = x(t) \\ y = y(t) \end{cases}, t \in [a, b]$

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{(x'(s_i))^2 + (y'(r_i))^2} \Delta t \\ &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt \end{aligned}$$

- If f has continuous derivative, we can compute the above limit.
- Let $\begin{cases} x_i = x(t_i) \\ y_i = y(t_i) \end{cases}$, and $\begin{cases} \Delta x = x_i - x_{i-1} = x(t_i) - x(t_{i-1}) \\ \Delta y = y_i - y_{i-1} = y(t_i) - y(t_{i-1}) \end{cases}$.
- Then $|P_iP_{i-1}| = \sqrt{(\Delta x)^2 + (\Delta y)^2}$.
- Mean Value Theorem: there exist numbers s_i and r_i between t_{i-1} and t_i such that $x(t_i) - x(t_{i-1}) = x'(s_i)(t_i - t_{i-1})$ and $y(t_i) - y(t_{i-1}) = y'(r_i)(t_i - t_{i-1})$.
- $\Delta x = x'(s_i)\Delta t$, $\Delta y = y'(r_i)\Delta t$.

$$\begin{aligned} |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(x'(s_i)\Delta t)^2 + (y'(r_i)\Delta t)^2} \\ &= \sqrt{(x'(s_i))^2 + (y'(r_i))^2} \sqrt{(\Delta t)^2} = \sqrt{(x'(s_i))^2 + (y'(r_i))^2} \Delta t \end{aligned}$$

The Arc Length Formula

Let $\gamma : \begin{cases} x = x(t) \\ y = y(t) \end{cases}, t \in [a, b]$.

Definition

Suppose $x'(t)$ and $y'(t)$ (exist and) are continuous on $[a, b]$. Then the length of the curve γ is defined as

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt \\ &= \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \quad \text{in Leibniz notation .} \end{aligned}$$

Arc length of graph of a function

Question

What is the length of the graph of the curve given by the graph of $y = f(x)$?

- The graph of $y = f(x)$ is written as a curve as

$$\gamma : \begin{cases} x = t \\ y = f(t) \end{cases}, t \in [a, b] .$$

- In other words, the question asks what is the length $L(\gamma)$ of γ . That is a straightforward computation:

$$L(\gamma) = \int \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int \sqrt{1 + (f'(t))^2} dt$$

The Arc Length Formula


Definition

Suppose f' exists and is continuous on $[a, b]$. Then the length of the curve $y = f(x)$, $a \leq x \leq b$, is

$$\begin{aligned} L &= \int_a^b \sqrt{1 + (f'(x))^2} \, dx \\ &= \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx \quad (\text{in Leibniz notation}) \end{aligned}$$

Example


Find the length of the arc of $y^2 = x^3$ between $(1, 1)$ and $(4, 8)$.




- For the top half of the curve we have:
- $y = x^{3/2}$ and $y' = \frac{3}{2}x^{1/2}$.
- $u = 1 + \frac{9}{4}x$ and $du = \frac{9}{4}dx$.
- When $x = 1$, $u = \frac{13}{4}$.
- When $x = 4$, $u = 10$.

$$\begin{aligned}
 L &= \int_1^4 \sqrt{1 + (y')^2} dx \\
 &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4}^{10} \frac{4}{9} \sqrt{u} du \\
 &= \frac{4}{9} \left[\frac{2}{3} u^{3/2} \right]_{13/4}^{10} = \frac{8}{27} \left(10^{3/2} - \left(\frac{13}{4} \right)^{3/2} \right)
 \end{aligned}$$

Example




Find the length of the arc of the parabola $y = x^2$ from $(0,0)$ to $(1,1)$.



$$\begin{aligned}
 L &= \int_{x=0}^{x=1} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_{x=0}^{x=1} \sqrt{1 + 4x^2} dx & \left| \text{Set } x = \frac{1}{2} \tan \theta \right. \\
 &= \int_{\theta=0}^{\theta=\arctan 2} \sqrt{1 + \tan^2 \theta} d\left(\frac{1}{2} \tan \theta\right) \\
 &= \int_{\theta=0}^{\theta=\arctan 2} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^{\alpha} \sec^3 \theta d\theta & \left| \text{Set } \alpha = \arctan 2 \right. \\
 &= \frac{1}{2} \cdot \left[\frac{1}{2} (\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|) \right]_{\theta=0}^{\theta=\alpha} \\
 &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \\
 &= \frac{1}{4} (2 \cdot \sqrt{5} + \ln |\sqrt{5} + 2|)
 \end{aligned}$$

Example



Find the length of the curve γ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2 \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that $x'(t) = \frac{1}{2\sqrt{t}} - 2$ and $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$.

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[\sqrt{t} + 2t\right]_1^4 = \sqrt{4} + 2 \cdot 4 - (\sqrt{1} + 2 \cdot 1) = 7. \end{aligned}$$

Example $((a + b)^2, (a - b)^2, 2ab = 1/2)$



Find the length of the arc of $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$ from $x = 0$ to $x = 1$.

$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$\begin{aligned}(y')^2 &= \frac{1}{4}e^{6x} - \frac{1}{4}e^{3x}e^{-3x} - \frac{1}{4}e^{3x}e^{-3x} + \frac{1}{4}e^{-6x} \\ &= \frac{1}{4}e^{6x} - \frac{1}{2} + \frac{1}{4}e^{-6x}.\end{aligned}$$

$$\begin{aligned}L &= \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{1 + \frac{1}{4}e^{6x} - \frac{1}{2} + \frac{1}{4}e^{-6x}} dx \\ &= \int_0^1 \sqrt{\frac{1}{4}e^{6x} + \frac{1}{2} + \frac{1}{4}e^{-6x}} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx \\ &= \int_0^1 \left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right) dx = \left[\frac{1}{6}e^{3x} - \frac{1}{6}e^{-3x}\right]_0^1 = \frac{e^3 - e^{-3}}{6}.\end{aligned}$$

Example



Find the length of one arch of the cycloid

$$x = r(\theta - \sin \theta), \quad y = r(1 - \cos \theta).$$

The first arch is $0 \leq \theta \leq 2\pi$.

$$\begin{aligned} L &= \int_0^{2\pi} \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta = \int_0^{2\pi} \sqrt{(r(1 - \cos \theta))^2 + (r \sin \theta)^2} d\theta \\ &= \int_0^{2\pi} \sqrt{r^2(1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta)} d\theta = r \int_0^{2\pi} \sqrt{2(1 - \cos \theta)} d\theta \end{aligned}$$

Use the identity $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$. Then

$$\sqrt{2(1 - \cos \theta)} = \sqrt{4 \sin^2(\theta/2)} = 2|\sin(\theta/2)| = 2 \sin(\theta/2)$$

$$L = r \int_0^{2\pi} 2 \sin(\theta/2) d\theta = r [-4 \cos(\theta/2)]_0^{2\pi} = 8r$$

Arc Length

To find the arc length of a polar curve $r = f(\theta)$, $a \leq \theta \leq b$, regard θ as a parameter. Then the derivatives of the parametric equations are

$$\frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta \quad \frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta$$


and

$$\begin{aligned} \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 &= \left(\frac{dr}{d\theta}\right)^2 \cos^2 \theta - 2r \frac{dr}{d\theta} \cos \theta \sin \theta + r^2 \sin^2 \theta \\ &\quad + \left(\frac{dr}{d\theta}\right)^2 \sin^2 \theta + 2r \frac{dr}{d\theta} \sin \theta \cos \theta + r^2 \cos^2 \theta \\ &= \left(\frac{dr}{d\theta}\right)^2 + r^2 \end{aligned}$$

The arc length is

$$L = \int_a^b \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta = \int_a^b \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Example



Find the length of the cardioid $r = 1 + \sin \theta$. The full length is given by $0 \leq \theta \leq 2\pi$.

$$\begin{aligned}
 L &= \int_0^{2\pi} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta = \int_0^{2\pi} \sqrt{(1 + \sin \theta)^2 + \cos^2 \theta} d\theta \\
 &= \int_0^{2\pi} \sqrt{2 + 2 \sin \theta} \frac{\sqrt{2 - 2 \sin \theta}}{\sqrt{2 - 2 \sin \theta}} d\theta = \int_0^{2\pi} \frac{\sqrt{4 - 4 \sin^2 \theta}}{\sqrt{2 - 2 \sin \theta}} d\theta \\
 &= \int_0^{2\pi} \frac{\sqrt{4 \cos^2 \theta}}{\sqrt{2 - 2 \sin \theta}} d\theta = \int_0^{2\pi} \frac{2|\cos \theta|}{\sqrt{2 - 2 \sin \theta}} d\theta \\
 &= \int_0^{\pi/2} \frac{2 \cos \theta}{\sqrt{2 - 2 \sin \theta}} d\theta + \int_{\pi/2}^{3\pi/2} \frac{-2 \cos \theta}{\sqrt{2 - 2 \sin \theta}} d\theta + \int_{3\pi/2}^{2\pi} \frac{2 \cos \theta}{\sqrt{2 - 2 \sin \theta}} d\theta \\
 &= \left[-2\sqrt{2 - 2 \sin \theta}\right]_0^{\pi/2} + \left[2\sqrt{2 - 2 \sin \theta}\right]_{\pi/2}^{3\pi/2} + \left[-2\sqrt{2 - 2 \sin \theta}\right]_{3\pi/2}^{2\pi} \\
 &= -2(0 - \sqrt{2}) + 2(2 - 0) - 2(\sqrt{2} - 2) = 8
 \end{aligned}$$