

Calculus II

Weierstrass substitution theory

Todor Milev

2019

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - Apply the substitution $\theta = 2 \arctan t$ to transform to integral of rational function.

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - Apply the substitution $\theta = 2 \arctan t$ to transform to integral of rational function.
 - Solve as previously studied.

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = ?$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z}{(\cos^2 z + \sin^2 z)}$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}}$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z) = ?$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad .$$

$$\cos(2z) = \cos^2 z - \sin^2 z$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad . \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z)}{(\cos^2 z + \sin^2 z)}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad . \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$.

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta, \cos \theta$?

$$\sin \theta =$$

$$\cos \theta =$$

Recall the expression of $\sin(2z), \cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t)$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)}$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2}$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2}$$

$$\cos \theta = \cos(2 \arctan t)$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)}\end{aligned}$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta, \cos \theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

$d\theta$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t)\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = ? \quad dt\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= ?\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= \tan\left(\frac{\theta}{2}\right)\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= \tan\left(\frac{\theta}{2}\right)\end{aligned}$$

Theorem

The substitution given above transforms $\int R(\cos \theta, \sin \theta) d\theta$ to an integral of a rational function of t .

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\int \sec \theta d\theta$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$.

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{\cancel{1+t^2}} dt \\ &= \int \frac{2}{1-t^2} dt \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \\ &= \ln \left| \frac{1+t}{1-t} \right| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\begin{aligned}
 \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\
 &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\
 &= -\ln |1-t| + \ln |1+t| + C \\
 &= \ln \left| \frac{1+t}{1-t} \right| + C \\
 &= \ln \left| \frac{1 + \tan\left(\frac{\theta}{2}\right)}{1 - \tan\left(\frac{\theta}{2}\right)} \right| + C
 \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \\ &= \ln \left| \frac{1+t}{1-t} \right| + C \\ &= \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta =$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2}))} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})) (\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2}))} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln |\tan \theta + \sec \theta| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$