
### Calculus I


# Reference: tangents to graphs of functions

**Todor Milev** 


2019




• How to find the tangent line to the curve y = f(x) at P = (a, f(a))?




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.



- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.

Let P = (a, f(a)). Suppose the limit  $m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$  exists. Define the tangent to y = f(x) at P to be the line passing through P with slope m




- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.

Let P = (a, f(a)). Suppose the limit  $m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$  exists. Define the tangent to y = f(x) at P to be the line passing through P with slope m, in other words, the line with equation



- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.

Let P = (a, f(a)). Suppose the limit  $m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$  exists. Define the tangent to y = f(x) at P to be the line passing through P with slope m, in other words, the line with equation y - f(a) = m(x - a).



- How to find the tangent line to the curve y = f(x) at P = (a, f(a))?
- Consider nearby point Q = (x, f(x)).
- Compute slope of secant line *PQ*:  $m_{PQ} = \frac{f(x) f(a)}{x a}$ .
- As x approaches a, the point Q approaches P.

Let P = (a, f(a)). Suppose the limit  $m = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$  exists. Define the tangent to y = f(x) at P to be the line passing through P with slope m, in other words, the line with equation y - f(a) = m(x - a).

**Note.** Even if the limit does not exist a reasonable notion of a tangent line may still exist.


### **Derivatives**

#### Definition (Derivative)

The derivative of a function f at a number a, denoted by f'(a), is

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h}$$

if the limit exists.




#### **Derivatives**

### Definition (Derivative)

The derivative of a function f at a number a, denoted by f'(a), is

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h} = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

if the limit exists.



 The two alternative formulas result in equivalent definitions.


### **Derivatives**

### Definition (Derivative)

The derivative of a function f at a number a, denoted by f'(a), is

$$f'(a) = \lim_{h \to 0} \frac{f(a+h) - f(a)}{h} = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

if the limit exists.



- The two alternative formulas result in equivalent definitions.
  - Equivalent formulation. The derivative f'(a) is the slope of the tangent line to y = f(x) at (a, f(a)), provided that tangent line exists and is non-vertical.