

Precalculus

Degree lowering formulas

Todor Milev

2019

Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned}\cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ \color{red}{2}\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{\color{red}{2}}\end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha - 1 = \cos(2\alpha)$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned} \quad \begin{aligned} \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\cos^2 \alpha - 1 &= \cos(2\alpha) \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha - 1 = \cos(2\alpha)$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

Corollary (Half-Angle Formulas)

$$\sin\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$

Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

Corollary (Half-Angle Formulas)

$$\sin\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$

Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express $\sin^k \alpha$ and $\cos^k \alpha$ via lower powers of the sin and cos functions (applied to angles other than α).

Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express $\sin^k \alpha$ and $\cos^k \alpha$ via lower powers of the sin and cos functions (applied to angles other than α).
- This technique will play a key role in integration (studied later/in another course).

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\sin^4 \alpha$$

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\sin^4 \alpha = \left(\sin^2 \alpha \right)^2$$

Recall the formulas: $\sin^2 \beta = ?$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned} \sin^4 \alpha &= \left(\sin^2 \alpha \right)^2 \\ &= \left(? \right)^2 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned} \sin^4 \alpha &= \left(\sin^2 \alpha \right)^2 \\ &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} \left(? \right) \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} (?) \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha))\end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = ?$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + ?)
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \cos^2(2\alpha) \right) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right)
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left(\frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right)
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left(\frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\
 &= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))
 \end{aligned}$$

Recall the formulas: $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$, $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$.

Example

Rewrite $\sin^4 \alpha$ in terms of first powers of the cosines and sines of multiples of the angle α .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left(\frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left(1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left(\frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\
 &= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))
 \end{aligned}$$