Calculus II Integrals of the form $\int \sqrt{ax^2 + bx + c} dx$, quadratic has no real roots.

Todor Miley

2019

Linear substitutions to simplify radicals $\sqrt{ay^2 + by + c}$

- Using linear substitutions, radicals of form $\sqrt{ay^2 + by + c}$, $a \neq 0$, $b^2 4ac \neq 0$ can be transformed to (multiple of):
 - $\sqrt{x^2 + 1}$
 - $\sqrt{-x^2+1}$
 - $\sqrt{x^2-1}$.
- We already studied how to do that using completing the square when dealing with rational functions.

Example

$$\sqrt{x^2 + x + 1} =$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + ? - ? + 1}$$

Example

$$\sqrt{x^2+x+1} = \sqrt{x^2+2\cdot\frac{1}{2}x+\frac{1}{4}-\frac{1}{4}+1}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

$$= \sqrt{\left(x + ?\right)^2 + ?}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$
$$= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

$$= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$= \sqrt{\frac{3}{4} \left(\frac{4}{3} \left(x + \frac{1}{2}\right)^2 + 1\right)}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

$$= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$= \sqrt{\frac{3}{4}\left(\frac{4}{3}\left(x + \frac{1}{2}\right)^2 + 1\right)}$$

$$= \frac{\sqrt{3}}{2}\sqrt{\left(\frac{2}{\sqrt{3}}\left(x + \frac{1}{2}\right)\right)^2 + 1}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

$$= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$= \sqrt{\frac{3}{4}\left(\frac{4}{3}\left(x + \frac{1}{2}\right)^2 + 1\right)}$$

$$= \frac{\sqrt{3}}{2}\sqrt{\left(\frac{2}{\sqrt{3}}\left(x + \frac{1}{2}\right)\right)^2 + 1}$$

Example

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

$$= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$= \sqrt{\frac{3}{4}\left(\frac{4}{3}\left(x + \frac{1}{2}\right)^2 + 1\right)}$$

$$= \frac{\sqrt{3}}{2}\sqrt{\left(\frac{2}{\sqrt{3}}\left(x + \frac{1}{2}\right)^2 + 1\right)}$$

$$= \frac{\sqrt{3}}{2}\sqrt{u^2 + 1},$$
where $u = \frac{2}{\sqrt{3}}\left(x + \frac{1}{2}\right) = \frac{2\sqrt{3}}{3}x + \frac{\sqrt{3}}{3}.$

where
$$u = \frac{2}{\sqrt{3}} \left(x + \frac{1}{2} \right) = \frac{2\sqrt{3}}{3} x + \frac{\sqrt{3}}{3}$$
.