

Precalculus

Equations involving logarithms and exponents

Todor Milev

2019

Outline

- 1 Equations involving logarithms
- 2 Equations involving exponents
- 3 Inverse function problems and exponents
- 4 Basic exponential inequalities

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Example

Solve the equation.

$$\begin{aligned}\log_3(2x^2 + 1) &= 2 && | \text{Exponentiate base 3} \\ 3^{\log_3(2x^2 + 1)} &= 3^2 \\ 2x^2 + 1 &= 9 \\ 2x^2 &= 8 \\ x^2 &= \frac{8}{2} = 4 \\ x &= \pm\sqrt{4} = \pm 2 \\ x = 2 \text{ or } x = -2 && | \text{final answer}\end{aligned}$$

The logarithmic property $\log_a(xy) = \log_a x + \log_a y$ holds only for positive x, y . Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

Example

Solve the equation.

$$\log_2(x+2) + \log_2(x-1) = 2$$

$$\log_2((x+2)(x-1)) = 2$$

$$(x+2)(x-1) = 2^2$$

$$x^2 + x - 2 = 4$$

$$x^2 + x - 6 = 0$$

$$(x-2)(x+3) = 0$$

$$x = 2 \quad \text{or} \quad x = -3$$

$x = -3$ not a solution (outside of domain)

Domain: $x > 1$

Exponentiate base 2

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned}16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \\ 2^{16t} &= 2^{3t-6} \\ 16t &= 3t - 6 \\ 13t &= -6 \\ t &= -\frac{6}{13}.\end{aligned}$$

Example

Solve the equation.

$$\begin{aligned}2^{1-5x} &= 12 && | \text{ apply } \log_2 \\ \log_2(2^{1-5x}) &= \log_2 12 \\ 1 - 5x &= \log_2 12 = \log_2(4 \cdot 3) \\ 1 - 5x &= \log_2 4 + \log_2 3 \\ 1 - 5x &= 2 + \log_2 3 \\ 5x &= 1 - (2 + \log_2 3) \\ &= -1 - \log_2 3 \\ x &= \frac{-1 - \log_2 3}{5} \\ \text{Calculator: } x &\approx -0.516993.\end{aligned}$$

Example

Solve the equation.

$$e^{x-3} = 2e^{2x-1}$$

Divide by e^{2x-1}

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

$$e^{-x-2} = 2$$

Apply \ln

$$-x - 2 = \ln 2$$

$$-x = \ln 2 + 2$$

$$x = -(\ln 2 + 2)$$

$$x = -\ln 2 - 2$$

Final answer

$$x \approx -2.693$$

Calculator

Example

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

Common base

$$(2^{\log_2 3})^{2x+5} = 5 \cdot 2^{-x+1}$$

$$a = b^{\log_b a}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

Apply \log_2

$$(\log_2 3)(2x + 5) + x - 1 = \log_2 5$$

$$x(2 \log_2 3 + 1) + 5 \log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1 - 5 \log_2 3}{2 \log_2 3 + 1}$$

$$x \approx -1.1038$$

Calculator

Example

Solve the equation.

$$e^{5-3x} = 10 \quad \text{apply } \ln$$

$$=$$

$$=$$

$$x =$$

Calculator: $x \approx 0.8991.$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $=$

 $=$ $=$ $=$ $=$ $=$ $t =$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\u^2 - 2u - 63 &= 0 \\(u - 9)(u + 7) &= 0\end{aligned}$$

$$\begin{aligned}u &= 9 \quad \text{or} \quad u = -7 \\3^x &= 9 \quad \text{or} \quad 3^x = -7 \\x &= 2 & \quad \text{no real solution}\end{aligned}$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

$$e^x = 4$$

or

$$e^x = -1$$

$$x = \ln 4$$

or

no real solution

$$x \approx 1.3863$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right) = \frac{\ln \left(\frac{3}{2} \right)}{\ln 2} \approx 0.58496 \quad \text{or} \quad \text{no real solution}$$

Example (Exponential equation that reduces to quadratic)

Solve the equation.

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^2 - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \quad \text{or} \quad u + 7 = 0$$

$$u = 9 \quad \text{or} \quad u = -7$$

$$3^{2x} = 9 \quad \text{or} \quad \text{no real solution}$$

$$2x = \log_3 9$$

$$2x = 2$$

$$x = 1$$

> 2 terms \Rightarrow

transfer one side

$$3^{2x} = u$$

$$3^{-2x} = (3^{2x})^{-1} = u^{-1}$$


Multiply $\cdot u$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$f = \tanh =$ hyperbolic tangent function.



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln \left(\frac{1 + y}{1 - y} \right)$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Take \ln

Example

Solve the inequality.

$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$-3x - 5 < \log_2 7$$

$$-3x < \log_2 7 + 5$$

$$x > -\frac{\log_2 7 + 5}{3}$$

$$x \in \left(-\frac{5 + \log_2 7}{3}, \infty \right)$$

Logarithms preserve inequalities: apply \log_2

Division by negative number flips inequalities

