

# Calculus II

## Trigonometry review

Todor Milev

2019

# Outline

- 1 Review of trigonometry
  - The Trigonometric Functions
  - Trigonometric Identities
  - Trigonometric Identities and Complex Numbers
  - Graphs of the Trigonometric Functions

# Outline

- 1 Review of trigonometry
  - The Trigonometric Functions
  - Trigonometric Identities
  - Trigonometric Identities and Complex Numbers
  - Graphs of the Trigonometric Functions
- 2 Inverse Trigonometric Functions

# License to use and redistribute


These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.


- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Geometric interpretation of all trigonometric functions




Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .

# Geometric interpretation of all trigonometric functions




Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ .

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ .

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$\sin \theta$

$\cos \theta$

$\tan \theta$


$\cot \theta$

$\sec \theta$

$\csc \theta$



# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$\sin \theta$

$\cos \theta$


$\tan \theta$

$\cot \theta$

$\sec \theta$

$\csc \theta$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$

$$\tan \theta$$


$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$



# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ . Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ . Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$


$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$



# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

# Geometric interpretation of all trigonometric functions



$\beta = ?$

$\angle OED = ?$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$\sec \theta$

$\csc \theta$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$


$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$



# Geometric interpretation of all trigonometric functions



$\angle OED = ?$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
 Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$



# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ . Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|}$$

$$\sec \theta$$

$$\csc \theta$$



# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|} = \frac{1}{|DE|}$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{opp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1}$$

$\csc \theta$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$



# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OB|}{|AC|}$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|} = \frac{1}{|DE|} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1}$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ . Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

# Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center  $O$ , coordinates  $(0, 0)$ .  
Let  $\angle DOB = \theta$ . Let  $OB$  intersect the circle at point  $A$ . Coordinates of  $A$  are  $(\cos \theta, \sin \theta)$ .

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$



# Trigonometric Identities

## Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

# Trigonometric Identities

## Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.


- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.

# Trigonometric Identities

## Definition (Trigonometric Identity)


A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.
- For example,  $\frac{\sin \theta}{\sin \theta} = 1$  is considered a valid trigonometric identity, although, when considered as a function, the left hand side is not defined for  $\theta \neq 0$ .




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- $\csc \theta = \frac{1}{\sin \theta}$
- $\sec \theta = \frac{1}{\cos \theta}$
- $\cot \theta = \frac{1}{\tan \theta}$
- $\tan \theta = \frac{\sin \theta}{\cos \theta}$
- $\cot \theta = \frac{\cos \theta}{\sin \theta}$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\sin^2 \theta + \cos^2 \theta$$

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2}\end{aligned}$$






$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$

Therefore  $\sin^2 \theta + \cos^2 \theta = 1$ .




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

### Example ( $\tan^2 \theta + 1 = \sec^2 \theta$ )

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

### Example ( $\tan^2 \theta + 1 = \sec^2 \theta$ )

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\sin^2 \theta + \cos^2 \theta = 1$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

### Example ( $\tan^2 \theta + 1 = \sec^2 \theta$ )

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

### Example ( $\tan^2 \theta + 1 = \sec^2 \theta$ )

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta} \\ \tan^2 \theta + 1 &= \sec^2 \theta\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$



The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

Substitute  $-y$  for  $y$ , and use the fact that  $\sin(-y) = -\sin y$  and  $\cos(-y) = \cos y$ :

$$\begin{aligned}\sin(x - y) &= \sin x \cos y - \cos x \sin y \\ \cos(x - y) &= \cos x \cos y + \sin x \sin y\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute  $x$  for  $y$ :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute  $x$  for  $y$ :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using  $\cos^2 x = 1 - \sin^2 x$  and  $\sin^2 x = 1 - \cos^2 x$ :

$$\cos(2x) = 2 \cos^2 x - 1$$

$$\cos(2x) = 1 - 2 \sin^2 x$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute  $x$  for  $y$ :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using  $\cos^2 x = 1 - \sin^2 x$  and  $\sin^2 x = 1 - \cos^2 x$ :

$$\cos(2x) = 2 \cos^2 x - 1$$

$$\cos(2x) = 1 - 2 \sin^2 x$$

To get the half-angle formulas, solve these equations for  $\cos^2 x$  and  $\sin^2 x$  respectively.

$$\cos^2 x = \frac{1 + \cos(2x)}{2}, \quad \sin^2 x = \frac{1 - \cos(2x)}{2}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

Divide the first equation by the second, and then cancel  $\cos x \cos y$  from the top and bottom:

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$


Divide the first equation by the second, and then cancel  $\cos x \cos y$  from the top and bottom:

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

Do the same for the subtraction formulas:


$$\tan(x - y) = \frac{\tan x - \tan y}{1 + \tan x \tan y}$$






- Positive angles are obtained by rotating counterclockwise.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$ .

$$\begin{aligned} \sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y} \end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$ .
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$ .

$$\begin{aligned} \sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y} \end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$ .
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$ .
- $\sin$  is an **odd function**.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If  $(x, y)$  is on the terminal arm of the angle  $\theta$ , then  $(x, -y)$  is on the terminal arm of  $-\theta$ .
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$ .
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$ .
- $\sin$  is an odd function.
- $\cos$  is an even function.






$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- $2\pi$  represents a full rotation.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- $2\pi$  represents a full rotation.
- $\theta + 2\pi$  has the same terminal arm as  $\theta$ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- $2\pi$  represents a full rotation.
- $\theta + 2\pi$  has the same terminal arm as  $\theta$ .
- $\theta + 2\pi$  uses the same point  $(x, y)$  and the same length  $r$ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- $2\pi$  represents a full rotation.
- $\theta + 2\pi$  has the same terminal arm as  $\theta$ .
- $\theta + 2\pi$  uses the same point  $(x, y)$  and the same length  $r$ .
- $\sin(\theta + 2\pi) = \sin \theta$ .
- $\cos(\theta + 2\pi) = \cos \theta$ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- $2\pi$  represents a full rotation.
- $\theta + 2\pi$  has the same terminal arm as  $\theta$ .
- $\theta + 2\pi$  uses the same point  $(x, y)$  and the same length  $r$ .
- $\sin(\theta + 2\pi) = \sin \theta$ .
- $\cos(\theta + 2\pi) = \cos \theta$ .
- We say  $\sin$  and  $\cos$  are  $2\pi$ -periodic.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit.

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1.$$

The number  $i$  is called the imaginary unit.

$$\pm \sqrt{-1} = i.$$



## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. **By definition,**  $\pm \sqrt{-1} = i$ .

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. **By definition,**  $\sqrt{-1} = i$ .

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$



## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2 = ac + adi + bci - bd$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = \textcolor{red}{ac} + adi + bci - \textcolor{red}{bd} \\ &= (\textcolor{red}{ac} - \textcolor{red}{bd}) + i(ad + bc)\end{aligned}$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + ad\textcolor{red}{i} + bc\textcolor{red}{i} - bd \\ &= (ac - bd) + \textcolor{red}{i}(ad + bc)\end{aligned}$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- **Complex addition/subtraction**

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- **Complex multiplication**

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + \textcolor{red}{ad}i + \textcolor{red}{bc}i - bd \\ &= (ac - bd) + i(\textcolor{red}{ad} + \textcolor{red}{bc})\end{aligned}$$

## Definition (Complex numbers)

The set of complex numbers  $\mathbb{C}$  is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number  $i$  is a number for which

$$i^2 = -1 \quad .$$

The number  $i$  is called the imaginary unit. By definition,  $\sqrt{-1} = i$ .

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + adi + bci - bd \\ &= (ac - bd) + i(ad + bc)\end{aligned}$$

# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant .

## Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

## Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + \frac{(-1)^n x^{2n+1}}{(2n+1)!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + \frac{(-1)^n x^{2n}}{(2n)!} + \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

## Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} + \frac{z^5}{5!} + \dots$$

Rearrange.



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

## Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix + \frac{(ix)^2}{2!} + \frac{(ix)^3}{3!} + \frac{(ix)^4}{4!} + \frac{(ix)^5}{5!} + \dots$$

Rearrange. **Plug-in**  $z = ix$ .



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

## Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in  $z = ix$ . Use  $i^2 = -1$ .



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

### Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in  $z = ix$ . Use  $i^2 = -1$ . Multiply  $\sin x$  by  $i$ .



# Euler's Formula

## Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where  $e \approx 2.71828$  is Euler's/Napier's constant.

### Proof.

Recall  $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ . Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in  $z = ix$ . Use  $i^2 = -1$ . Multiply  $\sin x$  by  $i$ . **Add to get**  
 $e^{ix} = \cos x + i \sin x.$



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

All trigonometric formulas can be easily derived using the above formulas.

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$e^{i(x+y)} =$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$e^{i(x+y)} = \cos(x+y) + i \sin(x+y)$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x$ ,  $\cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Compare coefficient in front of  $i$  and

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Compare coefficient in front of  $i$  and **remaining terms** to get the desired equalities.



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$1 = e^0$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = \end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = \end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\ &= (\cos x + i \sin x)(\cos x - i \sin x) = \end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\ &= (\cos x + i \sin x)(\cos x - i \sin x) = \cos^2 x - i^2 \sin^2 x \end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\sin^2 x + \cos^2 x = 1$$

## Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\ &= (\cos x + i \sin x)(\cos x - i \sin x) = \cos^2 x - i^2 \sin^2 x \\ &= \cos^2 x + \sin^2 x . \end{aligned}$$



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.



# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$e^{i(2x)} =$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$e^{i(2x)} = \cos(2x) + i \sin(2x)$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)(\cos x + i \sin x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Compare coefficient in front of  $i$  and

# Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$  (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$  (exponentiation rule: valid for  $\mathbb{C}$ ).
- $e^0 = 1$  (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$  (easy to remember).

## Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

## Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Compare coefficient in front of  $i$  and **remaining terms** to get the desired equalities. □

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .



- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) \\ &= e^{3ix} \end{aligned}$$

| Euler's f-la

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) \\ &= e^{3ix} \end{aligned}$$

| Euler's f-la

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) \\ &= e^{3ix} \\ &= (e^{ix})^3 \end{aligned}$$

| Euler's f-la

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) \\ &= e^{3ix} \\ &= (e^{ix})^3 \end{aligned} \quad \Bigg| \quad \text{Euler's f-la}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\cos(3x) + i \sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3 = (\cos x + i \sin x)^3$$

| Euler's f-la

| Euler's f-la

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = ?$

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\ &= e^{3ix} \\ &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\ &= e^{3ix} \\ &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\ &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3
 \end{aligned}$$



- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\ &= e^{3ix} \\ &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\ &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(\textcolor{red}{i} \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3\textcolor{red}{i}^2 \cos x \sin^2 x + i^3 \sin^3 x
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned} & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\ &= e^{3ix} \\ &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\ &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\ &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3 \cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3 \cos x \sin^2 x) + i(3 \cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$



- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example

Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$

The real parts of the starting and final expression must be equal;  
therefore:

$$\cos(3x) = \cos^3 x - 3\cos x \sin^2 x$$

- Recall Euler's formula:  $e^{i\alpha} = \cos \alpha + i \sin \alpha$ .
- Recall the formula:  $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ .

## Example


Express  $\sin(3x)$  and  $\cos(3x)$  via  $\cos x$  and  $\sin x$ .

$$\begin{aligned}
 & \cos(3x) + i \sin(3x) && | \text{ Euler's f-la} \\
 &= e^{3ix} \\
 &= (e^{ix})^3 = (\cos x + i \sin x)^3 && | \text{ Euler's f-la} \\
 &= \cos^3 x + 3\cos^2 x(i \sin x) + 3\cos x(i \sin x)^2 + (i \sin x)^3 \\
 &= \cos^3 x + 3i \cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x \\
 &= \cos^3 x + 3i \cos^2 x \sin x - 3\cos x \sin^2 x - i \sin^3 x && | \text{ Use } i^2 = -1 \\
 &= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)
 \end{aligned}$$

The real parts of the starting and final expression must be equal; likewise the imaginary parts must be equal; therefore:

$$\begin{aligned}
 \cos(3x) &= \cos^3 x - 3\cos x \sin^2 x \\
 \sin(3x) &= 3\cos^2 x \sin x - \sin^3 x
 \end{aligned}$$


# Graphs of the Trigonometric Functions



$$y = \sin x$$

$$y = \cos x$$

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .



# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .
- If we translate the graph of  $\cos x$  by  $\frac{\pi}{2}$  units to the right

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .
- If we translate the graph of  $\cos x$  by  $\frac{\pi}{2}$  units to the right

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .
- If we translate the graph of  $\cos x$  by  $\frac{\pi}{2}$  units to the right

# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .
- If we translate the graph of  $\cos x$  by  $\frac{\pi}{2}$  units to the right we get the graph of  $\sin x$ .


# Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$


- $\sin x$  has zeroes at  $n\pi$  for all integers  $n$ .
- $\cos x$  has zeroes at  $\frac{\pi}{2} + n\pi$  for all integers  $n$ .
- $-1 \leq \sin x \leq 1$ .
- $-1 \leq \cos x \leq 1$ .
- If we translate the graph of  $\cos x$  by  $\frac{\pi}{2}$  units to the right we get the graph of  $\sin x$ . This is a consequence of  $\cos\left(x - \frac{\pi}{2}\right) = \sin x$ .




$$y = \tan x$$



$$y = \cot x$$




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the  $x$  axis






$$y = \tan x$$




$$y = \cot x$$

If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the  $x$  axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the  $x$  axis, we get the graph of  $\cot x$ .




$$y = \tan x$$




$$y = \cot x$$


If we move the graph of  $\tan x$  by  $\frac{\pi}{2}$  units to the left (or right) and reflect across the  $x$  axis, we get the graph of  $\cot x$ . This follows from  $\tan\left(x \pm \frac{\pi}{2}\right) = -\cot x$ .




$$y = \csc x$$




$$y = \sec x$$




$$y = \csc x$$



$$y = \sec x$$




$$y = \csc x$$




$$y = \sec x$$



# Inverse Trigonometric Functions




# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.

# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .

# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .


# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .



# Inverse Trigonometric Functions



- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .
- $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .



## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .



## Observation

- $\arcsin y =$  *the appropriate angle* whose sine equals  $y$ .

## Example

Find  $\arcsin \left( \frac{1}{2} \right)$ .

- $\sin \left( ? \right) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin(?) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin \left( \frac{1}{2} \right)$ .

- $\sin \left( ? \right) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\textcolor{red}{?}\right) = \frac{1}{2}.$

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*
- *Important: the output angle must lie in the interval  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$ .
- $-\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*
- *Important: the output angle must lie in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .*


## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$ .
- $-\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}$ .
- Therefore  $\arcsin\left(\frac{1}{2}\right) = \frac{\pi}{6}$ .

## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .






## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with **opposite side 1** and hypotenuse 3.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and **hypotenuse 3**.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side = ?



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .

- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side =  $\sqrt{3^2 - 1^2}$






## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .
- Then  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right) = ?$



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .

- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .
- Then  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right) = \frac{1}{2\sqrt{2}}$ .



## Example

Find  $\arcsin(\sin(1.5))$ .

## Example

Find  $\arcsin(\sin(1.5))$ .

●  $\frac{\pi}{2} \approx ?$

## Example

Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .

## Example

Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .
- Therefore  $-\frac{\pi}{2} \leq 1.5 \leq \frac{\pi}{2}$ .

## Example



Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .
- Therefore  $-\frac{\pi}{2} \leq 1.5 \leq \frac{\pi}{2}$ .
- Therefore  $\arcsin(\sin 1.5) = 1.5$ .



## Example



Find  $\arcsin(\sin 2)$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  **$a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$**  for which  $\sin 2 = \sin a$ .







## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = ?$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\text{Therefore } \arcsin(\sin 2) = \arcsin(\sin a)$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  **$a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$**  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\begin{aligned} \text{Therefore } \arcsin(\sin 2) &= \arcsin(\sin a) \\ &= a \end{aligned}$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\begin{aligned} \text{Therefore } \arcsin(\sin 2) &= \arcsin(\sin a) \\ &= a = \pi - 2. \end{aligned}$$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

Proof.



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\quad = ?$





## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = ?$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = ?$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = 1$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = 1$

$$y' = \frac{1}{\cos y}$$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\cos y} \\ &= \frac{1}{\pm \sqrt{1 - \sin^2 y}} \end{aligned}$$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .

Differentiate implicitly:  $\cos y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\cos y} \\ &= \frac{1}{\pm \sqrt{1 - \sin^2 y}} \end{aligned}$$

But  $\cos y > 0$ : 
$$= \frac{1}{\sqrt{1 - \sin^2 y}}$$



## Theorem (The Derivative of $\arcsin x$ )

$$\frac{d}{dx} (\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

### Proof.

Let  $y = \arcsin x$ .

Then  $\sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .


Differentiate implicitly:  $\cos y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\cos y} \\ &= \frac{1}{\pm \sqrt{1 - \sin^2 y}} \end{aligned}$$

$$\text{But } \cos y > 0: \quad = \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}.$$




## Important facts about arcsin:



- 1 Domain: ?
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .




## Important facts about arcsin:




- 1 Domain: ?
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 **Range: ?**
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .


## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .




- Same for  $\cos x$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called  $\arccos$  or  $\cos^{-1}$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called  $\arccos$  or  $\cos^{-1}$ .








- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called  $\arccos$  or  $\cos^{-1}$ .
- $\arccos(x) = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .




## Important facts about arccos:




- 1 Domain:
- 2 Range:
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:




- 1 Domain: ?
- 2 Range:
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:




- 1 Domain:  $[-1, 1]$ .
- 2 Range:
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:




- 1 Domain:  $[-1, 1]$ .
- 2 Range: ?
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:



- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[0, \pi]$ .
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:



- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[0, \pi]$ .
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .  
(The proof is similar to the proof of the formula for the derivative of  $\arcsin x$ .)

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .

$$\sin(2 \arccos(x))$$



## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ .

$$\sin(2 \arccos(x))$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\sin(2 \arccos(x))$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\sin(2 \arccos(x)) = \sin(2y)$$

$$| \text{ Set } y = \arccos x$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= ?\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to **rewrite the expression only using the cos function**.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right)\end{aligned} \quad \left| \begin{array}{l} \text{Set } y = \arccos x \\ \text{Express via } \sin y, \cos y \\ \text{Express } \sin y \text{ via } \cos y \end{array} \right.$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y}\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$   
Express  $\sin y$  via  $\cos y$   
 $\sin y > 0$  because  
 $0 \leq y \leq \pi$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y}\end{aligned}$$

Set  $y = \arccos x$

Express via  $\sin y, \cos y$

Express  $\sin y$  via  $\cos y$

$\sin y > 0$  because

$0 \leq y \leq \pi$



## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y} \\ &= 2x \sqrt{1 - x^2}\end{aligned}$$

Set  $y = \arccos x$

Express via  $\sin y, \cos y$

Express  $\sin y$  via  $\cos y$

$\sin y > 0$  because

$0 \leq y \leq \pi$

use  $x = \cos y$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y} \\ &= 2x \sqrt{1 - x^2}\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$   
Express  $\sin y$  via  $\cos y$   
 $\sin y > 0$  because  
 $0 \leq y \leq \pi$   
use  $x = \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .

$$\cos(3 \arccos(x))$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ .

$$\cos(3 \arccos(x))$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x))$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) \quad \Big| \quad y = \arccos x$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) = \cos(2y + y) \quad \bigg| \quad y = \arccos x$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) = \cos(2y + y) \\ = ?$$

$y = \arccos x$   
Angle sum f-la



## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\ &= \cos(2y) \cos y - \sin(2y) \sin y\end{aligned} \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\text{?}) \cos y \\
 &\quad - \text{?} \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - \sin y \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - \sin y \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\ &= \cos(2y) \cos y - \sin(2y) \sin y \\ &= (\cos^2 y - \sin^2 y) \cos y \\ &\quad - 2 \sin y \cos y \sin y\end{aligned}$$

$y = \arccos x$   
Angle sum f-la  
Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$



## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(\text{?}) \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y \\
 &= 4\cos^3 y - 3 \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y \\
 &= 4\cos^3 y - 3\cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y \\
 &= 4\cos^3 y - 3\cos y \\
 &= 4x^3 - 3x
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

$x = \cos y$


## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ .


To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) & y &= \arccos x \\
 &= \cos(2y) \cos y - \sin(2y) \sin y & \text{Angle sum f-la} \\
 &= (\cos^2 y - \sin^2 y) \cos y & \text{Express via} \\
 &\quad - 2 \sin y \cos y \sin y & \sin y, \cos y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y & \text{Express } \sin y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y & \text{via } \cos y \\
 &= 4\cos^3 y - 3 \cos y \\
 &= 4x^3 - 3x & x = \cos y
 \end{aligned}$$


- $\tan x$  isn't one-to-one.









- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- **Domain of  $\arctan$ : ?**
- Range of  $\arctan$ :




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- **Domain of  $\arctan$ :  $(-\infty, \infty)$ .**
- Range of  $\arctan$ :



- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ : ?




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = ?$
- $\lim_{x \rightarrow -\infty} \arctan x =$






- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x =$




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x = ?$



- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x = -\frac{\pi}{2}$ .

## Example


Simplify the expression  $\cos(\arctan x)$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .
- Length of hypotenuse = ?



## Example

Simplify the expression  $\cos(\arctan x)$ .

- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent 1.
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .






## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent 1.
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .
- Then  $\cos(\arctan x) = ?$



## Example

Simplify the expression  $\cos(\arctan x)$ .

- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent 1.
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .
- Then  $\cos(\arctan x) = \frac{1}{\sqrt{1+x^2}}$ .



## Example

Evaluate

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right).$$

## Example

Evaluate

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right).$$

$$\frac{1}{x-2} \rightarrow \infty \quad \text{as} \quad x \rightarrow 2^+.$$

## Example

Evaluate

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right).$$

$$\frac{1}{x-2} \rightarrow \infty \quad \text{as} \quad x \rightarrow 2^+.$$

Therefore

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right) = ?$$

## Example

Evaluate

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right).$$

$$\frac{1}{x-2} \rightarrow \infty \quad \text{as} \quad x \rightarrow 2^+.$$

Therefore

$$\lim_{x \rightarrow 2^+} \arctan \left( \frac{1}{x-2} \right) = \frac{\pi}{2}.$$

## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

Proof.



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

## Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .





## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\quad = ?$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = ?$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = ?$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

## Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = 1$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = 1$

$$y' = \frac{1}{\sec^2 y}$$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\sec^2 y} \\ &= \frac{1}{?} \end{aligned}$$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\sec^2 y} \\ &= \frac{1}{1 + \tan^2 y} \end{aligned}$$



## Theorem (The Derivative of $\arctan x$ )

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}.$$

### Proof.

Let  $y = \arctan x$ .

Then  $\tan y = x$ .

Differentiate implicitly:  $\sec^2 y \cdot y' = 1$

$$\begin{aligned} y' &= \frac{1}{\sec^2 y} \\ &= \frac{1}{1 + \tan^2 y} \\ &= \frac{1}{1 + x^2}. \end{aligned}$$





The remaining inverse trigonometric functions aren't used as often:


$$\begin{aligned}y = \operatorname{arccsc} x \quad (|x| \geq 1) &\Leftrightarrow \csc y = x \quad \text{and} \quad y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right] \\y = \operatorname{arcsec} x \quad (|x| \geq 1) &\Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\y = \operatorname{arccot} x \quad (|x| \in \mathbb{R}) &\Leftrightarrow \cot y = x \quad \text{and} \quad y \in (0, \pi)\end{aligned}$$

The remaining inverse trigonometric functions aren't used as often:

$$\begin{aligned}y &= \operatorname{arccsc} x \quad (|x| \geq 1) \quad \Leftrightarrow \quad \csc y = x \quad \text{and} \quad y \in (0, \frac{\pi}{2}] \cup (\pi, \frac{3\pi}{2}] \\y &= \operatorname{arcsec} x \quad (|x| \geq 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{and} \quad y \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2}) \\y &= \operatorname{arccot} x \quad (|x| \in \mathbb{R}) \quad \Leftrightarrow \quad \cot y = x \quad \text{and} \quad y \in (0, \pi)\end{aligned}$$

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$


- Plot  $\sec x$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$

- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$




- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .
- $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

Table of derivatives of inverse trigonometric functions:

$$\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\operatorname{arccsc} x) = -\frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx}(\operatorname{arcsec} x) = \frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx}(\operatorname{arccot} x) = -\frac{1}{1+x^2}$$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = ?$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .



## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

$$= \begin{pmatrix} ? & \end{pmatrix} \begin{pmatrix} \end{pmatrix}$$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

$$= \left( -u^{-2} \right) \left( \quad \right)$$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

$$= \left( -u^{-2} \right) \left( ? \right)$$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

$$= \left( -u^{-2} \right) \left( \frac{1}{\sqrt{1-x^2}} \right)$$

## Example (Chain Rule)

Differentiate  $y = \frac{1}{\arcsin x}$ .

Let  $u = \arcsin x$ .

Then  $y = u^{-1}$ .

Chain Rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

$$= \left( -u^{-2} \right) \left( \frac{1}{\sqrt{1-x^2}} \right)$$
$$= - \frac{1}{(\arcsin x)^2 \sqrt{1-x^2}}.$$

All of the inverse trigonometric derivatives also give rise to integration formulas. These two are the most important:

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C.$$

$$\int \frac{1}{x^2+1} dx = \arctan x + C.$$