

## Precalculus

# Polynomial division and factorization of cubics with rational root

Todor Milev

2019

# Outline

## 1 Polynomial division

# Outline

- 1 Polynomial division
- 2 Factoring cubics with rational root

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$x - 1 \overline{) x^3 + 2x^2 \quad + 1}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \textcolor{red}{?} \\
 \textcolor{red}{x} - 1 \overline{) \textcolor{red}{x}^3 + 2x^2 \phantom{+ 0x} + 1}
 \end{array}$$

Divide  $\textcolor{red}{x}^3$  by  $\textcolor{red}{x}$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$x - 1 \overline{) \overset{x^2}{x^3 + 2x^2} + 1}$$

Divide  $x^3$  by  $x$ .



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1} \\
 \underline{\phantom{x^3} ? \phantom{+ 0x} ?} \phantom{+ 1} \\
 \phantom{x^3 + 2x^2} \phantom{+ 0x} \phantom{+ 1}
 \end{array}$$

Multiply  $x^2$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1} \\
 \underline{x^3 - x^2} \phantom{+ 0x} \\
 \phantom{x^3} 3x^2 \phantom{+ 0x} + 1
 \end{array}$$

Multiply  $x^2$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 \phantom{x^3} 3x^2 \phantom{+ 1} \\
 \phantom{x^3} \underline{3x^2 - 3x} \phantom{+ 1} \\
 \phantom{x^3} \phantom{3x^2} 3x + 1 \\
 \phantom{x^3} \phantom{3x^2} \underline{3x - 3} \\
 \phantom{x^3} \phantom{3x^2} \phantom{3x} 4
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \phantom{x^3} x^2 \quad ? \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1}
 \end{array}$$

Divide  $3x^2$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1
 \end{array}$$

Divide  $3x^2$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \phantom{3x^2} \underline{\phantom{+ 1} \text{?} \phantom{+ 1} \text{?}}
 \end{array}$$

Multiply  $3x$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 \phantom{3x^2} 4x + 1
 \end{array}$$

Multiply  $3x$  by divisor.



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 \phantom{3x^2} 3x \phantom{+ 1} \\
 \phantom{3x^2} \underline{3x - 3} \phantom{+ 1} \\
 \phantom{3x^2} \phantom{3x} 4 \phantom{+ 1}
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \quad ? \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1
 \end{array}$$

Divide  $3x$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 3x + 1
 \end{array}$$

Divide  $3x$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{\phantom{3x} 3 \phantom{+ 1}}
 \end{array}$$

Multiply 3 by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

Multiply 3 by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 ?
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

Subtract last two polynomials.



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \text{Quotient: } x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(x^3 + 2x^2 + 1) = (x^2 + 3x + 3) \cdot (x - 1) + 4$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \text{Quotient:} \quad x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

**Remainder:** 4

$$\begin{aligned}
 (\text{Dividend}) &= (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder}) \\
 (x^3 + 2x^2 + 1) &= (x^2 + 3x + 3) \cdot (x - 1) + 4
 \end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} \phantom{0} 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \end{array}$$

?

Divide  $6x^3$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3}$$

$3x^2$

Divide  $6x^3$  by  $2x$ .



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{\phantom{2x - 3} ? \phantom{00} ?}
 \end{array}$$

Multiply  $3x^2$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} 3x^2 \\ 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\ \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \end{array}$$

Multiply  $3x^2$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 \phantom{6x^3 - } 10x^2 + 17x - 3
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \quad ? \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Divide  $-10x^2$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Divide  $-10x^2$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \phantom{-} \quad ? \quad ?
 \end{array}$$

Multiply  $-5x$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Multiply  $-5x$  by divisor.



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 \phantom{-10x^2 + } 2x - 3 \\
 \phantom{-10x^2 + } \underline{2x - 3} \\
 \phantom{-10x^2 + } 0
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \quad ? \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Divide  $2x$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Divide  $2x$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{\phantom{2x} ? \phantom{0} ?}
 \end{array}$$

Multiply **1** by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

Multiply **1** by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 ?
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

Subtract last two polynomials.



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient: } 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient:} \quad 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

**Remainder:** 0

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. **Use your work to factor the cubic.** Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient:} \quad 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

**Remainder:** 0

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

$$= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = ?$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

$$= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. **Use your work to factor the cubic.** Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. **Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .**

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$6x^3 - 19x^2 + 17x - 3 = 0$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 &= 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \frac{3}{2}\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ \textcolor{red}{2}x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \textcolor{red}{\frac{3}{2}}\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \frac{3}{2}\end{aligned}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s).

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the x axis at: ?

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the x axis at:  $-1.5, -1, 2$ .



## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the x axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$?(x - ?)(x - ?)(x - ?)$$

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$?(x - (-1.5))(x - (-1))(x - 2)$$

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2)$$

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2) = (2x + 3)(x + 1)(x - 2)$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2) = (2x + 3)(x + 1)(x - 2)$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$


Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.



## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$\begin{aligned} 2x^3 + x^2 - 7x - 6 &= 0 \\ (2x + 3)(x + 1)(x - 2) &= 0 \end{aligned}$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$\begin{aligned}
 2x^3 + x^2 - 7x - 6 &= 0 \\
 (2x + 3)(x + 1)(x - 2) &= 0 \\
 x = -\frac{3}{2} \quad \text{or} \quad x = -1 \quad \text{or} \quad x = 2
 \end{aligned}$$

Make sure to practice with the graphing calculator you will on your exam(s). **The graph appears to intersect the x axis at:  $-1.5, -1, 2$ .** The left hand side should factor as:

$$\begin{aligned}
 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\
 &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\
 &= 2x^3 + x^2 - 7x - 6
 \end{aligned}$$

**Check work to make sure we guessed the roots correctly.**

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
?




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?




$$x - 3 \quad \overline{x^3 - x^2 - 8x + 6}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:   
 ? , ? , 3. What are the two roots besides 3?




$$x - 3 \overline{) x^3 - x^2 - 8x + 6}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?




$$x - 3 \quad \overline{) x^3 - x^2 - 8x + 6}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$x - 3 \overline{) \overset{?}{x^3} - x^2 - 8x + 6}$$


Divide  $x^3$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$x - 3 \overline{) \overset{x^2}{x^3 - x^2 - 8x + 6}}$$

Divide  $x^3$  by  $x$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{\phantom{x^3} ? \phantom{x^2} ?}
 \end{array}$$


Multiply  $x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$


Multiply  $x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 \phantom{x^3 - } 2x^2 - 8x + 6 \\
 \phantom{x^3 - } \underline{2x^2 - 6x} \phantom{+ 6} \\
 \phantom{x^3 - } \phantom{2x^2 - } 2x + 6 \\
 \phantom{x^3 - } \phantom{2x^2 - } \underline{2x + 6} \\
 \phantom{x^3 - } \phantom{2x^2 - } \phantom{2x + } 0
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) \begin{array}{l} x^3 - x^2 - 8x + 6 \\ x^3 - 3x^2 \end{array} } \\
 \hline
 2x^2 - 8x + 6
 \end{array}$$

Subtract last two polynomials.


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \quad ? \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{x^3 - x^2 - 8x + 6} \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{) x^3 - x^2 - 8x + 6} \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 \text{?} \quad \text{?}
 \end{array}$$


Multiply  $2x$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 \underline{x^3 - 3x^2} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \\
 6x + 6
 \end{array}$$

Multiply  $2x$  by divisor.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 \phantom{2}x^2 - 6x + 6 \\
 \phantom{2}x^2 - 6x + 6
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{x^3 - x^2 - 8x + 6} \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 2x^2 - 6x \\
 \hline
 -2x + 6
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \quad ? \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 2x^2 - 6x \\
 \hline
 -2x + 6
 \end{array}$$

Divide  $-2x$  by  $x$ .


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6
 \end{array}$$


Divide  $-2x$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{\phantom{-2x} 2x - 6} \\
 \phantom{-2x + 6} 12
 \end{array}$$


Multiply  $-2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


Multiply  $-2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 ?
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$

Subtract last two polynomials.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:   
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) + 0 = 0$$

The graph appears to intersect the x axis at:   
 ? , ? , 3. What are the two roots besides 3?



**Quotient:**  $x^2 + 2x - 2$

$$\begin{array}{r}
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) + 0 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



<b>Quotient:</b>	$x^2 + 2x - 2$
$x - 3$	$\begin{array}{r} \overline{) x^3 - x^2 - 8x + 6} \\ x^3 - 3x^2 \\ \hline 2x^2 - 8x + 6 \\ \phantom{2x^2 - } - 2x^2 + 6x \\ \hline \phantom{2x^2 - } - 2x + 6 \\ \phantom{2x^2 - } - 2x + 6 \\ \hline \phantom{2x^2 - } 0 \end{array}$
	<b>Remainder:</b> 0


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



<b>Quotient:</b>	$x^2 + 2x - 2$
$x - 3$	$\begin{array}{r} \overline{) x^3 - x^2 - 8x + 6} \\ x^3 - 3x^2 \\ \hline 2x^2 - 8x + 6 \\ \phantom{2x^2 - } - 2x^2 + 6x \\ \hline \phantom{2x^2 - } - 2x + 6 \\ \phantom{2x^2 - } - 2x + 6 \\ \hline \phantom{2x^2 - } 0 \end{array}$
	<b>Remainder: 0</b>


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x =$$



The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x =$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?





## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$


$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$


$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$


$$x = \frac{-2 \pm 2\sqrt{3}}{2}$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$


$$x = \frac{-2 \pm 2\sqrt{3}}{2} = -1 \pm \sqrt{3}.$$

The graph appears to intersect the x axis at:

**?**, **?**, 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$\begin{aligned}
 x^3 - x^2 - 8x + 6 &= 0 \\
 (x - 3)(x^2 + 2x - 2) &= 0 \\
 x - 3 = 0 \quad \text{or} \quad x &= \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1} \\
 x = 3 \quad \quad \quad x &= \frac{-2 \pm \sqrt{12}}{2} \\
 x &= \frac{-2 \pm 2\sqrt{3}}{2} = -1 \pm \sqrt{3}.
 \end{aligned}$$

The graph appears to intersect the x axis at:

$- \sqrt{3} - 1$ ,  $\sqrt{3} - 1$ , 3. What are the two roots besides 3?

Final answer:

$$x = 3 \quad \text{or} \quad x = -1 - \sqrt{3} \quad \text{or} \quad x = -1 + \sqrt{3}.$$

## Example


Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = ?$  .






## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?  
Is there another root (far away from 0)?




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?  
**Is there another root (far away from 0)?**




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$


Divide  $2x^3$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$

Divide  $2x^3$  by  $x$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$x - \frac{1}{2} \overline{) \begin{array}{l} 2x^3 + x^2 + 5x - 3 \\ \phantom{2x^3 +} 2x^2 \phantom{+ 5x - 3} \\ \hline \phantom{2x^3 +} ? \phantom{+ 5x - 3} \\ \phantom{2x^3 +} ? \phantom{+ 5x - 3} \end{array}}$$


Multiply  $2x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3}
 \end{array}$$


Multiply  $2x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 \phantom{2x^3 - } 2x^2 + 5x - 3
 \end{array}$$

? ? ?


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 \quad ? \\
 \hline
 x - \frac{1}{2} \quad \left| \begin{array}{l} 2x^3 + x^2 + 5x - 3 \\ 2x^3 - x^2 \end{array} \right. \\
 \hline
 2x^2 + 5x - 3
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x \\
 \hline
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \quad \quad \quad \underline{2x^3 - x^2} \\
 \quad \quad \quad \quad \quad 2x^2 + 5x - 3
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \phantom{2x^2 + } \underline{? \phantom{0} ?}
 \end{array}$$


Multiply  $2x$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x \\
 \hline
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \\
 \phantom{2x^2 + } 6x - 3
 \end{array}$$

Multiply  $2x$  by divisor.




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \\
 \phantom{x - \frac{1}{2}} 2x^2 + 5x - 3 \\
 \phantom{x - \frac{1}{2}} \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \\
 \phantom{x - \frac{1}{2}} \phantom{2x^2 + } 6x - 3 \\
 \phantom{x - \frac{1}{2}} \phantom{2x^2 + } \phantom{6x - } \textcolor{red}{?} \quad \textcolor{red}{?}
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{+ 5x - 3} \\
 \phantom{x - \frac{1}{2}} 2x^2 + 5x - 3 \\
 \phantom{x - \frac{1}{2}} \underline{\phantom{2x^2 + 5x - 3} 2x^2 - x} \phantom{- 3} \\
 \phantom{x - \frac{1}{2}} \phantom{2x^2 + 5x - 3} 6x - 3
 \end{array}$$

Subtract last two polynomials.

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 2x^2 + 2x \quad ? \\
 \hline
 x - \frac{1}{2} \quad \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3
 \end{array}$$


Divide  $6x$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3
 \end{array}$$


Divide  $6x$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} \phantom{6x} - 3} \\
 \phantom{x - \frac{1}{2}} 6x - 6 \\
 \underline{\phantom{x - \frac{1}{2}} \phantom{6x} 6x - 3} \\
 \phantom{x - \frac{1}{2}} 3
 \end{array}$$


Multiply 6 by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{2x^3} - 2x^3 + x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{2x^2} - 2x^2 + x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{6x} - 6x + 3} \\
 6x - 3
 \end{array}$$


Multiply 6 by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 ?
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:




$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

Subtract last two polynomials.



## Example




Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 6x - 3} \\
 0
 \end{array}$$

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2}) (2x^2 + 2x + 6) + 0 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:

**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) + 0 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:


**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

**Remainder:**

0

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:

**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

**Remainder:**

0


## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x =$$



We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example


Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x =$$

$$x = \frac{1}{2}$$



We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2}) (2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?



## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$


$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?





## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2} \quad x = \frac{-2 \pm \sqrt{-44}}{2 \cdot 2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?



## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2} \quad x = \frac{-2 \pm \sqrt{-44}}{2 \cdot 2}$$

no real solution

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

