

Calculus I

Reference: Indefinite integrals = antiderivatives

Todor Milev

2019

Indefinite Integrals

- The Evaluation Theorem establishes a connection between antiderivatives and definite integrals.
- It says that $\int_a^b f(x)dx$ equals $F(b) - F(a)$, where F is an antiderivative of f .
- We need convenient notation for writing antiderivatives.
- This is what the indefinite integral is.

Definition (Indefinite Integral)

The indefinite integral of f is another way of saying the antiderivative of f , and is written $\int f(x)dx$. In other words,

$$\int f(x)dx = F(x) \quad \text{means} \quad F'(x) = f(x).$$

Example

$$\int x^4 dx = ?$$

Example

$$\int x^4 dx = \frac{x^5}{5}$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.
- $\int \frac{1}{x} dx = \ln|x| + C$, and this is valid either on $(-\infty, 0)$ or $(0, \infty)$.