

# Precalculus

## Degree lowering formulas

Todor Milev

2019

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned}\cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ \color{red}{2}\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{\color{red}{2}}\end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$





## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha - 1 &= \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned} \quad \begin{aligned} \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\cos^2 \alpha - 1 &= \cos(2\alpha) \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha - 1 = \cos(2\alpha)$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

### Corollary (Half-Angle Formulas)

$$\sin \left( \frac{\beta}{2} \right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos \left( \frac{\beta}{2} \right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

### Corollary (Half-Angle Formulas)

$$\sin\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express  $\sin^k \alpha$  and  $\cos^k \alpha$  via lower powers of the sin and cos functions (applied to angles other than  $\alpha$ ).

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express  $\sin^k \alpha$  and  $\cos^k \alpha$  via lower powers of the sin and cos functions (applied to angles other than  $\alpha$ ).
- This technique will play a key role in integration (studied later/in another course).

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\sin^4 \alpha$$

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\sin^4 \alpha = \left( \sin^2 \alpha \right)^2$$

Recall the formulas:  $\sin^2 \beta = ?$  ,  $\cos^2 \beta = ?$  .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= \left( \sin^2 \alpha \right)^2 \\ &= \left( ? \right)^2\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= \left( \sin^2 \alpha \right)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} \left( ? \right) \end{aligned}$$



Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} ( ? ) \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + ?)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left( \frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left( \frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\
 &= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))
 \end{aligned}$$



Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left( \frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\
 &= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))
 \end{aligned}$$