

Calculus I

Derivative of rational function, part 2

Todor Milev

2019

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} \quad \left| \text{ (Leibniz notation) } \right.$$

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$$\begin{aligned} \frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) &= \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} & (\text{Leibniz notation}) \\ \left(\frac{f(x)}{g(x)} \right)' &= \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} & ' \text{ notation} \end{aligned}$$

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$$\begin{aligned} \frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) &= \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} && \text{(Leibniz notation)} \\ \left(\frac{f(x)}{g(x)} \right)' &= \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} && \text{' notation} \end{aligned}$$

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$$\begin{aligned} \frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) &= \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} && \text{(Leibniz notation)} \\ \left(\frac{f(x)}{g(x)} \right)' &= \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} && \text{' notation} \end{aligned}$$

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2}$$

(Leibniz notation)

$$\left(\frac{f(x)}{g(x)} \right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

' notation

$$\left(\frac{f}{g} \right)' = \frac{f'g - fg'}{g^2}$$

abbreviated

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2}$	<i>(Leibniz notation)</i> <i>' notation</i> <i>abbreviated</i>
$\left(\frac{f(x)}{g(x)} \right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$	
$\left(\frac{f}{g} \right)' = \frac{f'g - fg'}{g^2}$	

- The proof of the Quotient Rule is similar to the proof of the Product Rule.

Theorem (The Quotient Rule)

If f and g are differentiable and $g(x) \neq 0$, then

$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2}$	<i>(Leibniz notation)</i> <i>' notation</i> <i>abbreviated</i>
$\left(\frac{f(x)}{g(x)} \right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$	
$\left(\frac{f}{g} \right)' = \frac{f'g - fg'}{g^2}$	

- The proof of the Quotient Rule is similar to the proof of the Product Rule.
- There is an alternative algebraic proof via the Product Rule, the Power Rule and the (not yet studied) Chain Rule.

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}.$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\frac{dy}{dx} = \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\frac{dy}{dx} = \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(\textcolor{red}{?}) (-x^6 + 2) - (x^5 + 2x) (\textcolor{red}{?})}{(-x^6 + 2)^2}\end{aligned}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2)(-x^6 + 2) - (x^5 + 2x)(?)}{(-x^6 + 2)^2}\end{aligned}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2) (-x^6 + 2) - (x^5 + 2x) (\text{?})}{(-x^6 + 2)^2} \end{aligned}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2) (-x^6 + 2) - (x^5 + 2x) (-6x^5)}{(-x^6 + 2)^2}\end{aligned}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2) (-x^6 + 2) - (x^5 + 2x) (-6x^5)}{(-x^6 + 2)^2} \\ &= \frac{(-5x^{10} - 2x^6 + 10x^4 + 4) - (-6x^{10} - 12x^6)}{(-x^6 + 2)^2} \end{aligned}$$

Example (Quotient Rule, rational function)

Differentiate $y = \frac{x^5 + 2x}{-x^6 + 2}$.

Quotient Rule:

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\
 &= \frac{(5x^4 + 2) (-x^6 + 2) - (x^5 + 2x) (-6x^5)}{(-x^6 + 2)^2} \\
 &= \frac{(-5x^{10} - 2x^6 + 10x^4 + 4) - (-6x^{10} - 12x^6)}{(-x^6 + 2)^2} \\
 &= \frac{x^{10} + 10x^6 + 10x^4 + 4}{(-x^6 + 2)^2}.
 \end{aligned}$$