

Calculus II

Integral of rational function with cubic denominator, part 2

Todor Milev

2019

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$
- Factor denominator: $x^3 - x^2 - x + 1 = ?$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1)$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = ?$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$
- Plug-in $x = -1$:

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1 - 1)^2$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1 - 1)^2 \Rightarrow C = ?$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x - 1)^2(x + 1)} = \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 1}$$

$$-x^2 + 5x - 2 = A(x - 1)(x + 1) + B(x + 1) + C(x - 1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1 - 1)^2 \Rightarrow C = -2.$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$:

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1)$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = ?$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:

$$-2 = A(0-1)(0+1) + B(0+1) + C(0-1)^2$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + B(0+1) + C(0-1)^2$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = ?$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide: $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$

- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$

- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$

- Plug-in $x = 0$:

$$-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1.$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.

- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.

- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.

- Plug-in $x = 0$:

$$-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1.$$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1.$

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

= ?

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left(x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide: $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$.
- Factor denominator: $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$.
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in $x = -1$: $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$.
- Plug-in $x = 1$: $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$.
- Plug-in $x = 0$:
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$.

$Q(x)$ has linear factors with higher multiplicity

- Suppose $Q(x)$ is a product of linear factors, some of which appear with power greater than 1.

$Q(x)$ has linear factors with higher multiplicity

- Suppose $Q(x)$ is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power r , that is, $(a_1x + b_1)^r$ occurs in the factorization of $Q(x)$.

$Q(x)$ has linear factors with higher multiplicity

- Suppose $Q(x)$ is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power r , that is, $(a_1x + b_1)^r$ occurs in the factorization of $Q(x)$.
- Then instead of a single term $\frac{A}{a_1x+b_1}$ we use

$$\frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \cdots + \frac{A_r}{(a_1x + b_1)^r}$$

$Q(x)$ has linear factors with higher multiplicity

- Suppose $Q(x)$ is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power r , that is, $(a_1x + b_1)^r$ occurs in the factorization of $Q(x)$.
- Then instead of a single term $\frac{A}{a_1x+b_1}$ we use

$$\frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \cdots + \frac{A_r}{(a_1x + b_1)^r}$$

- In a similar fashion we add more partial fractions to account for all other terms of the form $(a_sx + b_s)^t$.