

**Arithmetics**  
**Fraction basics**  
**[calculator-algebra.org](http://calculator-algebra.org)**

Todor Milev

2019

# Division and the number line

- Numbers represent lengths by measuring distances.



# Division and the number line


- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.

# Division and the number line


- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.

# Division and the number line


- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.

# Division and the number line


- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.

# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

# Division and the number line

- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:


$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**



# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.

# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.
- Number on bottom: **denominator**

# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.
- Number on bottom: **denominator**, represents the number of segments we are dividing into.

# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.
- Number on bottom: **denominator**, represents the number of segments we are dividing into.

# Division and the number line

- Numbers represent lengths by measuring distances.




- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.
- Number on bottom: **denominator**, represents the number of segments we are dividing into.

# Division and the number line

- Numbers represent lengths by measuring distances.



- A segment can be divided into equal parts.
- The parts may no longer have lengths that are exact integers.
- Such lengths are represented by fractions.
- These lengths are called **fractions** and are denoted by:

$$\frac{a}{b} \quad \text{or} \quad a/b$$

- Number on top: **numerator**, represents length of divided segment.
- Number on bottom: **denominator**, represents the number of segments we are dividing into.

## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Example

Simplify.

$$\frac{5}{1} = ?$$

$$\frac{-3}{1} = ?$$



## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Example

Simplify.

$$\begin{aligned}\frac{5}{1} &= 5 \\ \frac{-3}{1} &= -3\end{aligned}$$

## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Example

Simplify.

$$\begin{aligned}\frac{5}{1} &= 5 \\ \frac{-3}{1} &= -3\end{aligned}$$

- Division by one corresponds to “division of a segment into one equal part”.

## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Example

Simplify.

$$\frac{5}{1} = 5$$
$$\frac{-3}{1} = -3$$

- Division by one corresponds to “division of a segment into one equal part”.
- “**Division of a segment into one equal part**” is interpreted as not dividing the segment at all.

## Observation

*For any number  $a$ , the following equality holds.*

$$\frac{a}{1} = a$$

## Example

Simplify.

$$\begin{aligned}\frac{5}{1} &= 5 \\ \frac{-3}{1} &= -3\end{aligned}$$

- Division by one corresponds to “division of a segment into one equal part”.
- “Division of a segment into one equal part” is interpreted as **not dividing the segment at all**.

## Example (Reading fractions)

**Read the fraction.** Honor your English dialect naming convention, if different from the one given here.

●  $\frac{1}{3}$  ?

●  $\frac{1}{2}$

●  $\frac{1}{4}$

●  $\frac{3}{4}$

●  $\frac{6}{7}$

●  $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$
- $\frac{1}{4}$
- $\frac{3}{4}$
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  ?
- $\frac{1}{4}$
- $\frac{3}{4}$
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$
- $\frac{3}{4}$
- $\frac{6}{7}$
- $\frac{11}{6}$



## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  ?
- $\frac{3}{4}$
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  ?
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  three quarters, three fourths.
- $\frac{6}{7}$
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  three quarters, three fourths.
- $\frac{6}{7}$  ?
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  three quarters, three fourths.
- $\frac{6}{7}$  six sevenths.
- $\frac{11}{6}$

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  three quarters, three fourths.
- $\frac{6}{7}$  six sevenths.
- $\frac{11}{6}$  ?

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; one fourth.
- $\frac{3}{4}$  three quarters, three fourths.
- $\frac{6}{7}$  six sevenths.
- $\frac{11}{6}$  eleven sixths.



## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
- $\frac{1}{2}$  one half.
- $\frac{1}{4}$  one quarter; **one fourth**.
- $\frac{3}{4}$  three quarters, **three fourths**.
- $\frac{6}{7}$  **six sevenths**.
- $\frac{11}{6}$  **eleven sixths**.

- Most frequently, a fraction  $\frac{a}{b}$  is read along the template “**a b-th(s)**”.

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  **one third** (can also say “a third”).
- $\frac{1}{2}$  **one half**.
- $\frac{1}{4}$  **one quarter**; one fourth.
- $\frac{3}{4}$  **three quarters**, three fourths.
- $\frac{6}{7}$  six sevenths.
- $\frac{11}{6}$  eleven sixths.

- **Most** frequently, a fraction  $\frac{a}{b}$  is read along the template “a b-th(s)”.

## Example (Reading fractions)

Read the fraction. Honor your English dialect naming convention, if different from the one given here.

- $\frac{1}{3}$  one third (can also say “a third”).
  - $\frac{1}{2}$  one half.
  - $\frac{1}{4}$  one quarter; one fourth.
  - $\frac{3}{4}$  three quarters, three fourths.
  - $\frac{6}{7}$  six sevenths.
  - $\frac{11}{6}$  eleven sixths.
- 
- Most frequently, a fraction  $\frac{a}{b}$  is read along the template “ $a$   $b$ -th(s)”.
  - Most important exceptions:  $\frac{1}{2}$  is read as “one half” (or just “half”).  
 $\frac{1}{4}$  is read as both “one fourth” and as “one quarter”.


## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$






## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$



## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$



## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$



## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$





## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$




## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

## Example

$$\frac{2}{3} + \frac{5}{3} = \frac{2+5}{3} = \frac{7}{3}$$



## Observation

*Fractions with same denominator are added by adding their numerators.*

$$\frac{a}{c} \pm \frac{b}{c} = \frac{a \pm b}{c}$$

*A similar rule holds for subtraction.*

## Example

$$\begin{array}{rclcl} \frac{2}{3} + \frac{5}{3} & = & \frac{2+5}{3} & = & \frac{7}{3} \\ \frac{4}{3} - \frac{1}{3} & = & \frac{4-1}{3} & = & \frac{3}{3} \end{array}$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} =$$

$$\frac{7}{6} + \frac{10}{6} =$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = ?$$

$$\frac{7}{6} + \frac{10}{6} =$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} =$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} =$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$



## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = ?$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} =$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = ?$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} =$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = ?$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$



## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} =$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = ?$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} =$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = ?$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8}$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8}$$

$$\frac{11}{13} - \frac{20}{13} =$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8} = -\frac{3}{8}$$

$$\frac{11}{13} - \frac{20}{13} =$$



## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8} = -\frac{3}{8}$$

$$\frac{11}{13} - \frac{20}{13} = ?$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8} = -\frac{3}{8}$$

$$\frac{11}{13} - \frac{20}{13} = \frac{-9}{13}$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8} = -\frac{3}{8}$$

$$\frac{11}{13} - \frac{20}{13} = \frac{-9}{13}$$

## Example (Add numbers with same denominator)

$$\frac{1}{4} + \frac{2}{4} = \frac{3}{4}$$

$$\frac{7}{6} + \frac{10}{6} = \frac{17}{6}$$

$$\frac{302}{111} + \frac{24}{111} = \frac{326}{111}$$

$$\frac{3}{6} - \frac{2}{6} = \frac{1}{6}$$

$$\frac{103}{101} - \frac{4}{101} = \frac{99}{101}$$

$$\frac{3}{8} - \frac{6}{8} = \frac{-3}{8} = -\frac{3}{8}$$

$$\frac{11}{13} - \frac{20}{13} = \frac{-9}{13} = -\frac{9}{13}$$

## Definition (Factor a number (properly))

To factor an integer  $a$  (properly) means to find integers  $b > 1$  and  $c > 1$  so that

$$a = \pm b \cdot c$$

The numbers  $b$  and  $c$  are called **factors**.

## Example (Proper factorization)

$$4 = 2 \cdot 2$$

$$6 = 3 \cdot 2 = 2 \cdot 3$$

$$-8 = -2 \cdot 4 = -4 \cdot 2 = -2 \cdot 2 \cdot 2$$

## Example (Not a proper factorization)

$$-3 = (-1) \cdot 3$$

$$4 = 1 \cdot 4$$

$$1 = 2 \cdot \frac{1}{2}$$

$$1 = 0.25 \cdot 4$$

## Definition

An integer greater than one is prime if it cannot be factored properly.

## Example

Prime?	Full factorization	Prime?	Full factorization
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.

## Example

Prime?	Full factorization	Prime?	Full factorization
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether **negatives can be prime** is usually left **undefined**.

## Example

Prime?	Full factorization	Prime?	Full factorization
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	



## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.

## Example

Prime?	Full factorization	Prime?	Full factorization
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

Prime?	Full factorization	Prime?	Full factorization
1		9	
2		10	
3		11	
4		12	
5		13	
6		14	
7		15	
8		16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	?		9		
2			10		
3			11		
4			12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer **greater than one** is prime if it cannot be factored properly.

- **0 and 1 are not prime by definition.**
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2			10		
3			11		
4			12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	?		10		
3			11		
4			12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3			11		
4			12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	?		11		
4			12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4			12		
5			13		
6			14		
7			15		
8			16		



## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	?		12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5			13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-		9	
2	yes	2		10	
3	yes	3		11	
4	no	$2 \cdot 2$		12	
5	?			13	
6				14	
7				15	
8				16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6			14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	?		14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7			15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	?		15		
8			16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8			16		



## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9		
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	?		16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-		9	
2	yes	2		10	
3	yes	3		11	
4	no	$2 \cdot 2$		12	
5	yes	5		13	
6	no	$2 \cdot 3$		14	
7	yes	7		15	
8	no	$2 \cdot 2 \cdot 2$		16	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	?	
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10		
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	?	
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11		
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	?	
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12		
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		



## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	?	
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13		
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	?	
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14		
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	?	
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	no	$2 \cdot 7$
7	yes	7	15		
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	no	$2 \cdot 7$
7	yes	7	15	?	
8	no	$2 \cdot 2 \cdot 2$	16		

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	no	$2 \cdot 7$
7	yes	7	15	no	$3 \cdot 5$
8	no	$2 \cdot 2 \cdot 2$	16		



## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	no	$2 \cdot 7$
7	yes	7	15	no	$3 \cdot 5$
8	no	$2 \cdot 2 \cdot 2$	16	?	

## Definition

An integer greater than one is prime if it cannot be factored properly.

- 0 and 1 are not prime by definition.
- Whether negatives can be prime is usually left undefined.
  - Both options could be made to make sense.
  - To avoid confusion: avoid question of prime negative numbers.

## Example

	Prime?	Full factorization		Prime?	Full factorization
1	no	-	9	no	$3 \cdot 3$
2	yes	2	10	no	$2 \cdot 5$
3	yes	3	11	yes	11
4	no	$2 \cdot 2$	12	no	$2 \cdot 2 \cdot 3$
5	yes	5	13	yes	13
6	no	$2 \cdot 3$	14	no	$2 \cdot 7$
7	yes	7	15	no	$3 \cdot 5$
8	no	$2 \cdot 2 \cdot 2$	16	no	$2 \cdot 2 \cdot 2 \cdot 2$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = ?$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = 2 \cdot 2 \cdot 2 \cdot 2$$

$$36 =$$



## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = 2 \cdot 2 \cdot 2 \cdot 2$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = 2 \cdot 2 \cdot 2 \cdot 2$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 =$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = ?$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = 4 \cdot 9 = 2 \cdot 2 \cdot 3 \cdot 3$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = 4 \cdot 9 = 2 \cdot 2 \cdot 3 \cdot 3$$



## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = 4 \cdot 9 = 2 \cdot 2 \cdot 3 \cdot 3$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = 4 \cdot 9 = \underbrace{2 \cdot 2}_{2 \text{ copies}} \cdot \underbrace{3 \cdot 3}_{2 \text{ copies}} = 2^2 \cdot 3^2$$

## Definition (Factor positive integer completely)

To factor a positive integer completely means to write it as a product of prime factors.

$$x = p_1 \cdot p_2 \cdots p_n$$

- It is best practice to sort (order) the prime factors. Most frequently used order: smaller factors come first.

## Lemma (Unique prime factorization)

*Up to shuffling prime factors, there is only one way to factor a number.*

- Consequence: two numbers are equal if and only if their sorted prime factorizations are equal.
- When factoring, we may or may not use exponent notation:

$$16 = 4 \cdot 4 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2}_{4 \text{ copies}} = 2^4$$

$$36 = 4 \cdot 9 = \underbrace{2 \cdot 2}_{2 \text{ copies}} \cdot \underbrace{3 \cdot 3}_{2 \text{ copies}} = 2^2 \cdot 3^2$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = ?$$

$$6 =$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 =$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. **If applicable, show two answers:** with and **without exponent notation** ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 =$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: **with** and without **exponent notation** ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 =$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = ?$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$



## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. **If applicable**, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3 \text{ [exponent notation not needed]}$$

$$7 =$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = ?$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 =$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = ?$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$



## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 =$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = ?$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 =$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = ?$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 =$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = ?$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$



## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 =$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = ?$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$



## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 =$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = ?$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 =$$

$$91 =$$



## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 =$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 = ?$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 = \text{already factored (prime number)}$$

$$91 =$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 = \text{already factored (prime number)}$$

$$91 = ?$$

## Example

Factor the number completely. If applicable, show two answers: with and without exponent notation ( $x^3$  vs  $x \cdot x \cdot x$ ).

$$4 = 2 \cdot 2 = 2^2$$

$$6 = 2 \cdot 3$$

$$7 = \text{already factored (prime number)}$$

$$8 = 2 \cdot 4 = 2 \cdot 2 \cdot 2 = 2^3$$

$$9 = 3 \cdot 3 = 3^2$$

$$15 = 3 \cdot 5$$

$$24 = 8 \cdot 3 = 2 \cdot 4 \cdot 3 = 2 \cdot 2 \cdot 2 \cdot 3 = 2^3 \cdot 3$$

$$36 = 6 \cdot 6 = 2 \cdot 3 \cdot 2 \cdot 3 = 2 \cdot 2 \cdot 3 \cdot 3 = 2^2 \cdot 3^2$$

$$52 = 4 \cdot 13 = 2 \cdot 2 \cdot 13 = 2^2 \cdot 13$$

$$67 = \text{already factored (prime number)}$$

$$91 = 7 \cdot 13$$

### Definition (Factor a number)

To factor an integer  $a$  means to find integers  $b > 1$  and  $c > 1$  so that

$$a = \pm b \cdot c$$

We say that  $b, c$  are **factors** of  $a$ .

### Definition (Prime number)

A number is prime if it cannot be factored.

### Definition (Complete factorization)

To find a complete factorization of an integer  $a$  means to find prime numbers  $p_1 > 1, p_2 > 1, \dots, p_n > 1$  with

$$a = \pm p_1 \cdot p_2 \cdots p_n$$

- Is the number 67414977753059 prime?

$$67414977753059 =$$

- Is the number 67414977753059 prime? **No**

$$\underbrace{67414977753059}_{\text{not prime}} =$$



- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} =$$

- Even when  $x$  is large, there exist **fast computer algorithms** to check whether  $x$  is prime.

- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} =$$

- Even when  $x$  is large, there exist fast computer algorithms to check whether  $x$  is prime.
- In other words, there exist fast algorithms for knowing whether a proper factorization exists.

- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} =$$

- Even when  $x$  is large, there exist fast computer algorithms to check whether  $x$  is prime.
- In other words, there exist fast algorithms for knowing whether a proper factorization exists.
- However, even when we know  $x$  can be factored, as of 2019, there are no known fast computer algorithms for finding an actual factorization.

- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} = \underbrace{11494253}_{\text{prime}} \cdot \underbrace{5865103}_{\text{prime}}$$

- Even when  $x$  is large, there exist fast computer algorithms to check whether  $x$  is prime.
- In other words, there exist fast algorithms for knowing whether a proper factorization exists.
- However, even when we know  $x$  can be factored, as of 2019, there are **no known fast computer algorithms** for **finding an actual factorization**.

- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} = \underbrace{11494253}_{\text{prime}} \cdot \underbrace{5865103}_{\text{prime}}$$

- Even when  $x$  is large, there exist fast computer algorithms to check whether  $x$  is prime.
- In other words, there exist fast algorithms for knowing whether a proper factorization exists.
- However, even when we know  $x$  can be factored, as of 2019, there are no known fast computer algorithms for finding an actual factorization.
- In fact, the number above was generated by first making two large primes and then multiplying them.

- Is the number 67414977753059 prime? No

$$\underbrace{67414977753059}_{\text{not prime}} = \underbrace{11494253}_{\text{prime}} \cdot \underbrace{5865103}_{\text{prime}}$$

- Even when  $x$  is large, there exist fast computer algorithms to check whether  $x$  is prime.
- In other words, there exist fast algorithms for knowing whether a proper factorization exists.
- However, even when we know  $x$  can be factored, as of 2019, there are no known fast computer algorithms for finding an actual factorization.
- In fact, the number above was generated by first making two large primes and then multiplying them.
- Each of the two known large primes above was in turn generated by trying large integers at random.


## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3}$$



## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3}$$





## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{\textcolor{red}{b}} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{\textcolor{red}{3}}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6}$$





## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\frac{2}{3} = \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{3 \cdot 2}{3 \cdot 3} \end{aligned}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{3 \cdot 2}{3 \cdot 3} \end{aligned}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{2 \cdot 2}{3 \cdot 2} \\ &= \frac{4}{\color{red}{3 \cdot 3}} \end{aligned}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{3 \cdot 2}{3 \cdot 3} = \frac{6}{9} \end{aligned}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{3 \cdot 2}{3 \cdot 3} = \frac{6}{9} \end{aligned}$$




## Observation

*Fractions do not change when we multiply their numerator and denominator by the same number.*

$$\frac{a}{b} = \frac{x \cdot a}{x \cdot b}$$

## Example

$$\begin{aligned} \frac{2}{3} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{4}{6} \\ &= \frac{3 \cdot 2}{3 \cdot 3} = \frac{6}{9} \end{aligned}$$



## Definition (Reduce a fraction)

A positive fraction  $\frac{a}{b}$  is a reduction of a positive fraction  $\frac{A}{B}$  when

$$\frac{A}{B} = \frac{a}{b}$$

and  $\frac{a}{b}$  has smaller numerator and denominator, i.e.,  $A > a$  and  $B > b$ .

Recall that  $\frac{x \cdot a}{x \cdot b} = \frac{a}{b}$ .

## Example

Reduce the fraction.

$$\begin{aligned}\frac{4}{6} &= \frac{2 \cdot 2}{2 \cdot 3} = \frac{2}{3} \\ \frac{2}{4} &= \frac{2 \cdot 1}{2 \cdot 2} = \frac{1}{2} \\ \frac{3}{9} &= \frac{3 \cdot 1}{3 \cdot 3} = \frac{1}{3}\end{aligned}$$



- To reduce a fraction, we use the rule:

$$\frac{x \cdot a}{x \cdot b} = \frac{a}{b}.$$

- To reduce a fraction, we use the rule:

$$\frac{x \cdot a}{x \cdot b} = \frac{\cancel{x} \cdot a}{\cancel{x} \cdot b} = \frac{a}{b}.$$

- To reduce excessive copying: use cancel notation.

- To reduce a fraction, we use the rule:

$$\frac{x \cdot a}{x \cdot b} = \frac{\cancel{x} \cdot a}{\cancel{x} \cdot b} = \frac{a}{b}.$$

- To reduce excessive copying: use cancel notation.
- The uses of the cancel notation will become apparent in examples.

- To reduce a fraction, we use the rule:

$$\frac{x \cdot a}{x \cdot b} = \frac{\cancel{x} \cdot a}{\cancel{x} \cdot b} = \frac{a}{b}.$$

- To reduce excessive copying: use cancel notation.
- The uses of the cancel notation will become apparent in examples.
- Rules.
  - Use a single slanted line.
  - Unless circumstances dictate otherwise, slant your line from lower left corner to top right corner.
  - Do not use crosses, smudges, or any other notation that obscures the expression below the cancel line.

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} =$$
$$\frac{14}{6} =$$
$$\frac{6}{2} =$$
$$\frac{6}{5} =$$
$$\frac{10}{15} =$$
$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} =$$
$$\frac{14}{6} =$$
$$\frac{6}{2} =$$
$$\frac{6}{5} =$$
$$\frac{10}{15} =$$
$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{3 \cdot 3}{3 \cdot 4}$$

$$\frac{14}{6} =$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\begin{array}{rcl} \frac{9}{12} & = & \frac{3 \cdot 3}{3 \cdot 4} \\ \frac{14}{6} & = & \\ \frac{6}{2} & = & \\ \frac{6}{5} & = & \\ \frac{10}{15} & = & \\ \frac{8}{6} & = & \end{array}$$



## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} =$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} =$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} =$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} = \frac{2 \cdot 7}{2 \cdot 3}$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{2 \cdot 7}{2 \cdot 7}$$

$$\frac{6}{6} = \frac{2 \cdot 3}{2 \cdot 3}$$

$$\frac{6}{2} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 3} = \frac{7}{3}$$

$$\frac{6}{2} =$$

$$\frac{6}{6} =$$

$$\frac{5}{10} =$$

$$\frac{15}{8} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 3} = \frac{7}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} =$$

$$\frac{2}{6} =$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{6}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{6} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 3} = \frac{7}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$



## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{\textcolor{red}{6}}{2} = \frac{\textcolor{red}{2} \cdot \textcolor{red}{3}}{2 \cdot 1}$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{2 \cdot 3}{2 \cdot 1}$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{6}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1}$$

$$\frac{5}{5} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\begin{array}{rcl}
 \frac{9}{12} & = & \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4} \\
 \frac{14}{14} & = & \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7} \\
 \frac{6}{6} & = & \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3} \\
 \frac{6}{2} & = & \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3 \\
 \frac{6}{5} & = & \\
 \frac{10}{15} & = & \\
 \frac{8}{6} & = &
 \end{array}$$

## Example

Simplify (reduce) the fraction.

$$\begin{array}{rcl}
 \frac{9}{12} & = & \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4} \\
 \frac{14}{14} & = & \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7} \\
 \frac{6}{6} & = & \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3} \\
 \frac{\color{red}{6}}{\color{red}{2}} & = & \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = \color{red}{3} \\
 \frac{6}{5} & = & \\
 \frac{10}{15} & = & \\
 \frac{8}{6} & = &
 \end{array}$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{\textcolor{red}{6}}{\textcolor{red}{5}} = \textcolor{red}{?}$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$



## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

$$\frac{10}{15} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{\cancel{5} \cdot 2}{\cancel{5} \cdot 3} = \frac{2}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{10}{15} = \frac{2 \cdot 5}{3 \cdot 5}$$

$$\frac{8}{6} = \frac{4 \cdot 2}{3 \cdot 2}$$

$$\frac{8}{6} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{4}{3}$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot 5}{3 \cdot 5}$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{14}{14} = \frac{\cancel{2} \cdot 7}{\cancel{2} \cdot 7} = \frac{7}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

$$\frac{8}{6} =$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*



## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{2 \cdot 3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{2 \cdot 3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{2 \cdot 3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{2 \cdot 3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{2 \cdot 3}$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{2 \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

## Example

Simplify (reduce) the fraction.

$$\frac{9}{12} = \frac{\cancel{3} \cdot 3}{\cancel{3} \cdot 4} = \frac{3}{4}$$

$$\frac{12}{14} = \frac{\cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{6}{7}$$

$$\frac{6}{6} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 3} = \frac{3}{3}$$

$$\frac{6}{2} = \frac{\cancel{2} \cdot 3}{\cancel{2} \cdot 1} = \frac{3}{1} = 3$$

$$\frac{6}{5} = \text{already simplified: } \frac{6}{5} = \frac{3 \cdot 2}{5}, \text{ no primes to cancel}$$

$$\frac{10}{15} = \frac{2 \cdot \cancel{5}}{3 \cdot \cancel{5}} = \frac{2}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

$$\frac{8}{6} = \frac{\cancel{2} \cdot 4}{\cancel{2} \cdot 3} = \frac{4}{3}$$

## Lemma

*If there are no common prime factors between the numerator and the denominator, the fraction is reduced.*

- There exists an algorithm for simplifying every fraction.

- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.

- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.
- Instead, the algorithm involves the notion of a greatest common divisor (GCD).

- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.
- Instead, the algorithm involves the notion of a greatest common divisor (GCD).
- We will study greatest common divisors (GCD) and the fraction simplification GCD algorithm later.



- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.
- Instead, the algorithm involves the notion of a greatest common divisor (GCD).
- We will study greatest common divisors (GCD) and the fraction simplification GCD algorithm later.
- That fraction simplification GCD algorithm is well-suited for and very fast on a computer.

- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.
- Instead, the algorithm involves the notion of a greatest common divisor (GCD).
- We will study greatest common divisors (GCD) and the fraction simplification GCD algorithm later.
- That fraction simplification GCD algorithm is well-suited for and very fast on a computer.
- This algorithm is also practical for use by hand.

- There exists an algorithm for simplifying every fraction.
- The algorithm is considerably different from what we exercised so far: it does not require us to factor the numerator and denominator.
- Instead, the algorithm involves the notion of a greatest common divisor (GCD).
- We will study greatest common divisors (GCD) and the fraction simplification GCD algorithm later.
- That fraction simplification GCD algorithm is well-suited for and very fast on a computer.
- This algorithm is also practical for use by hand.
- However on small examples the factorization-cancellation guess-work technique shown in examples is faster for a human.

## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*


## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$






## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{\color{red}3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$




## Observation

$$x \cdot \frac{a}{b} = \frac{x \cdot a}{b}$$

*Multiplying a fraction by a number is equivalent to multiplying its numerator by that number.*

## Example

$$2 \cdot \frac{2}{3} = \frac{4}{3}$$



## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} =$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} =$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$



## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3}$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} =$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{3 \cdot 2}{?}$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{3 \cdot 2}{?}$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{3 \cdot 2}{3 \cdot 5}$$

$$3 \cdot \frac{1}{3} =$$

$$7 \cdot \frac{3}{21} =$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned} 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\ 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\ 3 \cdot \frac{1}{3} &= \\ 7 \cdot \frac{3}{21} &= \\ 6 \cdot \frac{2}{15} &= \\ 4 \cdot \frac{5}{18} &= \end{aligned}$$



## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 \color{red}{3} \cdot \frac{1}{3} &= \frac{\color{red}{3}}{3} \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3} \cdot 1}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \frac{7 \cdot 3}{21}
 \end{aligned}$$

$$\begin{aligned}
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \frac{7 \cdot 3}{21} = \frac{21}{21}
 \end{aligned}$$

$$\begin{aligned}
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$



## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1 \\
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$\begin{aligned}
 2 \cdot \frac{2}{3} &= \frac{2 \cdot 2}{3} = \frac{4}{3} \\
 3 \cdot \frac{2}{15} &= \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5} \\
 3 \cdot \frac{1}{3} &= \frac{\cancel{3}}{\cancel{3}} = 1 \\
 7 \cdot \frac{3}{21} &= \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1
 \end{aligned}$$

$$\begin{aligned}
 6 \cdot \frac{2}{15} &= \\
 4 \cdot \frac{5}{18} &=
 \end{aligned}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{3}{\color{red}{3 \cdot 7}}$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7}$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = \textcolor{red}{7} \cdot \frac{1}{7} = \textcolor{red}{7}$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} =$$

$$4 \cdot \frac{5}{18} =$$



## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{\color{red}{6} \cdot 2}{15} = \frac{\color{red}{2} \cdot \color{red}{3} \cdot 2}{3 \cdot 5}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} =$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$\textcolor{red}{4} \cdot \frac{5}{18} = \frac{\textcolor{red}{4} \cdot 5}{?}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{?}$$



## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{\color{red}{3} \cdot \color{red}{6}}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot \cancel{6}} = \frac{2 \cdot 2 \cdot 5}{3 \cdot \cancel{2} \cdot 3}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot 6} = \frac{2 \cdot 2 \cdot 5}{3 \cdot 2 \cdot 3}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot 6} = \frac{2 \cdot \cancel{2} \cdot 5}{3 \cdot \cancel{2} \cdot 3} = \frac{2 \cdot 5}{3 \cdot 3}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot 6} = \frac{2 \cdot \cancel{2} \cdot 5}{3 \cdot \cancel{2} \cdot 3} = \frac{2 \cdot 5}{3 \cdot 3} = \frac{10}{9}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{21}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot 6} = \frac{2 \cdot \cancel{2} \cdot 5}{3 \cdot \cancel{2} \cdot 3} = \frac{2 \cdot 5}{3 \cdot 3} = \frac{10}{9}$$

## Example

Simplify the expression to a single reduced fraction.

$$2 \cdot \frac{2}{3} = \frac{2 \cdot 2}{3} = \frac{4}{3}$$

$$3 \cdot \frac{2}{15} = \frac{\cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

$$3 \cdot \frac{1}{3} = \frac{\cancel{3}}{\cancel{3}} = 1$$

$$7 \cdot \frac{3}{21} = \frac{7 \cdot 3}{21} = \frac{\cancel{7} \cdot \cancel{3}}{\cancel{21}} = 1$$

$$[\text{alternatively}] = 7 \cdot \frac{\cancel{3}}{\cancel{3} \cdot 7} = 7 \cdot \frac{1}{7} = \frac{\cancel{7}}{\cancel{7}} = 1$$

$$6 \cdot \frac{2}{15} = \frac{6 \cdot 2}{15} = \frac{2 \cdot \cancel{3} \cdot 2}{\cancel{3} \cdot 5} = \frac{4}{5}$$

$$4 \cdot \frac{5}{18} = \frac{4 \cdot 5}{3 \cdot 6} = \frac{2 \cdot \cancel{2} \cdot 5}{3 \cdot \cancel{2} \cdot 3} = \frac{2 \cdot 5}{3 \cdot 3} = \frac{10}{9}$$