

## Precalculus

# Compute trigonometric function of a complementary angle, part 1

Todor Milev

2019

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.
- The Quadrant I part of the proof serves as a visual aid for memorization.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.
- The Quadrant I part of the proof serves as a visual aid for memorization.
- There is an algebraically simpler (but theoretically advanced) way to prove the above identities through the angle sum formulas, derived in turn from Euler's formula (studied later/in another course).




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = \frac{|PQ|}{|OP|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \quad \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \quad \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} & \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|} \\ &= \cos \alpha & \left. \text{as desired} \right|\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$

□




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|}\end{aligned}$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R|}{|OP|} = -\frac{|OQ'|}{|OP|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$

as desired




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) =$$






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \Bigg| \quad \square ORPQ \\ &= -\frac{|P'Q|}{|OP'|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|} \\ &= -\sin \alpha.\end{aligned}$$

□




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.


$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} & \square ORPQ \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|} \\ &= -\sin \alpha. & \text{as desired } \square\end{aligned}$$

# Cofunction identities


## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

To memorize the cofunction identities it suffices to memorize the Quadrant I case via the two diagrams below.



$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \frac{|OQ|}{|OP|}\end{aligned}$$



$$\begin{aligned}\sin\left(\frac{\pi}{2} + \alpha\right) &= \frac{|OQ'|}{|OP|} \\ \cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|PQ'|}{|OP'|} = -\frac{|PR|}{|OP|}\end{aligned}$$