
Precalculus

Trigonometric functions with arguments translated by a multiple of $\frac{\pi}{2}$

Todor Milev

2019

$$\cos\left(\frac{\pi}{2}-x\right)$$




$$\cos\left(\frac{\pi}{2} - x\right) = \cos\left(\frac{\pi}{2}\right)\cos x + \sin\left(\frac{\pi}{2}\right)\sin x$$

$$= 0 \cdot \cos(x) + 1 \cdot \sin x$$

$$= \sin x$$

$$\cot\left(\frac{3\pi}{2}+x\right)$$



cot
$$\left(\frac{3\pi}{2} + x\right)$$
 = $\frac{\cos\left(\frac{3\pi}{2} + x\right)}{\sin\left(\frac{3\pi}{2} + x\right)}$ = $\frac{\cos\left(\frac{3\pi}{2} + x\right)}{\sin\left(\frac{3\pi}{2}\right)\cos x - \sin\left(\frac{3\pi}{2}\right)\sin x}$ = $\frac{\sin\left(\frac{3\pi}{2}\right)\cos x + \cos\left(\frac{3\pi}{2}\right)\sin x}{(-1)\cos x + \cos x}$ = $\frac{\sin x}{-\cos x}$ = $-\frac{\sin x}{\cos x}$ = $-\tan x$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$
$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$\tan(\pi + X) = \frac{\sin(\pi + X)}{\cos(\pi + X)}$$
$$= \frac{\sin \pi \cos X + \cos \pi \sin X}{\cos \pi \cos X - \sin \pi \sin X}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{? \cdot \cos x - ? \cdot \sin x}{? \cdot \cos x - ? \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x - \sin \pi \sin x}{2 \cdot \cos x - 2 \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + \mathbf{?} \cdot \sin x}{\mathbf{?} \cdot \cos x - \mathbf{?} \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{2 \cdot \cos x - 2 \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x - \sin \pi \sin x}{2 \cdot \cos x - 2 \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x - \sin \pi \sin x}{(-1) \cdot \cos x - ? \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - ? \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x - \sin \pi \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}$$

$$= \frac{-\sin x}{-\cos x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}$$

$$= \frac{-\sin x}{-\cos x}$$

$$= \frac{\sin x}{\cos x}$$

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}$$

$$= \frac{-\sin x}{-\cos x}$$

$$= \frac{\sin x}{\cos x}$$

$$= \tan x,$$

Show that $tan(\pi + x) = tan x$ using the angle sum formulas.

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

$$= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}$$

$$= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}$$

$$= \frac{-\sin x}{-\cos x}$$

$$= \frac{\sin x}{\cos x}$$

$$= \tan x,$$

as desired.

Proposition (tan, cot are π -periodic)

The tangent and cotangent functions are π -periodic, in other words,

$$\tan(\theta + \pi) = \tan \theta \\
\cot(\theta + \pi) = \cot \theta$$