

## Calculus II

Integrals of the form  $\int \frac{Ax + B}{ax^2 + bx + c} dx$ ,  
denominator has no real roots

Todor Milev

2019

# Linear substitutions leading to blocks IIa and IIIa

Building block IIa:  $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa:  $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let  $ax^2 + bx + c$  have no real roots.
- We can find  $p, q$  so that the linear substitution  $u = px + q$  transforms the quadratic to:

$$ax^2 + bx + c = r(u^2 + 1)$$

(where  $r$  is some number to be determined).

- To find  $p, q$ , we complete the square.
- In this way, integrals of the form  $\int \frac{Ax + B}{ax^2 + bx + c} dx$  are transformed to combinations of building blocks IIa and IIIa.
- We show examples; the general case is analogous and we leave it to the student.

# Linear substitutions leading to blocks IIa and IIIa

Building block IIa:  $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa:  $\int \frac{1}{1+x^2} dx = \arctan x + C.$

## Example

No real roots  $\Rightarrow$  complete the square. Let  $u = x + \frac{1}{2}$ , let  $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\ &= \int \frac{u - \frac{1}{2}}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \end{aligned}$$

# Linear substitutions leading to blocks IIa and IIIa

Building block IIa:  $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa:  $\int \frac{1}{1+x^2} dx = \arctan x + C.$

## Example

No real roots  $\Rightarrow$  complete the square. Let  $u = x + \frac{1}{2}$ , let  $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left( \frac{4}{3} u^2 + 1 \right)} du \\ &= \int \frac{1}{\frac{3}{4} \left( \left( \frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left( \frac{2u}{\sqrt{3}} \right) \\ &= \frac{2\sqrt{3}}{3} \int \frac{1}{z^2 + 1} dz = \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

# Linear substitutions leading to blocks IIa and IIIa

Building block IIa:  $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa:  $\int \frac{1}{1+x^2} dx = \arctan x + C.$

## Example

No real roots  $\Rightarrow$  complete the square. Let  $u = x + \frac{1}{2}$ , let  $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\begin{aligned} \int \frac{u}{u^2 + \frac{3}{4}} du &= \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{u^2}{2}\right) \\ &= \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} d\left(u^2 + \frac{3}{4}\right) = \frac{1}{2} \ln\left(u^2 + \frac{3}{4}\right) + C \end{aligned}$$

# Linear substitutions leading to blocks IIa and IIIa

Building block IIa:  $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa:  $\int \frac{1}{1+x^2} dx = \arctan x + C.$

## Example

No real roots  $\Rightarrow$  complete the square. Let  $u = x + \frac{1}{2}$ , let  $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left( u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \\ &= \frac{1}{2} \ln \left( \left( x + \frac{1}{2} \right)^2 + \frac{3}{4} \right) - \frac{\sqrt{3}}{3} \arctan \left( \frac{2u}{\sqrt{3}} \right) + C \\ &= \frac{1}{2} \ln \left( x^2 + x + 1 \right) - \frac{\sqrt{3}}{3} \arctan \left( \frac{2x+1}{\sqrt{3}} \right) + C \end{aligned}$$