

Precalculus Homework

Definition of the trigonometric functions and basic computations

1. Convert from degrees to radians.

(a) 15° .

ANSWER: $\frac{12}{\pi} \approx 0.261799388$

(b) 30° .

ANSWER: $\frac{6}{\pi} \approx 0.523598776$

(c) 36° .

ANSWER: $\frac{5}{\pi} \approx 0.628318531$

(d) 45° .

ANSWER: $\frac{4}{\pi} \approx 0.785398163$

(e) 60° .

ANSWER: $\frac{3}{\pi} \approx 1.047197551$

(f) 75° .

ANSWER: $\frac{12}{5\pi} \approx 1.308997$

(g) 90° .

ANSWER: $\frac{2}{\pi}$

(h) 120° .

(i) 135° .

(j) 150° .

(k) 180° .

(l) 225° .

(m) 270° .

(n) 305° .

ANSWER: $\frac{3}{2\pi}$

(o) 360° .

ANSWER: $\frac{4}{3\pi}$

(p) 405° .

ANSWER: $\frac{6}{5\pi}$

(q) 1200° .

ANSWER: π

(r) -900° .

ANSWER: $-\frac{4}{5\pi}$

(s) -2014° .

ANSWER: $\frac{7}{3\pi}$

ANSWER: $-\frac{1007}{90}\pi \approx -35.150931$

2. Convert from radians to degrees. The answer key has not been proofread, use with caution.

(a) 4π .

ANSWER: 720°

(b) $-\frac{7}{6}\pi$.

ANSWER: -210°

(c) $\frac{7}{12}\pi$.

ANSWER: 105°

(d) $\frac{4}{3}\pi$.

(e) $-\frac{3}{8}\pi$.

(f) 2014π .

ANSWER: 240°

ANSWER: -67.5°

ANSWER: 362520°

(g) 5 .

ANSWER: $\left(\frac{\pi}{900}\right)^\circ \approx 286^\circ$

(h) -2014 .

ANSWER: -362520°

3. Find the indicated circle arc-length. The answer key has not been proofread, use with caution.

(a) Circle of radius 3, arc of measure 36° .

ANSWER: $\frac{5}{3}\pi \approx 1.884956$

(b) Circle of radius $\frac{1}{2}$, arc of measure 100° .

ANSWER: $\frac{18}{5}\pi \approx 0.872665$


(c) Circle of radius 1, arc of measure 3 (radians).

ANSWER: 3

(d) Circle of radius 3, arc of measure 300° .

ANSWER: $5\pi \approx 15.707963$

4. Find the 6 trigonometric functions of the indicated angle in the indicated right triangle.



(a)

$$\frac{3}{\sqrt{13}} = \theta \csc \theta, \frac{2}{\sqrt{13}} = \theta \sec \theta, \frac{3}{2} = \theta \cot \theta, \frac{2}{3} = \theta \tan \theta, \frac{1}{\sqrt{13}} = \theta \cos \theta, \frac{3}{\sqrt{13}} = \theta \sin \theta$$



(b)


$$\sqrt{5} = \theta \csc \theta, \frac{2}{\sqrt{5}} = \theta \sec \theta, \frac{1}{2} = \theta \cot \theta, \frac{2}{1} = \theta \tan \theta, \frac{2}{\sqrt{5}} = \theta \cos \theta, \frac{1}{\sqrt{5}} = \theta \sin \theta$$



(c)

$$\frac{2}{\sqrt{29}} = \theta \csc \theta, \frac{5}{\sqrt{29}} = \theta \sec \theta, \frac{5}{2} = \theta \cot \theta, \frac{2}{5} = \theta \tan \theta, \frac{6\sqrt{29}}{2} = \theta \cos \theta, \frac{6\sqrt{29}}{5} = \theta \sin \theta$$

(d)



$$\frac{11}{\sqrt{13}} = \theta \csc \theta, \frac{11}{9} = \theta \sec \theta, \frac{11}{5} = \theta \cot \theta, \frac{9}{11} = \theta \tan \theta, \frac{9}{\sqrt{13}} = \theta \cos \theta, \frac{9}{11} = \theta \sin \theta$$

5. Find the exact value of the trigonometric function (using radicals).

(a) $\cos 135^\circ$.

ANSWER:

(b) $\sin 225^\circ$.

ANSWER:

(c) $\cos 495^\circ$.

ANSWER:

(d) $\sin 560^\circ$.

ANSWER:

(e) $\sin \left(\frac{3\pi}{2} \right)$.

ANSWER:

(f) $\cos \left(\frac{11\pi}{6} \right)$.

ANSWER:

$$(g) \sin\left(\frac{2015\pi}{3}\right).$$

$$(h) \cos\left(\frac{17\pi}{3}\right).$$

6. Find all solutions of the equation in the interval $[0, 2\pi)$. The answer key has not been proofread, use with caution.

$$(a) \sin x = -\frac{\sqrt{2}}{2}.$$

$$(b) \cos x = \frac{\sqrt{3}}{2}.$$


$$(c) \sin(3x) = \frac{1}{2}.$$

$$(d) \cos(7x) = 0.$$

$$(e) \cos\left(3x + \frac{\pi}{2}\right) = 0.$$

$$(f) \sin\left(5x - \frac{\pi}{3}\right) = 0.$$

Solution. 6.a



$$\sin x = -\frac{\sqrt{2}}{2}$$

Since $\sin x$ is negative it must be either in Quadrant III or IV. Therefore the angle x is coterminal either with $225^\circ = \frac{5\pi}{4}$ (Quadrant III) or $315^\circ = \frac{7\pi}{4}$ (Quadrant IV).

Case 1. x is coterminal with $225^\circ = \frac{5\pi}{4}$. We can compute

$$\begin{aligned} x &= \frac{5\pi}{4} + 2k\pi & \left| \begin{array}{l} k \text{ is any integer} \end{array} \right. \\ x &= \frac{5\pi}{4} + \frac{8k\pi}{4} \\ x &= \frac{5\pi + 8k\pi}{4} \\ x &= \frac{\pi(5 + 8k)}{4} \end{aligned}$$


We are looking for solutions in the interval $[0, 2\pi)$ and so we must discard those values of the integer k for which $\frac{\pi(5+8k)}{4}$ is negative or is greater than or equal to 2π . Therefore the only solution in this case is $x = \frac{5\pi}{4}$.

Case 2.

$$\begin{aligned}
 x &= \frac{7\pi}{4} + 2k\pi \\
 x &= \frac{7\pi}{4} + \frac{8k\pi}{4} \\
 x &= \frac{7\pi + 8k\pi}{4} \\
 x &= \frac{\pi(7 + 8k)}{4}
 \end{aligned}$$

We are looking for solutions in the interval $[0, 2\pi)$ and so we must discard those values of the integer k for which $\frac{\pi(7+8k)}{4}$ is negative or is greater than or equal to 2π . Therefore the only solution in this case is $x = \frac{7\pi}{4}$.

Solution. 6.f



$$\sin\left(5x - \frac{\pi}{3}\right) = 0$$

Since $\sin 0 = 0$ and $\sin 180^\circ = \sin \pi = 0$, the angle $5x - \frac{\pi}{3}$ must be coterminal with 0 or π .

Case 1. $5x - \frac{\pi}{3}$ is coterminal with 0. We compute

$$\begin{aligned}
 5x - \frac{\pi}{3} &= 0 + 2k\pi \\
 5x &= \frac{\pi}{3} + 2k\pi \\
 x &= \frac{\frac{\pi}{3} + 2k\pi}{5} \\
 x &= \frac{\frac{\pi}{3} + \frac{6k\pi}{3}}{5} \\
 x &= \frac{\frac{\pi + 6k\pi}{3}}{5} \\
 x &= \frac{\pi + 6k\pi}{15} \\
 x &= \frac{\pi(1 + 6k)}{15}
 \end{aligned}$$

$$x = \cancel{\frac{\pi}{15}}, \frac{\pi[1 + 6(0)]}{15}, \frac{\pi[1 + 6(1)]}{15}, \frac{\pi[1 + 6(2)]}{15}, \frac{\pi(1 + 12)}{15}, \frac{\pi[1 + 6(3)]}{15}, \frac{\pi[1 + 6(4)]}{15}, \cancel{\frac{\pi(1 + 24)}{15}}$$

$$x = \frac{\pi}{15}, \frac{7\pi}{15}, \frac{13\pi}{15}, \frac{19\pi}{15}, \frac{25\pi}{15}.$$

Discard other values of k as they yield angles outside of $[0, 2\pi)$

Case 2.

$$5x - \frac{\pi}{3} = \pi + 2k\pi$$

$$5x = \pi + \frac{\pi}{3} + 2k\pi$$

$$5x = \frac{4\pi}{3} + 2k\pi$$

$$x = \frac{\frac{4\pi}{3} + 2k\pi}{5}$$

$$x = \frac{\frac{4\pi}{3} + \frac{6k\pi}{3}}{5}$$

$$x = \frac{\frac{4\pi + 6k\pi}{3}}{5}$$

$$x = \frac{4\pi + 6k\pi}{15}$$

$$x = \frac{2\pi(2 + 3k)}{15}$$

$$x = \cancel{\frac{2\pi(2 + 3(0))}{15}}, \frac{2\pi(2 + 3(1))}{15}, \frac{2\pi(2 + 3(2))}{15}, \frac{2\pi(2 + 3(3))}{15}, \frac{2\pi(2 + 3(4))}{15}, \cancel{\frac{2\pi(2 + 3(5))}{15}}$$

$$x = \frac{4\pi}{15}, \frac{10\pi}{15}, \frac{16\pi}{15}, \frac{22\pi}{15}, \frac{28\pi}{15}.$$

Discard other values of k as they yield angles outside of $[0, 2\pi)$

Our final answer (combined from the two cases) is $x = \frac{\pi}{15}, \frac{4\pi}{15}, \frac{7\pi}{15}, \frac{2\pi}{3}, \frac{13\pi}{15}, \frac{16\pi}{15}, \frac{19\pi}{15}, \frac{22\pi}{15}, \frac{5\pi}{3}$ or $\frac{28\pi}{15}$.