

Precalculus

Definition of the trigonometric functions and basic computations

Todor Milev


2019

Outline

1 Trigonometry


- Definition of the Trigonometric Functions
- Basic Computations with Trigonometric Functions
- Reference Angles
- Geometric Interpretation of the Trigonometric Functions
- Periodicity and Symmetries of the Trig Functions

Definition of the trigonometric functions




- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.

Definition of the trigonometric functions




- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.

Definition of the trigonometric functions




- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.

Definition of the trigonometric functions




- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.

Definition of the trigonometric functions




- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.


Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by

$$\cos \theta = x$$

$$\sin \theta = y.$$

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.


Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by

$$\cos \theta = x$$

$$\sin \theta = y.$$

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.

Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by


$$\cos \theta = x \qquad \sin \theta = y.$$

Definition (additional trigonometric functions)

The functions **tangent**, cotangent, secant and cosecant of the angle θ , denoted by $\tan \theta$, $\cot \theta$, $\sec \theta$, $\csc \theta$, are defined by

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \csc \theta = \frac{1}{\sin \theta}.$$

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.

Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by


$$\cos \theta = x \qquad \sin \theta = y.$$

Definition (additional trigonometric functions)

The functions tangent, **cotangent**, secant and cosecant of the angle θ , denoted by $\tan \theta$, $\cot \theta$, $\sec \theta$, $\csc \theta$, are defined by

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \csc \theta = \frac{1}{\sin \theta}.$$

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.

Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by


$$\cos \theta = x \qquad \sin \theta = y.$$

Definition (additional trigonometric functions)

The functions tangent, cotangent, **secant** and cosecant of the angle θ , denoted by $\tan \theta$, $\cot \theta$, $\sec \theta$, $\csc \theta$, are defined by

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta} \qquad \text{sec } \theta = \frac{1}{\cos \theta} \qquad \csc \theta = \frac{1}{\sin \theta}.$$

Definition of the trigonometric functions



- For an angle-measure θ we selected geometric angle with initial arm on x axis and terminal arm selected by traveling θ units on the unit circle.
- Let (x, y) be the intersection of the terminal arm of the geometric angle with the unit circle.

Definition (sin and cos)

The sine and cosine functions of the angle θ , denoted by $\sin \theta$ and $\cos \theta$, are defined by

$$\cos \theta = x \qquad \sin \theta = y.$$


Definition (additional trigonometric functions)

The functions tangent, cotangent, secant and **cosecant** of the angle θ , denoted by $\tan \theta$, $\cot \theta$, $\sec \theta$, $\csc \theta$, are defined by

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \qquad \cot \theta = \frac{\cos \theta}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \csc \theta = \frac{1}{\sin \theta}.$$

Definition (Similar triangles)


We say that a triangle $\triangle ABC$ is similar to a triangle $\triangle A'B'C'$ if the two triangles have equal angles.



Definition (Similar triangles)

We say that a triangle $\triangle ABC$ is similar to a triangle $\triangle A'B'C'$ if the two triangles have equal angles.


- The equal angles are assumed given in the same order for both triangles, that is, $\angle ABC = \angle A'B'C'$, $\angle BCA = \angle B'C'A'$, $\angle CAB = \angle C'A'B'$.



Definition (Similar triangles)

We say that a triangle $\triangle ABC$ is similar to a triangle $\triangle A'B'C'$ if the two triangles have equal angles.


- The equal angles are assumed given in the same order for both triangles, that is, $\angle ABC = \angle A'B'C'$, $\angle BCA = \angle B'C'A'$, $\angle CAB = \angle C'A'B'$.



Definition (Similar triangles)

We say that a triangle $\triangle ABC$ is similar to a triangle $\triangle A'B'C'$ if the two triangles have equal angles.


- The equal angles are assumed given in the same order for both triangles, that is, $\angle ABC = \angle A'B'C'$, $\angle BCA = \angle B'C'A'$, $\angle CAB = \angle C'A'B'$.



Definition (Similar triangles)

We say that a triangle $\triangle ABC$ is similar to a triangle $\triangle A'B'C'$ if the two triangles have equal angles.

- The equal angles are assumed given in the same order for both triangles, that is, $\angle ABC = \angle A'B'C'$, $\angle BCA = \angle B'C'A'$, $\angle CAB = \angle C'A'B'$.




The following statement is proved in the subject of Euclidean (planar) geometry.

Theorem (Similar triangles have equal side ratios)

Let $\triangle ABC$ and $\triangle A'B'C'$ be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.

Theorem (Similar triangles have equal side ratios)

Let $\triangle ABC$ and $\triangle A'B'C'$ be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.

Theorem (Similar triangles have equal side ratios)

Let $\triangle ABC$ and $\triangle A'B'C'$ be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.



Theorem (Similar triangles have equal side ratios)

Let $\triangle ABC$ and $\triangle A'B'C'$ be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$





Trigonometric Functions and Right Angle Triangles

	
$\cos \theta$ $\sin \theta$ $\tan \theta$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.

Trigonometric Functions and Right Angle Triangles

 <p> $\cos \theta = \frac{x}{r}$ $\sec \theta$ $\sin \theta$ $\csc \theta$ $\tan \theta$ $\cot \theta$ </p>	<p>$0 < \theta < \frac{\pi}{2}$</p>  <p> $\cos \theta$ $\sec \theta$ $\sin \theta$ $\csc \theta$ $\tan \theta$ $\cot \theta$ </p>
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta$ $\csc \theta$ $\cot \theta$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

 <p>Diagram showing an angle θ in standard position. The terminal arm passes through the point (x, y). The distance from the origin to the point is r. The x-axis is highlighted in green, and the y-axis is highlighted in blue. A right angle is shown at the origin between the axes.</p>	 <p>Diagram showing a right triangle for an acute angle θ. The angle θ is at the bottom-left vertex. The side adjacent to θ is the horizontal base, the side opposite is the vertical height, and the hypotenuse is the slanted side. A right angle is shown at the bottom-right vertex.</p>
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta$ $\csc \theta$ $\cot \theta$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta$ $\csc \theta$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$0 < \theta < \frac{\pi}{2}$ $\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

 $\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$ $\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$ <p style="text-align: center;">All angles</p>	 $0 < \theta < \frac{\pi}{2}$ $\cos \theta$ $\sin \theta$ $\tan \theta$ $\sec \theta$ $\csc \theta$ $\cot \theta$ <p style="text-align: center;">Acute angles</p>
---	---



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of **acute θ (between 0 and $\frac{\pi}{2}$)** can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of **sides of right angle triangle** with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta$ $\tan \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$ $\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$ $\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles


- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles

- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Example




³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example




³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

hypotenuse = ?

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2}$$

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25}$$

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

$$\sin \theta = \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = ? \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = \frac{3}{5} \quad \cos \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = \frac{3}{5} \quad \cos \theta = ? \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \\ \csc \theta = & \sec \theta = & \cot \theta = \end{array}$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = \frac{3}{5} \quad \cos \theta = \frac{4}{5} \quad \tan \theta = ?$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = \frac{3}{5} \quad \cos \theta = \frac{4}{5} \quad \tan \theta = \frac{3}{4}$$

$$\csc \theta = \quad \sec \theta = \quad \cot \theta =$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\sin \theta = \frac{3}{5} \quad \cos \theta = \frac{4}{5} \quad \tan \theta = \frac{3}{4}$$

$$\text{csc } \theta = ? \quad \sec \theta = \quad \cot \theta =$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \text{csc } \theta = \frac{5}{3} & \sec \theta = & \cot \theta = \end{array}$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \csc \theta = \frac{5}{3} & \sec \theta = ? & \cot \theta = \end{array}$$

Example



³ Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \csc \theta = \frac{5}{3} & \sec \theta = \frac{5}{4} & \cot \theta = \end{array}$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .


To find the trigonometric functions, we need to know the length of the hypotenuse.

$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \csc \theta = \frac{5}{3} & \sec \theta = \frac{5}{4} & \cot \theta = ? \end{array}$$

Example



Let the angle θ be as indicated in the figure. Find the values of the six trigonometric functions of θ .

To find the trigonometric functions, we need to know the length of the hypotenuse.


$$\text{hypotenuse} = \sqrt{4^2 + 3^2} = \sqrt{25} = 5.$$

Using the right angle triangle ratio interpretations of the trig functions, we can compute:

$$\begin{array}{lll} \sin \theta = \frac{3}{5} & \cos \theta = \frac{4}{5} & \tan \theta = \frac{3}{4} \\ \csc \theta = \frac{5}{3} & \sec \theta = \frac{5}{4} & \cot \theta = \frac{4}{3} \end{array}$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.

$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.

$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = ?$, so $x = ?$.


$$\sin \theta = \quad \quad \tan \theta =$$

$$\csc \theta = \quad \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = ? \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = ?$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = ? \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = ?$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta = ?$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .




- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.

$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta = \frac{2}{\sqrt{21}}$$



Proposition

The angles of every triangle sum up to $\pi = 180^\circ$.

In other words, if α, β, γ are the angles indicated in the figure, then we have:

$$\alpha + \beta + \gamma = 180^\circ.$$


Example

Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

Example

Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle,



Example

Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle, adjacent side of length **1**.




Example

Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.



Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$\begin{aligned} 45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ \end{aligned}$$

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$\begin{aligned} 45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ = ? \end{aligned}$$

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$\begin{aligned} 45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ = 45^\circ. \end{aligned}$$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles \Rightarrow is **isosceles**

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$\begin{aligned}45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ = 45^\circ.\end{aligned}$$

- Triangle has two equal angles \Rightarrow is **isosceles (has two equal sides)**.

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1

Example




Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$\begin{aligned} 45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ = 45^\circ. \end{aligned}$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = ?

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.


- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$


- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

• $\sin 45^\circ = ?$

$\cos 45^\circ = ?$

$\tan 45^\circ = ?$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

- $\sin 45^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{\sqrt{2}}{2}$ $\cos 45^\circ = ?$

$$\tan 45^\circ = ?$$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$


- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

- $\sin 45^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{\sqrt{2}}{2}$

$$\cos 45^\circ = ?$$

$$\tan 45^\circ = ?$$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$


- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

$$\sin 45^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{\sqrt{2}}{2}$$

$$\cos 45^\circ = \frac{\text{adj}}{\text{hyp}} = \frac{\sqrt{2}}{2}$$

$$\tan 45^\circ = ?$$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :


$$\begin{aligned} 45^\circ + 90^\circ + \gamma &= 180^\circ \\ \gamma &= 180^\circ - 90^\circ - 45^\circ = 45^\circ. \end{aligned}$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.

$$\sin 45^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{\sqrt{2}}{2} \qquad \cos 45^\circ = \frac{\text{adj}}{\text{hyp}} = \frac{\sqrt{2}}{2}$$

$$\tan 45^\circ = ?$$

Example



Find the values of $\sin 45^\circ$, $\cos 45^\circ$, $\tan 45^\circ$.

- Draw the 45° angle in right angle triangle, adjacent side of length 1.
- Let γ be the angle indicated on the plot.
- Angles in triangle sum to 180° :

$$45^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 45^\circ = 45^\circ.$$

- Triangle has two equal angles \Rightarrow is isosceles (has two equal sides).
- \Rightarrow Opposite leg: length 1 \Rightarrow length(hyp) = $\sqrt{1^2 + 1^2} = \sqrt{2}$.


$$\sin 45^\circ = \frac{\text{opp}}{\text{hyp}} = \frac{\sqrt{2}}{2} \qquad \cos 45^\circ = \frac{\text{adj}}{\text{hyp}} = \frac{\sqrt{2}}{2}$$

$$\tan 45^\circ = \frac{\text{opp}}{\text{adj}} = \frac{1}{1} = 1.$$

Example

Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated:


Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ .


Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ .


Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ .

Example




Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles **in \triangle** sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

Example




Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$, $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle **sum to 180°** :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

Example




Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ$$

Example




Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = ?$$

Example




Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$.

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$.

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$.

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$.

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$.

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.


Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$)

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :


$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB|$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :


$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :


$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \bigg| \quad \text{Pythagorean theorem}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$


Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

$$= \sqrt{2^2 - 1^2}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$, $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles $60^\circ, 90^\circ, \gamma$. Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$


$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$\begin{aligned} |CH| &= \sqrt{|AC|^2 - |AH|^2} && \text{Pythagorean theorem} \\ &= \sqrt{2^2 - 1^2} \end{aligned}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$


Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$


Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = ?$$

$$\cos 60^\circ = ?$$

$$\tan 60^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = ?$$

$$\tan 60^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = ?$$

$$\tan 60^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$

$$\tan 60^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$

$$\tan 60^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$


$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$

$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = ?$$

$$\cos 30^\circ = ?$$

$$\tan 30^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = ?$$

$$\tan 30^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = ?$$

$$\tan 30^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\tan 30^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad | \text{ Pythagorean theorem}$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\tan 30^\circ = ?$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles
 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has
 three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\cos 60^\circ = \frac{1}{2}$$


$$\tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}$$

$$\sin 30^\circ = \frac{1}{2}$$

$$\cos 30^\circ = \frac{\sqrt{3}}{2}$$

$$\tan 30^\circ = \frac{1}{\sqrt{3}}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore


$$|AC| = |AB| = 1 + 1 = 2$$

$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\begin{aligned} \sin 60^\circ &= \frac{\sqrt{3}}{2} & \cos 60^\circ &= \frac{1}{2} & \tan 60^\circ &= \frac{\sqrt{3}}{1} = \sqrt{3} \\ \sin 30^\circ &= \frac{1}{2} & \cos 30^\circ &= \frac{\sqrt{3}}{2} & \tan 30^\circ &= \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}. \end{aligned}$$

Example



Find the values of $\sin 60^\circ$, $\cos 60^\circ$, $\tan 60^\circ$,
 $\sin 30^\circ$, $\cos 30^\circ$, $\tan 30^\circ$.

Construct a right angled $\triangle AHC$ as indicated: angles 60° , 90° , γ . Angles in \triangle sum to 180° :

$$60^\circ + 90^\circ + \gamma = 180^\circ$$

$$\gamma = 180^\circ - 90^\circ - 60^\circ = 30^\circ.$$

Construct $\triangle HBC$ as indicated so that $\triangle HBC \cong \triangle HAC$. $\triangle ABC$ has three equal angles ($= 60^\circ$) \Rightarrow its sides are of equal length. Therefore

$$|AC| = |AB| = 1 + 1 = 2$$


$$|CH| = \sqrt{|AC|^2 - |AH|^2} \quad \left| \text{Pythagorean theorem} \right.$$

$$= \sqrt{2^2 - 1^2} = \sqrt{3}$$

$$\begin{aligned} \sin 60^\circ &= \frac{\sqrt{3}}{2} & \cos 60^\circ &= \frac{1}{2} & \tan 60^\circ &= \frac{\sqrt{3}}{1} = \sqrt{3} \\ \sin 30^\circ &= \frac{1}{2} & \cos 30^\circ &= \frac{\sqrt{3}}{2} & \tan 30^\circ &= \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}. \end{aligned}$$

Observation

- *If the hypotenuse of a right angle triangle is twice larger than one of the sides, then the angle opposite to that side is 30° .*
- *Conversely, in a right angle triangle with angle 30° , the hypotenuse is twice longer than the shorter of the two legs.*




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.




To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.






To compute trigonometric functions from obtuse ($> 90^\circ$) or negative angles, we can use the following visual aid.

Definition (Reference Angle)

Let θ be an angle in standard position. Its reference angle is the acute positive angle formed by the terminal arm and the x axis.

The computation of the reference angle α depends on the quadrant.


<p>Quadrant II</p>  <p>$\alpha = \pi - \theta$ $\alpha = 180^\circ - \theta$</p>	<p>Quadrant III</p>  <p>$\alpha = \theta - \pi$ $\alpha = \theta - 180^\circ$</p>	<p>Quadrant IV</p>  <p>$\alpha = 2\pi - \theta$ $\alpha = 360^\circ - \theta$</p>
--	--	---

Observation

One can find the value of a trigonometric function of θ as follows.

- *Find the reference angle α associated to θ .*
- *Find the trig function of α .*
- *Use the quadrant in which θ lies to affix an appropriate sign to the function value.*

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = ?$$

$$\csc\left(\frac{2\pi}{3}\right) =$$


$$\cos\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) =$$


$$\cos\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = ?$$

$$\csc\left(\frac{2\pi}{3}\right) = \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2}$$

$$\cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) = \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = ?$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} \\ \csc\left(\frac{2\pi}{3}\right) &= & \sec\left(\frac{2\pi}{3}\right) &= \end{aligned}$$

$$\begin{aligned} \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \cot\left(\frac{2\pi}{3}\right) &= \end{aligned}$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\text{csc}\left(\frac{2\pi}{3}\right) = ? \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\text{csc}\left(\frac{2\pi}{3}\right) = \frac{2}{\sqrt{3}} \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) = \frac{2}{\sqrt{3}} \quad \sec\left(\frac{2\pi}{3}\right) = ?$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= \end{aligned}$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= ? \end{aligned}$$


Example



Find the exact values of the trigonometric functions of


$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= -\frac{1}{\sqrt{3}} \end{aligned}$$




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	?										
cos	?										




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0										
cos	1										




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	?									
cos	1	?									




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$									
cos	1	$\frac{\sqrt{3}}{2}$									




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$?								
cos	1	$\frac{\sqrt{3}}{2}$?								




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$								
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$								




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$?							
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$?							



- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$							
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$							




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Quadrant III


Quadrant IV

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$?						
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$?						




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1						
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0						




- One only needs to memorize sines and cosines in Quadrant I and on the axes.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	?					
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	?					




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of **the reference angle**

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	?					
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	?					




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of **the reference angle**

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	?					
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	?					




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the **sine/cosine of the reference angle**

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$					
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$					




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and **adjusting the sign according to the quadrant.**

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$					
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$					




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$?				
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$?				




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$				
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$				




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$?			
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$?			




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$			
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$			




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$?		
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$?		




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0		
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1		




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0		
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1		




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0		
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1		




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	?	
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	?	




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	0	




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	0	




- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	0	



- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	?
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	0	?



- One only needs to memorize sines and cosines in Quadrant I and on the axes.
- The remaining sines and cosines are extracted by
 - taking the sine/cosine of the reference angle
 - and adjusting the sign according to the quadrant.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	0
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1	0	1

Geometric interpretation of all trigonometric functions




Fix unit circle, center O , coordinates $(0, 0)$.

Geometric interpretation of all trigonometric functions




Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$.

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A .

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$\sin \theta$

$\cos \theta$


$\tan \theta$

$\cot \theta$

$\sec \theta$

$\csc \theta$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

Geometric interpretation of all trigonometric functions



$$\beta = ?$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

Geometric interpretation of all trigonometric functions



$\angle OED = ?$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$. Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OB|}{|AC|} = \frac{|OB|}{|AC|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|} = \frac{1}{|DE|} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$


$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$




- Positive angles are obtained by rotating counterclockwise.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.

$$\begin{aligned} \sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y} \end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.
- \sin is an **odd function**.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.
- \sin is an odd function.
- \cos is an even function.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- 2π represents a full rotation.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .
- $\sin(\theta + 2\pi) = \sin \theta$.
- $\cos(\theta + 2\pi) = \cos \theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .
- $\sin(\theta + 2\pi) = \sin \theta$.
- $\cos(\theta + 2\pi) = \cos \theta$.
- We say \sin and \cos are 2π -periodic.

$$\begin{aligned} \sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y} \end{aligned}$$

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.


- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.

Trigonometric Identities

Definition (Trigonometric Identity)


A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.
- For example, $\frac{\sin \theta}{\sin \theta} = 1$ is considered a valid trigonometric identity, although, when considered as a function, the left hand side is not defined for $\theta \neq 0$.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- $\csc \theta = \frac{1}{\sin \theta}$
- $\sec \theta = \frac{1}{\cos \theta}$
- $\cot \theta = \frac{1}{\tan \theta}$
- $\tan \theta = \frac{\sin \theta}{\cos \theta}$
- $\cot \theta = \frac{\cos \theta}{\sin \theta}$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\sin^2 \theta + \cos^2 \theta$$

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$

Therefore $\sin^2 \theta + \cos^2 \theta = 1$.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\sin^2 \theta + \cos^2 \theta = 1$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta} \\ \tan^2 \theta + 1 &= \sec^2 \theta\end{aligned}$$