

# Calculus I

## Maxima and minima over closed intervals

Todor Milev

2019

# Outline

- 1 One Variable Optimization Problems
  - The Closed Interval Method
  - Solving One Variable Optimization Problems

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

Fermat's Theorem suggests that we should look at three types of points to find local maxima and minima:

- 1 Points  $c$  for which  $f'(c) = 0$ .
- 2 Points  $c$  for which  $f'(c)$  doesn't exist.
- 3 Points  $c$  at ends of intervals where  $f$  is defined. Here, we need also that  $f$  be defined at  $c$ .

Fermat's Theorem suggests that we should look at three types of points to find local maxima and minima:

- 1 Points  $c$  for which  $f'(c) = 0$ .
- 2 Points  $c$  for which  $f'(c)$  doesn't exist.
- 3 Points  $c$  at ends of intervals where  $f$  is defined. Here, we need also that  $f$  be defined at  $c$ .

### Definition (Critical Number)

A critical number of a function  $f$  is a number  $c$  in the domain of  $f$  such that either  $f'(c) = 0$  or  $f'(c)$  doesn't exist.

Fermat's Theorem suggests that we should look at three types of points to find local maxima and minima:

- 1 Points  $c$  for which  $f'(c) = 0$ .
- 2 Points  $c$  for which  $f'(c)$  doesn't exist.
- 3 Points  $c$  at ends of intervals where  $f$  is defined. Here, we need also that  $f$  be defined at  $c$ .

### Definition (Critical Number)

A critical number of a function  $f$  is a number  $c$  in the domain of  $f$  such that either  $f'(c) = 0$  or  $f'(c)$  doesn't exist.

Fermat's Theorem says that if  $f$  has a local maximum or minimum at  $c$ , and  $c$  is not an endpoint, then  $c$  is a critical number for  $f$ .

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$\begin{aligned} f(x) &= x^{\frac{1}{4}} (4 - x^2) \\ &= 4x^{\frac{1}{4}} - x^{\frac{9}{4}} \end{aligned}$$



## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

Set  $f'(x) = 0$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$



## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

• Critical numbers occur:

- ① Where  $f'(x)$  isn't defined:
- ② Where  $f'(x) = 0$ :

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

- Critical numbers occur:

① Where  $f'(x)$  isn't defined:

② Where  $f'(x) = 0$ :

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

- Critical numbers occur:

① Where  $f'(x)$  isn't defined: 0.

② Where  $f'(x) = 0$ :

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

- Critical numbers occur:

① Where  $f'(x)$  isn't defined: 0.

② Where  $f'(x) = 0$ :

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

• Critical numbers occur:

① Where  $f'(x)$  isn't defined: 0.

② Where  $f'(x) = 0$ :  $\frac{2}{3}$  and  $-\frac{2}{3}$ .

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

● Critical numbers occur:

- ① Where  $f'(x)$  isn't defined: 0.
- ② Where  $f'(x) = 0$ :  $\frac{2}{3}$  and  $-\frac{2}{3}$ .

●  $f$  isn't defined at  $-\frac{2}{3}$ .

## Example

Find the critical numbers of  $f(x) = x^{\frac{1}{4}} (4 - x^2)$ .

$$f(x) = x^{\frac{1}{4}} (4 - x^2)$$

$$= 4x^{\frac{1}{4}} - x^{\frac{9}{4}}$$

$$\text{so } f'(x) = x^{-\frac{3}{4}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{1}{x^{\frac{3}{4}}} - \frac{9}{4}x^{\frac{5}{4}}$$

$$= \frac{4 - 9x^2}{4x^{\frac{3}{4}}}$$

$$\text{Set } f'(x) = 0$$

$$\frac{4 - 9x^2}{4x^{\frac{3}{4}}} = 0$$

$$4 - 9x^2 = 0$$

$$(2 - 3x)(2 + 3x) = 0$$

$$x = \pm \frac{2}{3}$$

- Critical numbers occur:

① Where  $f'(x)$  isn't defined: 0.

② Where  $f'(x) = 0$ :  $\frac{2}{3}$  and  $-\frac{2}{3}$ .

- $f$  isn't defined at  $-\frac{2}{3}$ . Therefore the critical numbers are 0 and  $\frac{2}{3}$ .

# The Closed Interval Method

We know from the Extreme Value Theorem that a continuous function attains its maximum and minimum on a closed interval  $[a, b]$ . The maximum might occur at an endpoint. The minimum might occur at an endpoint.



# The Closed Interval Method

We know from the Extreme Value Theorem that a continuous function attains its maximum and minimum on a closed interval  $[a, b]$ . The maximum might occur at an endpoint. The minimum might occur at an endpoint.


To find the maximum and minimum values of a continuous function  $f$  on a closed interval  $[a, b]$ :

- 1 Find the values of  $f$  at the critical numbers of  $f$  in  $[a, b]$ .
  - Find the values  $c$  with  $f'(c) = 0$ .
  - Find the values  $c$  where  $f'$  does not exist.
- 2 Find the values of  $f$  at the endpoints  $a$  and  $b$ .
- 3 The maximum of  $f$  is maximum of the preceding values; the minimum value is the minimum.

## Example

Find the maximum and minimum values of the function


$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .




$$f'(x) = -3x^2 + 4x + 4$$

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .




$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$f'(x) = -3x^2 + 4x + 4$$


$$= (-3x - 2)(x - 2)$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\&= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or **2**.

Need to check:

- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .


$x$	$f(x)$
<b>2</b>	



## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
2	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	
2	
3	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	?
2	
3	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	
3	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\&= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	?
3	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\&= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	?

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:

- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .


$x$	$f(x)$
1	0
2	3
3	-2



## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned}f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2)\end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	-2

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:

- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .


$x$	$f(x)$
1	0
2	3
3	-2

Maximum on  $[1, 3]$ : Minimum on  $[1, 3]$ :

## Example

Find the maximum and minimum values of the function

$f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	-2

Maximum on  $[1, 3]$ : 3. Minimum on  $[1, 3]$ :

## Example

Find the maximum and minimum values of the function  $f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .

Need to check:


- 1 The critical numbers of  $f$  in  $[a, b]$ .
- 2 The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	-2

Maximum on  $[1, 3]$ : 3. Minimum on  $[1, 3]$ :

## Example

Find the maximum and minimum values of the function  $f(x) = -x^3 + 2x^2 + 4x - 5$  on the interval  $[1, 3]$ .



$$\begin{aligned} f'(x) &= -3x^2 + 4x + 4 \\ &= (-3x - 2)(x - 2) \end{aligned}$$

If  $f'(x) = 0$ ,  $x = -\frac{2}{3}$  or  $2$ .


Need to check:

- ① The critical numbers of  $f$  in  $[a, b]$ .
- ② The endpoints  $a$  and  $b$ .

$x$	$f(x)$
1	0
2	3
3	-2


Maximum on  $[1, 3]$ : 3. Minimum on  $[1, 3]$ : -2.

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.


## Example



$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right)$$

Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

## Example




Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2})$$




## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} \left( e^{-x^2} \right)$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1) e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\ &= ? \cdot e^{-x^2} + (x + 1) ?\end{aligned}$$


## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1) e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\ &= 1 \cdot e^{-x^2} + (x + 1)?\end{aligned}$$


## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\ &= 1 \cdot e^{-x^2} + (x + 1) \text{?}\end{aligned}$$


## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\ &= 1 \cdot e^{-x^2} + (x + 1) e^{-x^2} (-x^2)'\end{aligned}$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (?)
 \end{aligned}$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x)
 \end{aligned}$$

# Example




Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x) \\
 &= (1 + (x + 1)(-2x)) e^{-x^2}
 \end{aligned}$$




# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x) \\
 &= (1 + (x + 1)(-2x)) e^{-x^2}
 \end{aligned}$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x) \\
 &= (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}
 \end{aligned}$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x) \\
 &= (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}
 \end{aligned}$$


# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\begin{aligned}
 \frac{d}{dx} \left( (x + 1)e^{-x^2} \right) &= \frac{d}{dx} (x + 1)e^{-x^2} + (x + 1) \frac{d}{dx} (e^{-x^2}) \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-x^2)' \\
 &= 1 \cdot e^{-x^2} + (x + 1)e^{-x^2} (-2x) \\
 &= (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}
 \end{aligned}$$


## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

## Example




Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.


$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

Div. by  $e^{-x^2}$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$


Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

Div. by  $e^{-x^2} \neq 0$



# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.


$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$\begin{aligned} (-2x^2 - 2x + 1)e^{-x^2} &= 0 \\ -2x^2 - 2x + 1 &= 0 \end{aligned}$$

$$\left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$


Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$\begin{aligned} (-2x^2 - 2x + 1)e^{-x^2} &= 0 \\ -2x^2 - 2x + 1 &= 0 \end{aligned}$$

$$\left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = ?$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$


Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$\begin{aligned} (-2x^2 - 2x + 1)e^{-x^2} &= 0 \\ -2x^2 - 2x + 1 &= 0 \end{aligned}$$

Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$


$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

$$= \frac{2 \pm \sqrt{12}}{-4}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$


Div. by  $e^{-x^2} \neq 0$

$$x_1, x_2 = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)}$$

$$= \frac{2 \pm \sqrt{12}}{-4}$$



# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$\begin{aligned} x_1, x_2 &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)} \\ &= \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} \end{aligned}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$\begin{aligned} x_1, x_2 &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)} \\ &= \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} \end{aligned}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$\left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$-2x^2 - 2x + 1 = 0$$

$$\begin{aligned} x_1, x_2 &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)} \\ &= \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2} \end{aligned}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0$$

$$-2x^2 - 2x + 1 = 0$$

Div. by  $e^{-x^2} \neq 0$

$$\begin{aligned} x_1, x_2 &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(-2) \cdot 1}}{2(-2)} \\ &= \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2} \end{aligned}$$

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.


$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	
$\frac{-1 - \sqrt{3}}{2}$	
$\frac{-1 + \sqrt{3}}{2}$	
$5$	

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at **the endpoints** and the critical points:

$x$	$f(x)$
<b><math>-5</math></b>	
$\frac{-1 - \sqrt{3}}{2}$	
$\frac{-1 + \sqrt{3}}{2}$	
<b><math>5</math></b>	

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . Use the given plot.

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$


$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	
$\frac{-1 - \sqrt{3}}{2}$	
$\frac{-1 + \sqrt{3}}{2}$	
$5$	



# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . **Use the given plot.**

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	close to 0 from plot
$\frac{-1 - \sqrt{3}}{2}$	
$\frac{-1 + \sqrt{3}}{2}$	
$5$	

# Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . **Use the given plot.**

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	close to 0 from plot
$\frac{-1 - \sqrt{3}}{2}$	<b>negative, min from plot</b>
$\frac{-1 + \sqrt{3}}{2}$	
$5$	

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . **Use the given plot.**

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	close to 0 from plot
$\frac{-1 - \sqrt{3}}{2}$	negative, min from plot
$\frac{-1 + \sqrt{3}}{2}$	<b>positive, max from plot</b>
$5$	

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . **Use the given plot.**

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :


$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	close to 0 from plot
$\frac{-1-\sqrt{3}}{2}$	negative, min from plot
$\frac{-1+\sqrt{3}}{2}$	positive, max from plot
$5$	close to 0 from plot

## Example



Find the value of  $x$  for which  $f(x) = (x + 1)e^{-x^2}$  attains its maximum in the interval  $[-5, 5]$ . **Use the given plot.**

$$\frac{d}{dx} \left( (x + 1)e^{-x^2} \right) = (1 + (x + 1)(-2x)) e^{-x^2} = (-2x^2 - 2x + 1) e^{-x^2}$$

Find critical points: set  $f'(x) = 0$  and solve for  $x$ :

$$(-2x^2 - 2x + 1)e^{-x^2} = 0 \quad \left| \text{Div. by } e^{-x^2} \neq 0 \right.$$

$$x_1, x_2 = \frac{2 \pm \sqrt{12}}{-4} = \frac{-2 \mp 2\sqrt{3}}{4} = \frac{-1 \pm \sqrt{3}}{2}$$

Compare the values of  $f$  at the endpoints and the critical points:

$x$	$f(x)$
$-5$	close to 0 from plot
$\frac{-1 - \sqrt{3}}{2}$	negative, min from plot
<b>Final answer: <math>\frac{-1 + \sqrt{3}}{2}</math></b>	<b>positive, max from plot</b>
$5$	close to 0 from plot

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in **the above form**.



# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

- 1 Draw a picture of the problem. Assign variable names to the involved quantities.

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

- 1 Draw a picture of the problem. Assign variable names to the involved quantities. Determine **which quantity** is being maximized/minimized.

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

- 1 Draw a picture of the problem. Assign variable names to the involved quantities. Determine which quantity is being maximized/minimized.
- 2 Express all involved quantities in terms of **only one of them**.

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

- 1 Draw a picture of the problem. Assign variable names to the involved quantities. Determine which quantity is being maximized/minimized.
- 2 Express all involved quantities in terms of only one of them. If you cannot do that then the problem is not in one variable (i.e., lies outside of the scope of Calculus I).

# One variable optimization Problems

The problem of finding minimum/maximum of a differentiable one-variable function often arises in practice.

## Problem (One variable optimization problem statement)

*Given a function  $f(x)$ , find the maximum and/or the minimum of  $f(x)$ , and the values of  $x$  for which the minima/maxima are achieved.*

Optimization problems are usually not formulated directly in the above form. Solving an optimization problem involves the following steps.

- 1 Draw a picture of the problem. Assign variable names to the involved quantities. Determine which quantity is being maximized/minimized.
- 2 Express all involved quantities in terms of only one of them. If you cannot do that then the problem is not in one variable (i.e., lies outside of the scope of Calculus I).
- 3 Use the closed interval method to find the maximum/minimum value of the desired quantity.


## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?




$$\text{Area} = 200 \cdot 2000 = 400,000\text{ft}^2$$



## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 300 \cdot 1800 = 540,000\text{ft}^2$$

## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 500 \cdot 1400 = 700,000\text{ft}^2$$

## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 600 \cdot 1200 = 720,000\text{ft}^2$$

## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 800 \cdot 800 = 640,000\text{ft}^2$$

## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 900 \cdot 600 = 540,000\text{ft}^2$$

## Example


A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = 1100 \cdot 200 = 220,000\text{ft}^2$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?




Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$\text{Area} = A = xy$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.


$$2x + y = 2400$$

$$\text{Area} = A = xy$$



## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.


$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$\text{Area} = A = xy$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$


$$y = 2400 - 2x$$

$$A = xy$$

$$\text{Area} = A = xy$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$


$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$\text{Area} = A = xy$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$


$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$


$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$


Notice that  $0 \leq x \leq 1200$ .

Maximize the function  $A(x)$ :

$$A'(x) = ?$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$


Notice that  $0 \leq x \leq 1200$ .

Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .

Maximize the function  $A(x)$ :


$$A'(x) = 2400 - 4x$$

Critical number:  $x = ?$



## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



$$\text{Area} = A = xy$$

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	
600	
1200	

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	?
600	
1200	

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	
1200	

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	?
1200	

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	720,000
1200	

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	720,000
1200	?

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .


Maximize the function  $A(x)$ :

$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	720,000
1200	0

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .

Maximize the function  $A(x)$ :


$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .



## Example

A farmer has 2400 ft of fencing and wants to fence off a rectangular field that borders a straight river. He doesn't need to put fencing along the river. What are the dimensions of the field with the largest area?



Area =  $A = xy$

$x$	$A(x)$
0	0
600	720,000
1200	0

Let  $x$  and  $y$  denote the depth and width of the rectangle (in feet). Let  $A$  be its area.

$$2x + y = 2400$$

$$y = 2400 - 2x$$

$$A = xy = x(2400 - 2x)$$

$$= 2400x - 2x^2$$

Notice that  $0 \leq x \leq 1200$ .

Maximize the function  $A(x)$ :


$$A'(x) = 2400 - 4x$$

Critical number:  $x = 600$ .

Therefore the maximum area occurs when  $x = 600\text{ft}$  and  $y = 1200\text{ft}$ .


## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



## Example


Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .






## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



## Example


Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin.

## Example


Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle.

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= ? \cdot ?$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height} \\ = ? \cdot y$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.


$$A = \text{base} \cdot \text{height}$$

$$= ? \cdot y$$



## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$\begin{aligned} A &= \text{base} \cdot \text{height} \\ &= 2x \cdot y \end{aligned}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .




Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$\begin{aligned} A &= \text{base} \cdot \text{height} \\ &= 2x \cdot y = 2x \cdot ? \end{aligned}$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$


$$= 2x \cdot y = 2x \cdot ?$$

To eliminate  $y$ , use that  
 $(x, y)$  lies on the semicircle.

$$x^2 + y^2 = r^2$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$


$$= 2x \cdot y = 2x \cdot ?$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$x^2 + y^2 = r^2$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$


$$= 2x \cdot y = 2x \cdot ?$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.


$$\begin{aligned} A &= \text{base} \cdot \text{height} \\ &= 2x \cdot y = 2x \cdot ? \end{aligned}$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$\begin{aligned} y^2 &= r^2 - x^2 \\ y &= \pm \sqrt{r^2 - x^2} \end{aligned}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.


$$\begin{aligned} A &= \text{base} \cdot \text{height} \\ &= 2x \cdot y = 2x \cdot ? \end{aligned}$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$\begin{aligned} y^2 &= r^2 - x^2 \\ y &= \pm \sqrt{r^2 - x^2} \quad | \quad y > 0 \end{aligned}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot ?$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.


$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2} \mid y > 0$$



## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$A' = ?$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$A' = 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}}$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} \end{aligned}$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$A' = 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}}$$

$$= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}}$$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

Critical numbers:  $x = ?$


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

Critical numbers:  $x = \frac{r}{\sqrt{2}}$  and  $r$ .

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.


$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$



## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

Critical numbers:  $x = \frac{r}{\sqrt{2}}$  and  $r$ .


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

Critical numbers:  $x = \frac{r}{\sqrt{2}}$  and  $r$ .


To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

Critical numbers:  $x = \frac{r}{\sqrt{2}}$  and  $r$ .

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.


$$y^2 = r^2 - x^2$$

$$y = \sqrt{r^2 - x^2}$$

We have  $0 \leq x \leq r$  and so the critical numbers together with the endpoints are  $x = 0, \frac{r}{\sqrt{2}}, r$ .

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

$$\text{Critical numbers: } x = \frac{r}{\sqrt{2}} \text{ and } r.$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$


$$y = \sqrt{r^2 - x^2}$$

We have  $0 \leq x \leq r$  and so the critical numbers together with the endpoints are  $x = 0, \frac{r}{\sqrt{2}}, r$ . Since  $A(0) = 0 = A(r)$ , the max is achieved

$$\text{at } x = y = \frac{r}{\sqrt{2}}.$$

## Example

Find the largest possible area of a rectangle inscribed in a semicircle of radius  $r$ .



Let the semicircle have center at the origin. Let  $(x, y)$  -coord. of top right corner of rectangle. Let  $A$  be its area.

$$A = \text{base} \cdot \text{height}$$

$$= 2x \cdot y = 2x \cdot \sqrt{r^2 - x^2}$$

$$\begin{aligned} A' &= 2\sqrt{r^2 - x^2} - \frac{2x^2}{\sqrt{r^2 - x^2}} \\ &= \frac{2(r^2 - x^2)}{\sqrt{r^2 - x^2}} - \frac{2x^2}{\sqrt{r^2 - x^2}} = \frac{2(r^2 - 2x^2)}{\sqrt{r^2 - x^2}} \end{aligned}$$

$$\text{Critical numbers: } x = \frac{r}{\sqrt{2}} \text{ and } r.$$

To eliminate  $y$ , use that  $(x, y)$  lies on the semicircle.

$$y^2 = r^2 - x^2$$



$$y = \sqrt{r^2 - x^2}$$

We have  $0 \leq x \leq r$  and so the critical numbers together with the endpoints are  $x = 0, \frac{r}{\sqrt{2}}, r$ . Since  $A(0) = 0 = A(r)$ , the max is achieved

at  $x = y = \frac{r}{\sqrt{2}}$ . The max area is  $A(\frac{r}{\sqrt{2}}) = 2 \cdot \frac{r}{\sqrt{2}} \sqrt{r^2 - \frac{r^2}{2}} = r^2$ .

## Example



A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



Set  $h$  - cone height,



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.

Set  $h$  - cone height,  $t$  - cone radius.







## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.


Set  $h$  - cone height,  $t$  - cone radius. Then

$V =$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then

$$V = \frac{1}{3}(\text{area cone base})h$$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then

$$V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h.$$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  
 $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge.




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc}AB =$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  
 $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc}AB = r\theta$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  
 $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} =$




## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ .






## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ .





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$h =$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2}$$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2}$$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2} = \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2},$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





Set  $h$  - cone height,  $t$  - cone radius. Then

$V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2} = \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2},$$


and therefore

$$V = \frac{1}{3}\pi t^2 h =$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2} = \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2},$$


and therefore

$$V = \frac{1}{3}\pi t^2 h = \frac{1}{3}\pi \left(\frac{r\theta}{2\pi}\right)^2 \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2}$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2} = \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2},$$

and therefore


$$V = \frac{1}{3}\pi t^2 h = \frac{1}{3}\pi \left(\frac{r\theta}{2\pi}\right)^2 \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2}$$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





Set  $h$  - cone height,  $t$  - cone radius. Then  $V = \frac{1}{3}(\text{area cone base})h = \frac{1}{3}\pi t^2 h$ . Let  $\theta$  - angle of the wedge. Then  $\text{arc } AB = r\theta = \text{perimeter cone base} = 2\pi t$ . Therefore  $t = \frac{r\theta}{2\pi}$ . Then

$$h = \sqrt{r^2 - t^2} = \sqrt{r^2 - \left(\frac{r\theta}{2\pi}\right)^2} = \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2},$$

and therefore

$$\begin{aligned} V &= \frac{1}{3}\pi t^2 h = \frac{1}{3}\pi \left(\frac{r\theta}{2\pi}\right)^2 \frac{r}{2\pi} \sqrt{4\pi^2 - \theta^2} \\ &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}. \end{aligned}$$






## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.

We reduced the problem to: find the maximum of

$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2},$$

as function of  $\theta$  (using the closed interval method).






## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.

We reduced the problem to: find the maximum of

$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad \leq \theta \leq$$

as function of  $\theta$  (using the **closed interval** method).






## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.

We reduced the problem to: find the maximum of


$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

as function of  $\theta$  (using the **closed interval** method).



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





We reduced the problem to: find the maximum of

$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$


as function of  $\theta$  (using the closed interval method).

We need to find the critical points of  $V$ , i.e., the values of  $\theta$  for which  $\frac{dV}{d\theta} = 0$  and the values of  $\theta$  for which  $\frac{dV}{d\theta}$  is not defined.





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} =$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2})$$



## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{-\theta}{\sqrt{4\pi^2 - \theta^2}} \right)
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{-\theta}{\sqrt{4\pi^2 - \theta^2}} \right)
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{-\theta}{\sqrt{4\pi^2 - \theta^2}} \right)
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{1}{2} \frac{\frac{d}{d\theta} (-\theta^2)}{\sqrt{4\pi^2 - \theta^2}} \right)
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{1}{2} \frac{\frac{d}{d\theta}(-\theta^2)}{\sqrt{4\pi^2 - \theta^2}} \right) \\
 &= \left( \frac{r^3}{24\pi^2} \right) \frac{2\theta(4\pi^2 - \theta^2) - \theta^3}{\sqrt{4\pi^2 - \theta^2}}
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{1}{2} \frac{\frac{d}{d\theta}(-\theta^2)}{\sqrt{4\pi^2 - \theta^2}} \right) \\
 &= \left( \frac{r^3}{24\pi^2} \right) \frac{2\theta(4\pi^2 - \theta^2) - \theta^3}{\sqrt{4\pi^2 - \theta^2}}
 \end{aligned}$$

## Example




A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{1}{2} \frac{\frac{d}{d\theta}(-\theta^2)}{\sqrt{4\pi^2 - \theta^2}} \right) \\
 &= \left( \frac{r^3}{24\pi^2} \right) \frac{2\theta(4\pi^2 - \theta^2) - \theta^3}{\sqrt{4\pi^2 - \theta^2}}
 \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




$$\begin{aligned}
 V &= \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, & 0 \leq \theta \leq 2\pi \\
 \frac{dV}{d\theta} &= \left( \frac{r^3}{24\pi^2} \right) \frac{d}{d\theta} (\theta^2) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \frac{d}{d\theta} (\sqrt{4\pi^2 - \theta^2}) \\
 &= \left( \frac{r^3}{24\pi^2} \right) (2\theta) \sqrt{4\pi^2 - \theta^2} \\
 &\quad + \left( \frac{r^3}{24\pi^2} \right) \theta^2 \left( \frac{1}{2} \frac{\frac{d}{d\theta}(-\theta^2)}{\sqrt{4\pi^2 - \theta^2}} \right) \\
 &= \left( \frac{r^3}{24\pi^2} \right) \frac{2\theta(4\pi^2 - \theta^2) - \theta^3}{\sqrt{4\pi^2 - \theta^2}} \\
 &= \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}
 \end{aligned}$$





## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.




$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$



## Example



A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$


$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$

We have that  $\frac{dV}{d\theta} = 0$  when



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$


We have that  $\frac{dV}{d\theta} = 0$  when

$$8\theta\pi^2 - 3\theta^3 = 0$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.





$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$




We have that  $\frac{dV}{d\theta} = 0$  when

$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \end{aligned}$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$




$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$

We have that  $\frac{dV}{d\theta} = 0$  when

$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \\ -3\theta \left( \theta - \sqrt{\frac{8}{3}}\pi \right) \left( \theta + \sqrt{\frac{8}{3}}\pi \right) &= 0. \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$




We have that  $\frac{dV}{d\theta} = 0$  when

$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \\ -3\theta \left( \theta - \sqrt{\frac{8}{3}}\pi \right) \left( \theta + \sqrt{\frac{8}{3}}\pi \right) &= 0. \end{aligned}$$

Therefore  $\theta$  is critical point for  $V$  if  $\theta = 0$ ,  $\theta = \sqrt{\frac{8}{3}}\pi$ ,  
or  $\theta = 2\pi$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$




We have that  $\frac{dV}{d\theta} = 0$  when

$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \\ -3\theta \left( \theta - \sqrt{\frac{8}{3}}\pi \right) \left( \theta + \sqrt{\frac{8}{3}}\pi \right) &= 0. \end{aligned}$$

Therefore  $\theta$  is critical point for  $V$  if  $\theta = 0$ ,  $\theta = \sqrt{\frac{8}{3}}\pi$ ,  
or  $\theta = 2\pi$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$

We have that  $\frac{dV}{d\theta} = 0$  when




$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \\ -3\theta \left( \theta - \sqrt{\frac{8}{3}}\pi \right) \left( \theta + \sqrt{\frac{8}{3}}\pi \right) &= 0. \end{aligned}$$

Therefore  $\theta$  is critical point for  $V$  if  $\theta = 0$ ,  $\theta = \sqrt{\frac{8}{3}}\pi$ ,  
or  $\theta = 2\pi$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}, \quad 0 \leq \theta \leq 2\pi$$

$$\frac{dV}{d\theta} = \left( \frac{r^3}{24\pi^2} \right) \frac{8\theta\pi^2 - 3\theta^3}{\sqrt{4\pi^2 - \theta^2}}$$




We have that  $\frac{dV}{d\theta} = 0$  when

$$\begin{aligned} 8\theta\pi^2 - 3\theta^3 &= 0 \\ \theta(8\pi^2 - 3\theta^2) &= 0 \\ -3\theta \left( \theta - \sqrt{\frac{8}{3}}\pi \right) \left( \theta + \sqrt{\frac{8}{3}}\pi \right) &= 0. \end{aligned}$$

Therefore  $\theta$  is critical point for  $V$  if  $\theta = 0$ ,  $\theta = \sqrt{\frac{8}{3}}\pi$ , or  $\theta = 2\pi$  (note  $\theta = -\sqrt{\frac{8}{3}}\pi$  is outside of the domain of  $V$ ).


# Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



## Example



A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$




Finally, the answer to the problem is

$$V_{\max} = V\left(\sqrt{\frac{8}{3}}\pi\right)$$



## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$

Finally, the answer to the problem is

$$\begin{aligned} V_{\max} &= V\left(\sqrt{\frac{8}{3}}\pi\right) \\ &= \frac{r^3}{24\pi^2} \left(\sqrt{\frac{8}{3}}\pi\right)^2 \sqrt{4\pi^2 - \left(\sqrt{\frac{8}{3}}\pi\right)^2} \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$

Finally, the answer to the problem is

$$\begin{aligned} V_{\max} &= V\left(\sqrt{\frac{8}{3}}\pi\right) \\ &= \frac{r^3}{24\pi^2} \left(\sqrt{\frac{8}{3}}\pi\right)^2 \sqrt{4\pi^2 - \left(\sqrt{\frac{8}{3}}\pi\right)^2} \\ &= \frac{r^3}{9} \pi \sqrt{4 - \frac{8}{3}} \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$

Finally, the answer to the problem is

$$\begin{aligned} V_{\max} &= V\left(\sqrt{\frac{8}{3}}\pi\right) \\ &= \frac{r^3}{24\pi^2} \left(\sqrt{\frac{8}{3}}\pi\right)^2 \sqrt{4\pi^2 - \left(\sqrt{\frac{8}{3}}\pi\right)^2} \\ &= \frac{r^3}{9} \pi \sqrt{4 - \frac{8}{3}} \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.






$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$

Finally, the answer to the problem is

$$\begin{aligned} V_{\max} &= V\left(\sqrt{\frac{8}{3}}\pi\right) \\ &= \frac{r^3}{24\pi^2} \left(\sqrt{\frac{8}{3}}\pi\right)^2 \sqrt{4\pi^2 - \left(\sqrt{\frac{8}{3}}\pi\right)^2} \\ &= \frac{r^3}{9} \pi \sqrt{4 - \frac{8}{3}} \\ &= \pi \frac{r^3}{9} \sqrt{\frac{4}{3}} \end{aligned}$$

## Example

A cone is folded from a wedge-shaped profile of radius  $r$ . Find the maximal possible volume  $V$  of such a cone.



$$V(\theta) = \frac{r^3}{24\pi^2} \theta^2 \sqrt{4\pi^2 - \theta^2}$$

Finally, the answer to the problem is

$$\begin{aligned} V_{\max} &= V\left(\sqrt{\frac{8}{3}}\pi\right) \\ &= \frac{r^3}{24\pi^2} \left(\sqrt{\frac{8}{3}}\pi\right)^2 \sqrt{4\pi^2 - \left(\sqrt{\frac{8}{3}}\pi\right)^2} \\ &= \frac{r^3}{9} \pi \sqrt{4 - \frac{8}{3}} \\ &= \pi \frac{r^3}{9} \sqrt{\frac{4}{3}} = \frac{2\pi r^3}{9\sqrt{3}} \end{aligned}$$