

Calculus II

Power series expansion of logarithms, part 1

Todor Milev

2019

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\ln(1 - x)$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\ln(1 - x) = \int d(\ln(1 - x)) \quad \Bigg| \quad \text{up to const.}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\ln(1 - x) = \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx \quad \Big| \text{ up to const.}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned} \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx \quad \Bigg| \text{ up to const.} \\ &= \int ? \quad dx \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}\ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx \quad \Big| \text{ up to const.} \\ &= \int \left(-\frac{1}{1 - x} \right) dx\end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx \quad \Bigg| \text{ up to const.} \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(? \right) dx
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(? \right) dx && \left| \text{for } |x| < 1 \right.
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right.
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} ?
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} ?
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} \frac{x^n}{n}
 \end{aligned}$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\ln(1 - x) = \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx \quad \left| \text{up to const.} \right.$$

$$= \int \left(-\frac{1}{1 - x} \right) dx$$

$$= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx \quad \left| \text{for } |x| < 1 \right.$$

$$= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C$$

$$= C - \sum_{n=1}^{\infty} \frac{x^n}{n}$$

- To find C , plug in $x = 0$: $C = ?$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} \frac{x^n}{n}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} \frac{x^n}{n}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.
- Therefore the theorem on integrating power series implies that

$$\ln(1 - x) = - \sum_{n=1}^{\infty} \frac{x^n}{n}, \text{ for } |x| < 1.$$

Example

Find a power series for $\ln(1 - x)$ and state its radius of convergence.

$$\begin{aligned}
 \ln(1 - x) &= \int d(\ln(1 - x)) = \int (\ln(1 - x))' dx && \left| \text{up to const.} \right. \\
 &= \int \left(-\frac{1}{1 - x} \right) dx \\
 &= - \int \left(1 + x + x^2 + x^3 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= - \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + C \\
 &= C - \sum_{n=1}^{\infty} \frac{x^n}{n}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.
- Therefore the theorem on integrating power series implies that

$$\ln(1 - x) = - \sum_{n=1}^{\infty} \frac{x^n}{n}, \text{ for } |x| < 1.$$

- By the same theorem, the radius of convergence remains $R = 1$.