

# Calculus II

## Integrals of involving radicals of quadratics

Todor Milev

2019

# Outline

- 1 Integrals of form  $\int R(x, \sqrt{ax^2 + bx + c})dx$ ,  $R$  - rational function
  - Transforming to the forms  $\sqrt{x^2 + 1}$ ,  $\sqrt{-x^2 + 1}$ ,  $\sqrt{x^2 - 1}$
  - Table of Euler and trig substitutions
  - The case  $\sqrt{x^2 + 1}$
  - The case  $\sqrt{-x^2 + 1}$
  - The case  $\sqrt{x^2 - 1}$

# Outline

- 1 Integrals of form  $\int R(x, \sqrt{ax^2 + bx + c})dx$ ,  $R$  - rational function
  - Transforming to the forms  $\sqrt{x^2 + 1}$ ,  $\sqrt{-x^2 + 1}$ ,  $\sqrt{x^2 - 1}$
  - Table of Euler and trig substitutions
  - The case  $\sqrt{x^2 + 1}$
  - The case  $\sqrt{-x^2 + 1}$
  - The case  $\sqrt{x^2 - 1}$
- 2 Rationalizing Substitutions

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:



# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
  - Use trigonometric substitution or Euler substitution to transform to trigonometric or rational function integral (no radicals).

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

*Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?*

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
  - Use trigonometric substitution or Euler substitution to transform to trigonometric or rational function integral (no radicals).
  - Solve as previously studied.

# Integrals of form $\int R(x, \sqrt{ax^2 + bx + c})dx$ , $R$ - rational function

Let  $R(x, y)$  be an arbitrary rational expression in two variables (quotient of polynomials in two variables).

## Question

Can we integrate  $\int R(x, \sqrt{ax^2 + bx + c}) dx$ ?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
  - Use linear substitution to transform to one of three integrals:  $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
  - Use trigonometric substitution or Euler substitution to transform to trigonometric or rational function integral (no radicals).
  - Solve as previously studied.
- We motivate why we need **such integrals** by examples such as computing the area of an ellipse.

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.



# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

$$\sqrt{a^2 - x^2} =$$

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- **Instead, substitute  $x = a \sin \theta$ .**

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} =$$

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2(1 - \sin^2 \theta)} =$$

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2(1 - \sin^2 \theta)} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|.$$

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2(1 - \sin^2 \theta)} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|.$$

- With  $u = a^2 - x^2$ , the new variable is a function of the old one.

# Trigonometric Substitution

- To find the area of a circle or ellipse, one needs to compute  $\int \sqrt{a^2 - x^2} dx$ .
- For  $\int x\sqrt{a^2 - x^2} dx$ , the substitution  $u = a^2 - x^2$  would work.
- For  $\int \sqrt{a^2 - x^2} dx$ , we need a more elaborate substitution.
- Instead, substitute  $x = a \sin \theta$ .

$$\sqrt{a^2 - x^2} = \sqrt{a^2 - a^2 \sin^2 \theta} = \sqrt{a^2(1 - \sin^2 \theta)} = \sqrt{a^2 \cos^2 \theta} = a|\cos \theta|.$$

- With  $u = a^2 - x^2$ , the new variable is a function of the old one.
- With  $x = a \sin \theta$ , the old variable is a function of the new one.



# Linear substitutions to simplify radicals $\sqrt{ay^2 + by + c}$

- Using linear substitutions, radicals of form  $\sqrt{ay^2 + by + c}$ ,  $a \neq 0$ ,  $b^2 - 4ac \neq 0$  can be transformed to (multiple of):
  - $\sqrt{x^2 + 1}$
  - $\sqrt{-x^2 + 1}$
  - $\sqrt{x^2 - 1}$ .
- We already studied how to do that using completing the square when dealing with rational functions.

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\sqrt{x^2 + x + 1} =$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + ? - ? + 1}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\sqrt{x^2 + x + 1} = \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}\sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\ &= \sqrt{\left(x + ?\right)^2 + ?}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}\sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\ &= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\
 &= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} \\
 &= \sqrt{\frac{3}{4} \left(\frac{4}{3} \left(x + \frac{1}{2}\right)^2 + 1\right)}
 \end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\
 &= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} \\
 &= \sqrt{\frac{3}{4} \left(\frac{4}{3} \left(x + \frac{1}{2}\right)^2 + 1\right)} \\
 &= \frac{\sqrt{3}}{2} \sqrt{\left(\frac{2}{\sqrt{3}} \left(x + \frac{1}{2}\right)\right)^2 + 1}
 \end{aligned}$$



Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\
 &= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} \\
 &= \sqrt{\frac{3}{4} \left(\frac{4}{3} \left(x + \frac{1}{2}\right)^2 + 1\right)} \\
 &= \frac{\sqrt{3}}{2} \sqrt{\left(\frac{2}{\sqrt{3}} \left(x + \frac{1}{2}\right)\right)^2 + 1}
 \end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear substitution to transform  $\sqrt{x^2 + x + 1}$  to multiple of  $\sqrt{u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{x^2 + x + 1} &= \sqrt{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} \\
 &= \sqrt{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} \\
 &= \sqrt{\frac{3}{4} \left(\frac{4}{3} \left(x + \frac{1}{2}\right)^2 + 1\right)} \\
 &= \frac{\sqrt{3}}{2} \sqrt{\left(\frac{2}{\sqrt{3}} \left(x + \frac{1}{2}\right)\right)^2 + 1} \\
 &= \frac{\sqrt{3}}{2} \sqrt{u^2 + 1},
 \end{aligned}$$

where  $u = \frac{2}{\sqrt{3}} \left(x + \frac{1}{2}\right) = \frac{2\sqrt{3}}{3}x + \frac{\sqrt{3}}{3}$ .

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\sqrt{-2x^2 + x + 1} =$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\sqrt{-2x^2 + x + 1} = \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \quad - \quad - \frac{1}{2}\right)}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + ? - ? - \frac{1}{2}\right)}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)}\end{aligned}$$



Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)} \\
 &= \sqrt{\frac{9}{8}\left(-\frac{16}{9}\left(x - \frac{1}{4}\right)^2 + 1\right)}
 \end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}\sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\ &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)} \\ &= \sqrt{\frac{9}{8}\left(-\frac{16}{9}\left(x - \frac{1}{4}\right)^2 + 1\right)}\end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)} \\
 &= \sqrt{\frac{9}{8}\left(-\frac{16}{9}\left(x - \frac{1}{4}\right)^2 + 1\right)} \\
 &= \frac{3}{\sqrt{8}}\sqrt{-\left(\frac{4}{3}\left(x - \frac{1}{4}\right)\right)^2 + 1}
 \end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)} \\
 &= \sqrt{\frac{9}{8}\left(-\frac{16}{9}\left(x - \frac{1}{4}\right)^2 + 1\right)} \\
 &= \frac{3}{\sqrt{8}}\sqrt{-\left(\frac{4}{3}\left(x - \frac{1}{4}\right)\right)^2 + 1}
 \end{aligned}$$

Recall: linear substitution is subst. of the form  $u = px + q$ .

## Example

Use linear subst. to transform  $\sqrt{-2x^2 + x + 1}$  to multiple of  $\sqrt{-u^2 + 1}$ .

$$\begin{aligned}
 \sqrt{-2x^2 + x + 1} &= \sqrt{-2\left(x^2 - \frac{1}{2}x - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(x^2 - 2\frac{1}{4}x + \frac{1}{16} - \frac{1}{16} - \frac{1}{2}\right)} \\
 &= \sqrt{-2\left(\left(x - \frac{1}{4}\right)^2 - \frac{9}{16}\right)} \\
 &= \sqrt{\frac{9}{8}\left(-\frac{16}{9}\left(x - \frac{1}{4}\right)^2 + 1\right)} \\
 &= \frac{3}{\sqrt{8}}\sqrt{-\left(\frac{4}{3}\left(x - \frac{1}{4}\right)\right)^2 + 1} \\
 &= \frac{3}{\sqrt{8}}\sqrt{-u^2 + 1},
 \end{aligned}$$

where  $u = \frac{4}{3}\left(x - \frac{1}{4}\right) = \frac{4}{3}x - \frac{1}{3}$ .

- Let  $R$  be a rational function in two variables.

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .



- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .



- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .
- The Euler substitutions directly transform the integral to a rational function integral.

- Let  $R$  be a rational function in two variables.
- So far, with linear transformations we converted all integrals of the form  $\int R(x, \sqrt{ax^2 + bx + c})dx$  to one of the three forms:  
 $\int R(x, \sqrt{x^2 + 1})dx$ ,  $\int R(x, \sqrt{-x^2 + 1})dx$ ,  $\int R(x, \sqrt{x^2 - 1})dx$ .
- Each of the above integrals can be transformed to a rational trigonometric integral using 3 pairs of substitutions:  
 $x = \tan \theta$ ,  $x = \cot \theta$ ;  $x = \sin \theta$ ,  $x = \cos \theta$ ;  $x = \csc \theta$ ,  $x = \sec \theta$ .
- We studied that trigonometric integrals are converted to rational function integrals via  $\theta = 2 \arctan t$ .
- The resulting 3 pairs of substitutions are called Euler substitutions:  
 $x = \tan(2 \arctan t)$ ,  $x = \cot(2 \arctan t)$ ;  $x = \sin(2 \arctan t)$ ,  
 $x = \cos(2 \arctan t)$ ;  $x = \csc(2 \arctan t)$ ,  $x = \sec(2 \arctan t)$ .
- The Euler substitutions directly transform the integral to a rational function integral.
- We will demonstrate that the Euler substitutions are **rational**.

# Trigonometric substitution and Euler substitution

Expression	Substitution	Variable range	Relevant identity
$\sqrt{x^2 + 1}$	$x = \tan \theta$	$\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$	$1 + \tan^2 \theta = \sec^2 \theta$
	$x = \cot \theta$	$\theta \in (0, \pi)$	$1 + \cot^2 \theta = \csc^2 \theta$
$\sqrt{-x^2 + 1}$	$x = \sin \theta$	$\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$1 - \sin^2 \theta = \cos^2 \theta$
	$x = \cos \theta$	$\theta \in (0, \pi)$	$1 - \cos^2 \theta = \sin^2 \theta$
$\sqrt{x^2 - 1}$	$x = \csc \theta$	$\theta \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$	$\csc^2 \theta - 1 = \cot^2 \theta$
	$x = \sec \theta$	$\theta \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$	$\sec^2 \theta - 1 = \tan^2 \theta$

Euler substitution by applying in addition  $\theta = 2 \arctan t$

$\sqrt{x^2 + 1}$	$x = \frac{2t}{1-t^2}$	$-1 < t < 1$	(?)
	$x = \frac{1}{2} \left( \frac{1}{t} - t \right)$	$0 < t$	(?)
$\sqrt{-x^2 + 1}$	$x = \frac{2t}{1+t^2}$	$-1 \leq t \leq 1$	(?)
	$x = \frac{1-t^2}{1+t^2}$	$0 < t$	(?)
$\sqrt{x^2 - 1}$	$x = \frac{1}{2} \left( \frac{1}{t} + t \right)$	$t \in (-\infty, -1) \cup [0, 1)$	(?)
	$x = \frac{1+t^2}{1-t^2}$	$t \in (-\infty, -1) \cup [0, 1)$	(?)

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} =$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \sqrt{\cot^2 \theta + 1}$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}\sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\ &= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1}\end{aligned}$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}\sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\ &= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1} \\ &= \sqrt{\frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta}}\end{aligned}$$



# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}\sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\&= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1} \\&= \sqrt{\frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta}} \\&= \sqrt{\frac{1}{\sin^2 \theta}} = \frac{1}{\sqrt{\sin^2 \theta}}\end{aligned}$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}
 \sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\
 &= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1} \\
 &= \sqrt{\frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta}} \\
 &= \sqrt{\frac{1}{\sin^2 \theta}} = \frac{1}{\sqrt{\sin^2 \theta}} \\
 &= \frac{1}{\sin \theta}
 \end{aligned}$$

when  $\theta \in (0, \pi)$  we have  
 $\sin \theta \geq 0$  and so  
 $\sqrt{\sin^2 \theta} = \sin \theta$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}
 \sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\
 &= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1} \\
 &= \sqrt{\frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta}} \\
 &= \sqrt{\frac{1}{\sin^2 \theta}} = \frac{1}{\sqrt{\sin^2 \theta}} \\
 &= \frac{1}{\sin \theta} = \csc \theta .
 \end{aligned}$$

when  $\theta \in (0, \pi)$  we have  
 $\sin \theta \geq 0$  and so  
 $\sqrt{\sin^2 \theta} = \sin \theta$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\begin{aligned}
 \sqrt{x^2 + 1} &= \sqrt{\cot^2 \theta + 1} \\
 &= \sqrt{\frac{\cos^2 \theta}{\sin^2 \theta} + 1} \\
 &= \sqrt{\frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta}} \\
 &= \sqrt{\frac{1}{\sin^2 \theta}} = \frac{1}{\sqrt{\sin^2 \theta}} \\
 &= \frac{1}{\sin \theta} = \csc \theta .
 \end{aligned}$$

when  $\theta \in (0, \pi)$  we have  
 $\sin \theta \geq 0$  and so  
 $\sqrt{\sin^2 \theta} = \sin \theta$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ .

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ . To summarize:

## Definition

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$  is given by:

$$\begin{aligned} x &= \cot \theta \\ \sqrt{x^2 + 1} &= \frac{1}{\sin \theta} = \csc \theta \\ dx &= \quad \quad \quad ?d\theta \\ \theta &= \operatorname{arccot} x . \end{aligned}$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ . To summarize:

## Definition

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$  is given by:

$$\begin{aligned} x &= \cot \theta \\ \sqrt{x^2 + 1} &= \frac{1}{\sin \theta} = \csc \theta \\ dx &= \quad \quad \quad ?d\theta \\ \theta &= \operatorname{arccot} x . \end{aligned}$$



# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ . To summarize:

## Definition

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$  is given by:

$$\begin{aligned} x &= \cot \theta \\ \sqrt{x^2 + 1} &= \frac{1}{\sin \theta} = \csc \theta \\ dx &= \quad \quad \quad ?d\theta \\ \theta &= \operatorname{arccot} x . \end{aligned}$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ . To summarize:

## Definition

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$  is given by:

$$\begin{aligned} x &= \cot \theta \\ \sqrt{x^2 + 1} &= \frac{1}{\sin \theta} = \csc \theta \end{aligned}$$

$$dx = \quad \quad \quad ?d\theta$$

$$\theta = \operatorname{arccot} x .$$

# Trigonometric substitution $x = \cot \theta$ for $\sqrt{x^2 + 1}$

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$ :

$$\sqrt{x^2 + 1} = \frac{1}{\sin \theta} = \csc \theta .$$

The differential  $dx$  can be expressed via  $d\theta$  from  $x = \cot \theta$ . To summarize:

## Definition

The trigonometric substitution  $x = \cot \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 + 1}$  is given by:

$$\begin{aligned} x &= \cot \theta \\ \sqrt{x^2 + 1} &= \frac{1}{\sin \theta} = \csc \theta \\ dx &= -\frac{d\theta}{\sin^2 \theta} = -\csc^2 \theta \, d\theta \\ \theta &= \operatorname{arccot} x . \end{aligned}$$

## Example

$$\int \frac{1}{x^2 \sqrt{x^2 + 9}} dx$$

## Example

$$\int \frac{1}{x^2 \sqrt{x^2 + 9}} dx = \int \frac{1}{x^2 \cdot 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx$$

## Example

$$\int \frac{1}{x^2 \sqrt{x^2 + 9}} dx = \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx$$

Set

$$\frac{x}{3} = \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\int \frac{1}{x^2 \sqrt{x^2 + 9}} dx = \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta)
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$



## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(\textcolor{red}{3} \cot \theta)^2 \textcolor{red}{3} \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{\textcolor{red}{27} \cot^2 \theta \sqrt{\textcolor{red}{?}}} \left( \textcolor{red}{?} \right) d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{?}} \left( ? \right) d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} \left( ? \right) d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} \left( ? \right) d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow$$

$$\csc \theta > 0$$

## Example

$$\int \frac{1}{x^2 \sqrt{x^2 + 9}} dx = \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx$$

$$= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta)$$

$$= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta$$

$$= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta$$

$$= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$



## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(?)
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta)
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2}
 \end{aligned}$$

Set

$\frac{x}{3} = \cot \theta$   
 $x = 3 \cot \theta$   
 $\theta \in (0, \pi)$   
 $\theta \in (0, \pi) \Rightarrow \csc \theta > 0$

Set  $u = \cos \theta$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = ? + C
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

$$\text{Set } u = \cos \theta$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

$$\text{Set } u = \cos \theta$$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C
 \end{aligned}$$

Set

$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

Set  $u = \cos \theta$

## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C
 \end{aligned}$$

Set


$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

Set  $u = \cos \theta$



## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C
 \end{aligned}$$

Set


$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

Set  $u = \cos \theta$





## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C
 \end{aligned}$$

Set


$$\frac{x}{3} = \cot \theta$$

$$x = 3 \cot \theta$$

$$\theta \in (0, \pi)$$

$$\theta \in (0, \pi) \Rightarrow \csc \theta > 0$$

Set  $u = \cos \theta$



## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C
 \end{aligned}$$


Set

$$\begin{aligned}
 \frac{x}{3} &= \cot \theta \\
 x &= 3 \cot \theta
 \end{aligned}$$

$$\theta \in (0, \pi)$$

$$\begin{aligned}
 \theta &\in (0, \pi) \Rightarrow \\
 \csc \theta &> 0
 \end{aligned}$$

Set  $u = \cos \theta$



## Example

$$\begin{aligned}
 \int \frac{1}{x^2 \sqrt{x^2 + 9}} dx &= \int \frac{1}{x^2 3 \sqrt{\left(\frac{x}{3}\right)^2 + 1}} dx \\
 &= \int \frac{1}{(3 \cot \theta)^2 3 \sqrt{\cot^2 \theta + 1}} d(3 \cot \theta) \\
 &= \int \frac{1}{27 \cot^2 \theta \sqrt{\csc^2 \theta}} (-3 \csc^2 \theta) d\theta \\
 &= \frac{1}{9} \int \frac{-\csc^2 \theta}{\cot^2 \theta \csc \theta} d\theta \\
 &= \frac{1}{9} \int \frac{-\sin \theta}{\cos^2 \theta} d\theta = \frac{1}{9} \int \frac{1}{\cos^2 \theta} d(\cos \theta) \\
 &= \frac{1}{9} \int \frac{du}{u^2} = -\frac{1}{9u} + C = -\frac{\sec \theta}{9} + C \\
 &= -\frac{\sqrt{x^2 + 9}}{9x} + C
 \end{aligned}$$


Set

$$\begin{aligned}
 \frac{x}{3} &= \cot \theta \\
 x &= 3 \cot \theta
 \end{aligned}$$

$$\theta \in (0, \pi)$$

$$\begin{aligned}
 \theta \in (0, \pi) &\Rightarrow \\
 \csc \theta &> 0
 \end{aligned}$$

Set  $u = \cos \theta$



# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t$ , transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t$ , transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above?

## Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? **We get the Euler substitution:**

$$x =$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}x &= \cot \theta \\&= \cot (2 \arctan t)\end{aligned}$$



# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$= \cot(2 \arctan t) \quad | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$= \cot(2 \arctan t) \quad \left| \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \right.$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) & | \text{Recall: } \cot(2z) &= \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)}
 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) && | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)} \\
 &= \frac{1 - t^2}{2t}
 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) && | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)} \\
 &= \frac{1 - t^2}{2t} \\
 &= \frac{1}{2} \left( \frac{1}{t} - t \right) .
 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) && | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)} \\
 &= \frac{1 - t^2}{2t} \\
 &= \frac{1}{2} \left( \frac{1}{t} - t \right) .
 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) && | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)} \\
 &= \frac{1 - t^2}{2t} \\
 &= \frac{1}{2} \left( \frac{1}{t} - t \right) .
 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cot \theta$$

$$\begin{aligned}
 &= \cot(2 \arctan t) && | \text{Recall: } \cot(2z) = \frac{\cos(2z)}{\sin(2z)} = \frac{1 - \tan^2 z}{2 \tan z} \\
 &= \frac{1 - \tan^2(\arctan t)}{2 \tan(\arctan t)} \\
 &= \frac{1 - t^2}{2t} \\
 &= \frac{1}{2} \left( \frac{1}{t} - t \right) .
 \end{aligned}$$



# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} =$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} - t \right)^2 + 4} \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} - t \right)^2 + 4} \quad \mid \quad \left( \frac{1}{t} - t \right)^2 + 4 = \left( \frac{1}{t} + t \right)^2 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} - t \right)^2 + 4} \quad \Bigg| \quad \left( \frac{1}{t} - t \right)^2 + 4 = \left( \frac{1}{t} + t \right)^2 \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} + t \right)^2} \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} + t \right)^2} \quad \left| \begin{array}{l} \sqrt{\left( \frac{1}{t} + t \right)^2} = \frac{1}{t} + t \\ \text{because } t > 0 \end{array} \right. \end{aligned}$$



# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} + t \right)^2} \quad \left| \begin{array}{l} \sqrt{\left( \frac{1}{t} + t \right)^2} = \frac{1}{t} + t \\ \text{because } t > 0 \end{array} \right. \\ &= \frac{1}{2} \left( \frac{1}{t} + t \right) . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\begin{aligned} \sqrt{x^2 + 1} &= \sqrt{\frac{1}{4} \left( \frac{1}{t} - t \right)^2 + 1} \\ &= \frac{1}{2} \sqrt{\left( \frac{1}{t} + t \right)^2} \quad \left| \begin{array}{l} \sqrt{\left( \frac{1}{t} + t \right)^2} = \frac{1}{t} + t \\ \text{because } t > 0 \end{array} \right. \\ &= \frac{1}{2} \left( \frac{1}{t} + t \right) . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$dx =$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$dx = d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right)$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$dx = d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) =$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$dx = d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) . \end{aligned}$$



# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) = \sqrt{x^2 + 1} - x . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) = \sqrt{x^2 + 1} - x . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1}{2} \left( \frac{1}{t} - t \right) .$$

We can furthermore compute

$$\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right) .$$

Finally compute

$$\begin{aligned} dx &= d \left( \frac{1}{2} \left( \frac{1}{t} - t \right) \right) = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \frac{1}{2} \left( \frac{1}{t} + t \right) - \frac{1}{2} \left( \frac{1}{t} - t \right) = \sqrt{x^2 + 1} - x . \end{aligned}$$

# Euler subst. for $\sqrt{x^2 + 1}$ corresponding to $x = \cot \theta$

- $x = \cot \theta$  transforms  $dx, x, \sqrt{x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

## Definition

The Euler substitution for  $\sqrt{x^2 + 1}$  corresponding to  $x = \cot \theta$  is given by:

$$\begin{aligned} x &= \frac{1}{2} \left( \frac{1}{t} - t \right), & t > 0 \\ \sqrt{x^2 + 1} &= \frac{1}{2} \left( \frac{1}{t} + t \right) \\ dx &= -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ t &= \sqrt{x^2 + 1} - x \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\int \sqrt{x^2 + 1} \, dx =$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\int \sqrt{x^2 + 1} \, dx = - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$$



Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\int \sqrt{x^2 + 1} \, dx = - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\ &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\ &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \end{aligned}$$



Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\ &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} (t^{-1} - t) \frac{1}{2} (t^{-1} + t) \right) - \frac{1}{2} \ln t + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned} \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\ &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\ &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\ &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C
 \end{aligned}$$



Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \frac{1}{\left( \sqrt{x^2 + 1} - x \right)} + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \frac{\sqrt{x^2 + 1} + x}{\left( \sqrt{x^2 + 1} - x \right) \left( \sqrt{x^2 + 1} + x \right)} + C
 \end{aligned}$$

Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \frac{\sqrt{x^2 + 1} + x}{\left( \sqrt{x^2 + 1} - x \right) \left( \sqrt{x^2 + 1} + x \right)} + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \left( \sqrt{x^2 + 1} + x \right) + C
 \end{aligned}$$


Euler substitution:  $x = \frac{1}{2} \left( \frac{1}{t} - t \right)$ ,  $\sqrt{x^2 + 1} = \frac{1}{2} \left( \frac{1}{t} + t \right)$ ,  
 $t = \sqrt{x^2 + 1} - x$ ,  $dx = -\frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt$ . Recall  $t > 0$ .

## Example

$$\begin{aligned}
 \int \sqrt{x^2 + 1} \, dx &= - \int \frac{1}{2} \left( \frac{1}{t} + t \right) \frac{1}{2} \left( \frac{1}{t^2} + 1 \right) dt \\
 &= -\frac{1}{4} \int \left( \frac{1}{t^3} + 2\frac{1}{t} + t \right) dt \\
 &= -\frac{1}{4} \left( -\frac{t^{-2}}{2} + 2 \ln |t| + \frac{t^2}{2} \right) + C \\
 &= \frac{1}{2} \left( \frac{1}{2} \left( t^{-1} - t \right) \frac{1}{2} \left( t^{-1} + t \right) \right) - \frac{1}{2} \ln t + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} - \frac{1}{2} \ln \left( \sqrt{x^2 + 1} - x \right) + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \frac{\sqrt{x^2 + 1} + x}{\left( \sqrt{x^2 + 1} - x \right) \left( \sqrt{x^2 + 1} + x \right)} + C \\
 &= \frac{1}{2} x \sqrt{x^2 + 1} + \frac{1}{2} \ln \left( \sqrt{x^2 + 1} + x \right) + C
 \end{aligned}$$


## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




why do we call  
 $y = \sqrt{x^2 + 1}$  hyperbola?



## Example


Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola?

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2\end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1\end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$(y - x)(y + x) = 1$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{1}{2} (y - x) (y + x) = \frac{1}{2}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) = \frac{1}{2}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:


$$\begin{aligned}
 \sqrt{x^2 + 1} &= y \\
 x^2 + 1 &= y^2 \\
 y^2 - x^2 &= 1 \\
 \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\
 uv &= \frac{1}{2}
 \end{aligned}$$

where 
$$\begin{cases}
 u = \frac{\sqrt{2}}{2}(y - x) \\
 v = \frac{\sqrt{2}}{2}(y + x)
 \end{cases}$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}.$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}
 \sqrt{x^2 + 1} &= y \\
 x^2 + 1 &= y^2 \\
 y^2 - x^2 &= 1 \\
 \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\
 uv &= \frac{1}{2} \\
 v &= \frac{1}{2u},
 \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned}
 \sqrt{x^2 + 1} &= y \\
 x^2 + 1 &= y^2 \\
 y^2 - x^2 &= 1 \\
 \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\
 uv &= \frac{1}{2} \\
 v &= \frac{1}{2u},
 \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line  
 $u = 0$  equals


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2}$$

We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:


$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



distance b-n  $(x, y)$  and line

$u = 0$  equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} = \pm \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



**Signed** distance b-n  $(x, y)$  and line  $u = 0$  equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) = u. \end{aligned}$$


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ .


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that **signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .**


We studied  $v = \frac{1}{u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



Signed distance b-n  $(x, y)$  and line  $u = 0$  equals  $u$ . Similarly compute that signed distance b-n  $(x, y)$  and the line  $v = 0$  equals  $v$ .  
 $\Rightarrow y^2 - x^2 = 1$  is the **hyperbola**  
 $v = \frac{1/2}{u}$  in the  $(u, v)$ -plane.

We studied  $v = \frac{1}{2u}$  is called a hyperbola: why do we call  $y = \sqrt{x^2 + 1}$  hyperbola? Compute:


$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where  $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$ . Consider an arbitrary point  $(x, y)$ .



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned} & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\ &= 2 \left[ x\sqrt{x^2 + 1} \right. \\ & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) - \left( -2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} - \ln \left( \sqrt{(2\sqrt{2})^2 + 1} - 2\sqrt{2} \right) \right) \right)
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} \right. \\
 & \quad \left. + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right)
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2})
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



- Recall: integral can be solved via  $x = \tan \theta$ .


The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$



## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- Geometric interpretation of  $\theta$ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- **Geometric interpretation of  $\theta$ ?**

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2 \ln \left( 3 + 2\sqrt{2} \right) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .




- Recall: integral can be solved via  $x = \tan \theta$ .
- Geometric interpretation of  $\theta$ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find the area locked b-n the hyperbolas  $y = \pm\sqrt{x^2 + 1}$  and  $x = \pm 2\sqrt{2}$ .



- Recall: integral can be solved via  $x = \tan \theta$ .
- Geometric interpretation of  $\theta$ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[ x\sqrt{x^2 + 1} + \ln \left( \sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left( 2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left( \sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u =$
- $du =$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u =$
- $du =$



## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du =$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du =$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx =$$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx = \frac{1}{2} \int \frac{du}{\sqrt{u}} =$$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx = \frac{1}{2} \int \frac{du}{\sqrt{u}} =$$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx = \frac{1}{2} \int \frac{du}{\sqrt{u}} =$$

## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx = \frac{1}{2} \int \frac{du}{\sqrt{u}} = \sqrt{u} + C =$$



## Example

Find  $\int \frac{x}{\sqrt{x^2+4}} dx$ .

- We could use the trig substitution  $x = 2 \tan \theta$ .
- But there is an easier way:
- $u = x^2 + 4$ .
- $du = 2x dx$ .

$$\int \frac{x}{\sqrt{x^2+4}} dx = \frac{1}{2} \int \frac{du}{\sqrt{u}} = \sqrt{u} + C = \sqrt{x^2+4} + C$$

# Trigonometric substitution $x = \cos \theta$ for $\sqrt{-x^2 + 1}$

The trigonometric substitution  $x = \cos \theta$ ,  $\theta \in [0, \pi]$  for  $\sqrt{-x^2 + 1}$ :

$$\sqrt{-x^2 + 1} =$$

# Trigonometric substitution $x = \cos \theta$ for $\sqrt{-x^2 + 1}$

The trigonometric substitution  $x = \cos \theta$ ,  $\theta \in [0, \pi]$  for  $\sqrt{-x^2 + 1}$ :

$$\sqrt{-x^2 + 1} = \sqrt{1 - \cos^2 \theta}$$

# Trigonometric substitution $x = \cos \theta$ for $\sqrt{-x^2 + 1}$

The trigonometric substitution  $x = \cos \theta$ ,  $\theta \in [0, \pi]$  for  $\sqrt{-x^2 + 1}$ :

$$\begin{aligned}\sqrt{-x^2 + 1} &= \sqrt{1 - \cos^2 \theta} \\ &= \sqrt{\sin^2 \theta} \\ &= \sin \theta \quad .\end{aligned}$$

# Trigonometric substitution $x = \cos \theta$ for $\sqrt{-x^2 + 1}$

The trigonometric substitution  $x = \cos \theta$ ,  $\theta \in [0, \pi]$  for  $\sqrt{-x^2 + 1}$ :

$$\begin{aligned}
 \sqrt{-x^2 + 1} &= \sqrt{1 - \cos^2 \theta} \\
 &= \sqrt{\sin^2 \theta} \\
 &= \sin \theta \quad .
 \end{aligned}
 \quad \left| \begin{array}{l} \text{when } \theta \in [0, \pi] \text{ we have} \\ \sin \theta \geq 0 \text{ and so } \sqrt{\sin^2 \theta} = \sin \theta \end{array} \right.$$

To summarize:


## Definition

The trigonometric substitution  $x = \cos \theta$ ,  $\theta \in [0, \pi]$  for  $\sqrt{-x^2 + 1}$  is given by:

$$\begin{aligned}
 x &= \cos \theta \\
 \sqrt{-x^2 + 1} &= \sin \theta \\
 dx &= -\sin \theta d\theta \\
 \theta &= \arccos x \quad .
 \end{aligned}$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .




## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x =$

- Then  $dx =$

$$\sqrt{9-x^2} =$$




## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x =$

- Then  $dx =$

$$\sqrt{9-x^2} =$$






## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx =$   
 $\sqrt{9-x^2} =$




## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .

- Then  $dx =$

$$\sqrt{9-x^2} =$$




## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .

$$\sqrt{9-x^2} =$$




## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .


$$\sqrt{9 - x^2} = \sqrt{9 - 9 \sin^2 \theta} =$$



## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} =$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| =$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .



$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$


$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta$$



## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\int \frac{\sqrt{9-x^2}}{x^2} dx = \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta \end{aligned}$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta = -\cot\theta - \theta + C \end{aligned}$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta = -\cot\theta - \theta + C \end{aligned}$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .



$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$


$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta = -\cot\theta - \theta + C \end{aligned}$$



## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta = -\cot\theta - \theta + C \end{aligned}$$

## Example

Evaluate  $\int \frac{\sqrt{9-x^2}}{x^2} dx$ .

- Let  $x = 3 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ .
- Then  $dx = 3 \cos \theta d\theta$ .




$$\sqrt{9-x^2} = \sqrt{9-9\sin^2\theta} = \sqrt{9\cos^2\theta} = 3|\cos\theta| = 3\cos\theta$$

$$\begin{aligned} \int \frac{\sqrt{9-x^2}}{x^2} dx &= \int \frac{3\cos\theta}{9\sin^2\theta} 3\cos\theta d\theta = \int \cot^2\theta d\theta \\ &= \int (\csc^2\theta - 1) d\theta = -\cot\theta - \theta + C \\ &= -\frac{\sqrt{9-x^2}}{x} - \arcsin\left(\frac{x}{3}\right) + C \end{aligned}$$


## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .




Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .




Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$
$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Express  $y$  via  $x$ :


$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$


$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b \sqrt{1 - \frac{x^2}{a^2}}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$


$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b \sqrt{1 - \frac{x^2}{a^2}}$$



## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .




Express  $y$  via  $x$ :

$$\begin{aligned}\frac{x^2}{a^2} + \frac{y^2}{b^2} &= 1 \\ \frac{y^2}{b^2} &= 1 - \frac{x^2}{a^2} \\ y^2 &= b^2 \left( 1 - \frac{x^2}{a^2} \right) \\ y &= \pm b \sqrt{1 - \frac{x^2}{a^2}}\end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



The area in question is

Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$


$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b \sqrt{1 - \frac{x^2}{a^2}}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



The area in question is

$$\int_{?}^{?} 2b\sqrt{1 - \frac{x^2}{a^2}} dx$$

Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$


$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b\sqrt{1 - \frac{x^2}{a^2}}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



The area in question is

$$\int_{-a}^a 2b\sqrt{1 - \frac{x^2}{a^2}} dx$$

Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$


$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b\sqrt{1 - \frac{x^2}{a^2}}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



The area in question is


$$\int_{-a}^a 2b\sqrt{1 - \frac{x^2}{a^2}} dx$$

Express  $y$  via  $x$ :

$$\begin{aligned}\frac{x^2}{a^2} + \frac{y^2}{b^2} &= 1 \\ \frac{y^2}{b^2} &= 1 - \frac{x^2}{a^2} \\ y^2 &= b^2 \left(1 - \frac{x^2}{a^2}\right) \\ y &= \pm b\sqrt{1 - \frac{x^2}{a^2}}\end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



The area in question is

$$\int_{-a}^a 2b\sqrt{1 - \frac{x^2}{a^2}} dx$$

$$= 4 \int_0^a b\sqrt{1 - \frac{x^2}{a^2}} dx.$$

Express  $y$  via  $x$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$


$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \left( 1 - \frac{x^2}{a^2} \right)$$

$$y = \pm b\sqrt{1 - \frac{x^2}{a^2}}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .




The area in question is

$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx$$

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx$$


The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$



## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}}$


$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx$$

The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta}$ .


$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx$$

The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ .


$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx$$

The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ .


$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx = \int \cos \theta d(a \sin \theta)$$

The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx = \int_0^{\frac{\pi}{2}} \cos \theta d(a \sin \theta)$$

The area in question is

$$\begin{aligned} & \int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta d(\textcolor{red}{a} \sin \theta) \\ &= \textcolor{red}{a} \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .

$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \end{aligned}$$


The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$



## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\ &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\ &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\ &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \\ &= a \left( 0 + \frac{\pi}{4} - (0 + 0) \right) \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\ &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \\ &= a \left( 0 + \frac{\pi}{4} - (0 + 0) \right) \\ &= \frac{a\pi}{4} \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .


$$\begin{aligned}
 \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta \, d(a \sin \theta) \\
 &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\
 &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\
 &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \\
 &= a \left( 0 + \frac{\pi}{4} - (0 + 0) \right) \\
 &= \frac{a\pi}{4}
 \end{aligned}$$

The area in question is

$$\begin{aligned}
 &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\
 &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \\
 &= 4b \frac{a\pi}{4}
 \end{aligned}$$

## Example

Find the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a, b > 0$ .



Trig subst.: set  $x = a \sin \theta$ ,  $\theta \in (0, \frac{\pi}{2})$ .

Compute:  $\sqrt{1 - \frac{x^2}{a^2}} = \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} = \sqrt{1 - \sin^2 \theta} = \cos \theta$ . When  $x = 0$ ,  $\theta = 0$  and when  $x = a$ ,  $\theta = \frac{\pi}{2}$ .

$$\begin{aligned} \int_0^a \sqrt{1 - \frac{x^2}{a^2}} dx &= \int_0^{\frac{\pi}{2}} \cos \theta d(a \sin \theta) \\ &= a \int_0^{\frac{\pi}{2}} \cos^2 \theta d\theta \\ &= a \int_0^{\frac{\pi}{2}} \frac{\cos(2\theta) + 1}{2} d\theta \\ &= a \left[ \frac{\sin(2\theta)}{4} + \frac{\theta}{2} \right]_{\theta=0}^{\theta=\frac{\pi}{2}} \\ &= a \left( 0 + \frac{\pi}{4} - (0 + 0) \right) \\ &= \frac{a\pi}{4} \end{aligned}$$

The area in question is

$$\begin{aligned} &\int_{-a}^a 2b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4 \int_0^a b \sqrt{1 - \frac{x^2}{a^2}} dx \\ &= 4b \frac{a\pi}{4} = \pi ab \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:



## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 - (x^2 + 2x \quad) =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u =$  . Then  $du =$  and  $x =$  .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u =$  . Then  $du =$  and  $x =$  .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du =$       and  $x =$       .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du =$       and  $x =$       .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$



## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x =$  .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x =$  .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u =$  Then  $du =$
- $\sqrt{4 - u^2} =$



## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u =$  Then  $du =$
- $\sqrt{4 - u^2} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du =$
- $\sqrt{4 - u^2} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du =$
- $\sqrt{4 - u^2} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| =$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$



## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$

$$\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .

- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$

$$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\ &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C \end{aligned}$$



## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$

$$\begin{aligned}
 \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\
 &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\
 &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C
 \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\ &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .

- $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$

$$\begin{aligned}
 \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\
 &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\
 &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C \\
 &= -\sqrt{3 - 2x - x^2} - \sin^{-1} \left( \frac{x+1}{2} \right) + C
 \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
  - $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
  - Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
  - $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
  - Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .
  - $\sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$
- $$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\ &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C \\ &= -\sqrt{3 - 2x - x^2} - \sin^{-1} \left( \frac{x+1}{2} \right) + C \end{aligned}$$

## Example

Evaluate  $\int \frac{x}{\sqrt{3-2x-x^2}} dx$ .

- Complete the square under the root sign:
- $3 - 2x - x^2 = 3 + 1 - (x^2 + 2x + 1) = 4 - (x + 1)^2$
- Substitute  $u = x + 1$ . Then  $du = dx$  and  $x = u - 1$ .
- $\int \frac{x}{\sqrt{3-2x-x^2}} dx = \int \frac{x}{\sqrt{4-(x+1)^2}} dx = \int \frac{u-1}{\sqrt{4-u^2}} du$
- Let  $u = 2 \sin \theta$ , where  $-\pi/2 \leq \theta \leq \pi/2$ . Then  $du = 2 \cos \theta d\theta$ .

$$\bullet \sqrt{4 - u^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 |\cos \theta| = 2 \cos \theta$$

$$\begin{aligned} \int \frac{x}{\sqrt{3-2x-x^2}} dx &= \int \frac{u-1}{\sqrt{4-u^2}} du = \int \frac{2 \sin \theta - 1}{2 \cos \theta} 2 \cos \theta d\theta \\ &= \int (2 \sin \theta - 1) d\theta = -2 \cos \theta - \theta + C \\ &= -\sqrt{4 - u^2} - \sin^{-1} \left( \frac{u}{2} \right) + C \\ &= -\sqrt{3 - 2x - x^2} - \sin^{-1} \left( \frac{x+1}{2} \right) + C \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t$ , transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t$ , transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above?

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? **We get the Euler substitution:**

$$x =$$



# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cos \theta$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}x &= \cos \theta \\&= \cos(2 \arctan t)\end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned} x &= \cos \theta \\ &= \cos(2 \arctan t) \end{aligned} \quad \left| \quad \cos(2z) = \right.$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \cos \theta$$

$$= \cos(2 \arctan t)$$

$$\left| \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \cos \theta \\
 &= \cos(2 \arctan t) \\
 &= \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z}
 \right.$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \cos \theta \\
 &= \cos(2 \arctan t) & \left| \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right. \\
 &= \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} \\
 &= \frac{1 - t^2}{1 + t^2}
 \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \cos \theta \\
 &= \cos(2 \arctan t) & \left| \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right. \\
 &= \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} \\
 &= \frac{1 - t^2}{1 + t^2}
 \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$



# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---

$$\sqrt{-x^2 + 1} =$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---

$$\sqrt{-x^2 + 1} = \sqrt{1 - \left( \frac{1 - t^2}{1 + t^2} \right)^2}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned}\sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}}\end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} \quad | \quad (1 + t^2)^2 - (1 - t^2)^2 = ? \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} \quad | \quad (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} \quad | \quad (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \\ &= \sqrt{\frac{4t^2}{(1 + t^2)^2}} \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} & | (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \\ &= \sqrt{\frac{4t^2}{(1 + t^2)^2}} & | \sqrt{4t^2} = 2t \text{ because } t > 0 \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} & | (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \\ &= \sqrt{\frac{4t^2}{(1 + t^2)^2}} & | \sqrt{4t^2} = 2t \text{ because } t > 0 \\ &= \frac{2t}{1 + t^2} \end{aligned}$$



# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---


$$\begin{aligned} \sqrt{-x^2 + 1} &= \sqrt{1 - \left(\frac{1 - t^2}{1 + t^2}\right)^2} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 + t^2)^2}} & | (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \\ &= \sqrt{\frac{4t^2}{(1 + t^2)^2}} & | \sqrt{4t^2} = 2t \text{ because } t > 0 \\ &= \frac{2t}{1 + t^2} \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

$$t^2(\textcolor{red}{x} + \textcolor{red}{1}) = 1 - x$$

$$t^2 = \frac{1 - x}{\textcolor{red}{1} + \textcolor{red}{x}}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

$$t^2 = \frac{1 - x}{1 + x}$$

$$t = \frac{\sqrt{1 - x}}{\sqrt{1 + x}}$$

we use  $t > 0$



# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

$$t^2 = \frac{1 - x}{1 + x}$$

$$t = \frac{\sqrt{1 - x}}{\sqrt{1 + x}} \frac{\sqrt{1 + x}}{\sqrt{1 + x}}$$

we use  $t > 0$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

$$t^2 = \frac{1 - x}{1 + x}$$

$$t = \frac{\sqrt{1 - x} \sqrt{1 + x}}{\sqrt{1 + x} \sqrt{1 + x}} = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

we use  $t > 0$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$(1 + t^2)x = 1 - t^2$$

$$t^2(x + 1) = 1 - x$$

$$t^2 = \frac{1 - x}{1 + x}$$

$$t = \frac{\sqrt{1 - x}}{\sqrt{1 + x}} = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

we use  $t > 0$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

---

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

---

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$


---

$dx$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

$$dx = d\left(\frac{1 - t^2}{1 + t^2}\right)$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$


---

$$dx = d\left(\frac{1 - t^2}{1 + t^2}\right) = d\left(\frac{2 - (1 + t^2)}{1 + t^2}\right)$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

$$\begin{aligned} dx &= d\left(\frac{1 - t^2}{1 + t^2}\right) = d\left(\frac{2 - (1 + t^2)}{1 + t^2}\right) \\ &= d\left(\frac{2}{1 + t^2} - 1\right) \end{aligned}$$



# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$


---

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$


---

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$


---

$$\begin{aligned} dx &= d\left(\frac{1 - t^2}{1 + t^2}\right) = d\left(\frac{2 - (1 + t^2)}{1 + t^2}\right) \\ &= d\left(\frac{2}{1 + t^2} - 1\right) = -\frac{4t}{(1 + t^2)^2} dt \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

$$x = \frac{1 - t^2}{1 + t^2}$$

$$\sqrt{-x^2 + 1} = \frac{2t}{1 + t^2}$$

$$t = \frac{\sqrt{-x^2 + 1}}{x + 1}$$

$$\begin{aligned} dx &= d\left(\frac{1 - t^2}{1 + t^2}\right) = d\left(\frac{2 - (1 + t^2)}{1 + t^2}\right) \\ &= d\left(\frac{2}{1 + t^2} - 1\right) = -\frac{4t}{(1 + t^2)^2} dt \end{aligned}$$

# Euler subst. for $\sqrt{-x^2 + 1}$ corresponding to $x = \cos \theta$

- $x = \cos \theta$  transforms  $dx, x, \sqrt{-x^2 + 1}$  to trig form.
- $\theta = 2 \arctan t, t > 0$  transforms  $d\theta, \cos \theta, \sin \theta$  to rational form.

What if we compose the above? We get the Euler substitution:

## Definition

The Euler substitution for  $\sqrt{-x^2 + 1}$  corresponding to  $x = \cos \theta$  is given by:

$$\begin{aligned} x &= \frac{1 - t^2}{1 + t^2}, & t > 0 \\ \sqrt{-x^2 + 1} &= \frac{2t}{1 + t^2} \\ dx &= -\frac{4t}{(t^2 + 1)^2} dt \\ t &= \frac{\sqrt{-x^2 + 1}}{x + 1}. \end{aligned}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} =$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \sqrt{\sec^2 \theta - 1}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}\sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\ &= \sqrt{\frac{1}{\cos^2 \theta} - 1}\end{aligned}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}\sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\ &= \sqrt{\frac{1}{\cos^2 \theta} - 1} \\ &= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}}\end{aligned}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}\sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\ &= \sqrt{\frac{1}{\cos^2 \theta} - 1} \\ &= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}} \\ &= \sqrt{\tan^2 \theta}\end{aligned}$$



# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}
 \sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\
 &= \sqrt{\frac{1}{\cos^2 \theta} - 1} \\
 &= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}} \\
 &= \sqrt{\tan^2 \theta}
 \end{aligned}$$

when  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$  we have  
 $\tan \theta \geq 0$  and so  $\sqrt{\tan^2 \theta} = \tan \theta$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}
 \sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\
 &= \sqrt{\frac{1}{\cos^2 \theta} - 1} \\
 &= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}} \\
 &= \sqrt{\tan^2 \theta} \\
 &= \tan \theta .
 \end{aligned}$$

when  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$  we have  
 $\tan \theta \geq 0$  and so  $\sqrt{\tan^2 \theta} = \tan \theta$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\begin{aligned}
 \sqrt{x^2 - 1} &= \sqrt{\sec^2 \theta - 1} \\
 &= \sqrt{\frac{1}{\cos^2 \theta} - 1} \\
 &= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}} \\
 &= \sqrt{\tan^2 \theta} \\
 &= \tan \theta .
 \end{aligned}$$

when  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$  we have  
 $\tan \theta \geq 0$  and so  $\sqrt{\tan^2 \theta} = \tan \theta$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$

## Definition

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 - 1}$  is given by:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} & \theta &\in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\ \sqrt{x^2 - 1} &= \tan \theta \\ dx &= ? & d\theta \\ \theta &= \operatorname{arcsec} x . \end{aligned}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$

## Definition

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 - 1}$  is given by:

$$x = \sec \theta = \frac{1}{\cos \theta} \quad \theta \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$

$$\sqrt{x^2 - 1} = \tan \theta$$

$$dx = ? \quad d\theta$$

$$\theta = \operatorname{arcsec} x .$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$

## Definition

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 - 1}$  is given by:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} & \theta &\in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\ \sqrt{x^2 - 1} &= \tan \theta \\ dx &= ? & d\theta \\ \theta &= \operatorname{arcsec} x . \end{aligned}$$

# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$

## Definition

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 - 1}$  is given by:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} & \theta &\in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\ \sqrt{x^2 - 1} &= \tan \theta \\ dx &= ? & d\theta \\ \theta &= \operatorname{arcsec} x . \end{aligned}$$



# Trigonometric substitution $x = \sec \theta$ for $\sqrt{x^2 - 1}$

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ :

$$\sqrt{x^2 - 1} = \tan \theta .$$


## Definition

The trigonometric substitution  $x = \sec \theta$ ,  $\theta \in (0, \pi)$  for  $\sqrt{x^2 - 1}$  is given by:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} & \theta &\in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\ \sqrt{x^2 - 1} &= \tan \theta \\ dx &= \frac{\sin \theta}{\cos^2 \theta} d\theta = \sec \theta \tan \theta d\theta \\ \theta &= \operatorname{arcsec} x . \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}, a > 0$ .




## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}, a > 0$ .

•  $x =$

•  $dx =$   
 $\sqrt{x^2 - a^2} =$




## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}, a > 0$ .

•  $x =$


•  $dx =$   
 $\sqrt{x^2 - a^2} =$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}, a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .
- $dx =$   
 $\sqrt{x^2 - a^2} =$




## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .


- $dx =$   
 $\sqrt{x^2 - a^2} =$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .
- $dx = a \sec \theta \tan \theta d\theta$ .  
 $\sqrt{x^2 - a^2} =$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}, a > 0$ .


- $x = a \sec \theta,$

$$0 < \theta < \pi/2 \text{ or}$$

$$\pi < \theta < 3\pi/2.$$

- $dx = a \sec \theta \tan \theta d\theta.$

$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} =$$






## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .

- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} =$$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .

$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| =$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .


- $x = a \sec \theta$ ,

$$0 < \theta < \pi/2 \text{ or}$$

$$\pi < \theta < 3\pi/2.$$

- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .

$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$


$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C \end{aligned}$$



## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a} \right| + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a} \right| + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a} \right| + C \\ &= \ln |x + \sqrt{x^2 - a^2}| - \ln a + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .


$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a} \right| + C \\ &= \ln |x + \sqrt{x^2 - a^2}| - \ln a + C \end{aligned}$$

## Example

Find  $\int \frac{dx}{\sqrt{x^2 - a^2}}$ ,  $a > 0$ .

- $x = a \sec \theta$ ,  
 $0 < \theta < \pi/2$  or  
 $\pi < \theta < 3\pi/2$ .



- $dx = a \sec \theta \tan \theta d\theta$ .

$$\sqrt{x^2 - a^2} = \sqrt{a^2 \sec^2 \theta - a^2} = \sqrt{a^2 \tan^2 \theta} = a |\tan \theta| = a \tan \theta$$

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - a^2}} &= \int \frac{a \sec \theta \tan \theta d\theta}{a \tan \theta} = \int \sec \theta d\theta \\ &= \ln |\sec \theta + \tan \theta| + C = \ln \left| \frac{x}{a} + \frac{\sqrt{x^2 - a^2}}{a} \right| + C \\ &= \ln |x + \sqrt{x^2 - a^2}| + C_1 \end{aligned}$$



# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ , rationalizes  $d\theta, \cos \theta, \sin \theta$ .

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ , rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above?

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? **We get the Euler substitution:**

$$x =$$

## Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = \sec \theta = \frac{1}{\cos \theta}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} \\ &= \frac{1}{\cos(2 \arctan t)} \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \right.$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned} x &= \sec \theta = \frac{1}{\cos \theta} \\ &= \frac{1}{\cos(2 \arctan t)} \end{aligned}$$

$$\left| \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)} \\
 &= \frac{1 + \tan^2(\arctan t)}{1 - \tan^2(\arctan t)}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$



# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)} \\
 &= \frac{1 + \tan^2(\arctan t)}{1 - \tan^2(\arctan t)} \\
 &= \frac{1 + t^2}{1 - t^2}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)} \\
 &= \frac{1 + \tan^2(\arctan t)}{1 - \tan^2(\arctan t)} \\
 &= \frac{1 + t^2}{1 - t^2} = \frac{2 - (1 - t^2)}{1 - t^2}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)} \\
 &= \frac{1 + \tan^2(\arctan t)}{1 - \tan^2(\arctan t)} \\
 &= \frac{1 + t^2}{1 - t^2} = \frac{2 - (1 - t^2)}{1 - t^2} \\
 &= -1 + \frac{2}{1 - t^2}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$\begin{aligned}
 x &= \sec \theta = \frac{1}{\cos \theta} \\
 &= \frac{1}{\cos(2 \arctan t)} \\
 &= \frac{1 + \tan^2(\arctan t)}{1 - \tan^2(\arctan t)} \\
 &= \frac{1 + t^2}{1 - t^2} = \frac{2 - (1 - t^2)}{1 - t^2} \\
 &= -1 + \frac{2}{1 - t^2}
 \end{aligned}
 \quad \left| \quad \cos(2z) = \frac{1 - \tan^2 z}{1 + \tan^2 z} \right.$$

## Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$


---

$$\sqrt{x^2 - 1} =$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\sqrt{x^2 - 1} = \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\begin{aligned} \sqrt{x^2 - 1} &= \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 - t^2)^2}} \end{aligned}$$



# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\begin{aligned} \sqrt{x^2 - 1} &= \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 - t^2)^2}} \quad | \quad (1 + t^2)^2 - (1 - t^2)^2 = ? \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\begin{aligned} \sqrt{x^2 - 1} &= \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 - t^2)^2}} \quad | \quad (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\sqrt{x^2 - 1} = \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1}$$

$$= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 - t^2)^2}}$$

$$= \sqrt{\frac{4t^2}{(1 - t^2)^2}}$$

$$(1 + t^2)^2 - (1 - t^2)^2 = 4t^2$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\begin{aligned} \sqrt{x^2 - 1} &= \sqrt{\left(\frac{1 + t^2}{1 - t^2}\right)^2 - 1} \\ &= \sqrt{\frac{(1 + t^2)^2 - (1 - t^2)^2}{(1 - t^2)^2}} \quad \left| (1 + t^2)^2 - (1 - t^2)^2 = 4t^2 \right. \\ &= \sqrt{\frac{4t^2}{(1 - t^2)^2}} \quad \left| \begin{array}{l} t, 1 - t^2 \text{ have same sign} \\ \text{when } t \in (-\infty, -1) \cup [0, 1) \end{array} \right. \\ &= \frac{2t}{1 - t^2} \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

---

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$


---

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$


---

$$x = \frac{1 + t^2}{1 - t^2}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$


---

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$


---

$$x = \frac{1 + t^2}{1 - t^2}$$

$$(1 - t^2)x = 1 + t^2$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$


---

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$


---

$$x = \frac{1 + t^2}{1 - t^2}$$

$$(1 - t^2)x = 1 + t^2$$

$$(1 + x)t^2 = x - 1$$



# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$x = \frac{1 + t^2}{1 - t^2}$$

$$(1 - t^2)x = 1 + t^2$$

$$(1 + x)t^2 = x - 1$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

---


$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

---


$$x = \frac{1 + t^2}{1 - t^2}$$

$$(1 - t^2)x = 1 + t^2$$

$$(1 + x)t^2 = x - 1$$

$$t^2 = \frac{x - 1}{x + 1}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$x = \frac{1 + t^2}{1 - t^2}$$

$$(1 - t^2)x = 1 + t^2$$

$$(1 + x)t^2 = x - 1$$

$$t^2 = \frac{x - 1}{x + 1}$$

$$t = \pm \sqrt{\frac{x - 1}{x + 1}}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$


---

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$


---

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$


---

$$dx =$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$

$$dx = d\left(-1 + \frac{2}{1 - t^2}\right)$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$

$$\begin{aligned} dx &= d\left(-1 + \frac{2}{1 - t^2}\right) \\ &= ? \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$

$$\begin{aligned} dx &= d\left(-1 + \frac{2}{1 - t^2}\right) \\ &= \frac{4t}{(1 - t^2)^2} dt \end{aligned}$$



# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx, x, \sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta, \cos \theta, \sin \theta$ .

What if we compose the above? We get the Euler substitution:

$$x = -1 + \frac{2}{1 - t^2}$$

$$\sqrt{x^2 - 1} = \frac{2t}{1 - t^2}$$

$$t = \pm \sqrt{\frac{x-1}{x+1}}$$

$$\begin{aligned} dx &= d\left(-1 + \frac{2}{1 - t^2}\right) \\ &= \frac{4t}{(1 - t^2)^2} dt \end{aligned}$$

# Euler substitution $x = \sec \theta$ , $\theta = 2 \arctan t$

- $x = \sec \theta$  transforms  $dx$ ,  $x$ ,  $\sqrt{x^2 - 1}$  to trig form.
- $\theta = 2 \arctan t$ ,  $t \in (-\infty, -1) \cup [0, 1)$  rationalizes  $d\theta$ ,  $\cos \theta$ ,  $\sin \theta$ .

What if we compose the above? We get the Euler substitution:

## Definition

The Euler substitution for  $\sqrt{x^2 - 1}$  corresponding to  $x = \sec \theta$  is given by:

$$\begin{aligned} x &= \frac{1 + t^2}{1 - t^2}, & t &\in (-\infty, -1) \cup [0, 1) \\ \sqrt{x^2 - 1} &= \frac{2t}{1 - t^2} \\ dx &= \frac{4t}{(1 - t^2)^2} dt \\ t &= \pm \frac{\sqrt{x^2 - 1}}{x + 1} \end{aligned}$$

# Rationalizing Substitutions

Some non-rational fractions can be changed into rational fractions by means of appropriate substitutions. In particular, when an integrand contains an expression of the form  $\sqrt[n]{g(x)}$ , the substitution  $u = \sqrt[n]{g(x)}$  may be effective.

## Example

$$\int \frac{\sqrt{x+4}}{x} dx$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = ?$  and  $dx = ?$ .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{?}{?} ?$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x + 4$ , so  $x = ?$  and  $dx = ?$ .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{u}{?} ?$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = ?$  and  $dx = ?$ .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{u}{?} ?$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = ?$  .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{u}{u^2 - 4} ?$$



## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = ?$  .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{u}{u^2 - 4} ?$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\int \frac{\sqrt{x+4}}{x} dx = \int \frac{u}{u^2 - 4} 2u du$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}\int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\ &= 2 \int \frac{u^2}{u^2-4} du\end{aligned}$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}\int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\ &= 2 \int \frac{u^2}{u^2-4} du \\ &= 2 \int \left( 1 + \frac{4}{u^2-4} \right) du\end{aligned}$$

| long division

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}\int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\&= 2 \int \frac{u^2}{u^2-4} du \\&= 2 \int \left( 1 + \frac{4}{u^2-4} \right) du && \left| \begin{array}{l} \text{long division} \end{array} \right. \\&= 2 \int du + 8 \int \frac{du}{u^2-4}\end{aligned}$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}\int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\&= 2 \int \frac{u^2}{u^2-4} du \\&= 2 \int \left( 1 + \frac{4}{u^2-4} \right) du && \left| \text{long division} \right. \\&= 2 \int du + 8 \int \frac{du}{u^2-4} \\&= 2 \int du + 8 \int \left( \frac{\frac{1}{4}}{u-2} - \frac{\frac{1}{4}}{u+2} \right) du && \left| \text{partial fractions} \right.\end{aligned}$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}\int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\&= 2 \int \frac{u^2}{u^2-4} du \\&= 2 \int \left( 1 + \frac{4}{u^2-4} \right) du && \left| \begin{array}{l} \text{long division} \end{array} \right. \\&= 2 \int du + 8 \int \frac{du}{u^2-4} \\&= 2 \int du + 8 \int \left( \frac{\frac{1}{4}}{u-2} - \frac{\frac{1}{4}}{u+2} \right) du && \left| \begin{array}{l} \text{partial fractions} \end{array} \right. \\&= 2u + 2(\ln |u-2| - \ln |u+2|) + C\end{aligned}$$

## Example

Let  $u = \sqrt{x+4}$ . Then  $u^2 = x+4$ , so  $x = u^2 - 4$  and  $dx = 2u du$ .

$$\begin{aligned}
 \int \frac{\sqrt{x+4}}{x} dx &= \int \frac{u}{u^2-4} 2u du \\
 &= 2 \int \frac{u^2}{u^2-4} du \\
 &= 2 \int \left( 1 + \frac{4}{u^2-4} \right) du && \left| \text{long division} \right. \\
 &= 2 \int du + 8 \int \frac{du}{u^2-4} \\
 &= 2 \int du + 8 \int \left( \frac{\frac{1}{4}}{u-2} - \frac{\frac{1}{4}}{u+2} \right) du && \left| \text{partial fractions} \right. \\
 &= 2u + 2(\ln |u-2| - \ln |u+2|) + C \\
 &= 2\sqrt{x+4} + 2 \ln \left| \frac{\sqrt{x+4}-2}{\sqrt{x+4}+2} \right| + C
 \end{aligned}$$