#### Calculus II

# Power series expansion of rational functions with linear denominator, part 2

**Todor Milev** 

2019

$$\frac{1}{2+x} \quad = \quad \frac{1}{2\left(1+\frac{x}{2}\right)}$$

$$\frac{1}{2+x} = \frac{1}{2\left(1+\frac{x}{2}\right)}$$
$$= \frac{1}{2} \cdot \frac{1}{\left(1-\left(-\frac{x}{2}\right)\right)}$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} (-\frac{x}{2})^n \quad | \text{ if \& only if } |-\frac{x}{2}| < 1$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} (-\frac{x}{2})^{n} \quad | \text{ if & only if } |-\frac{x}{2}| < 1$$

$$= \sum_{n=0}^{\infty} \frac{?}{?} ?$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} (-\frac{x}{2})^{n} \quad | \text{ if & only if } |-\frac{x}{2}| < 1$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{?}?$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^{n} \quad \left| \begin{array}{c} \text{if \& only if } \\ |-\frac{x}{2}| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{n}$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^{n} \quad \left| \begin{array}{c} \text{if \& only if } \\ |-\frac{x}{2}| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2} x^{n}$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^n \quad \left| \begin{array}{c} \text{if \& only if } \\ |-\frac{x}{2}| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{?} x^n$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^n \quad \left| \begin{array}{c} \text{if & endy if } \\ |-\frac{x}{2}| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$\frac{1}{2+x} = \frac{1}{2(1+\frac{x}{2})}$$

$$= \frac{1}{2} \cdot \frac{1}{(1-(-\frac{x}{2}))} = \frac{1}{2} \sum_{n=0}^{\infty} (-\frac{x}{2})^n \quad | \text{ if & only if } | -\frac{x}{2}| < 1$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} + \dots$$

Find a power series representation for  $\frac{1}{x+2}$ .

$$\frac{1}{2+x} = \frac{1}{2\left(1+\frac{x}{2}\right)}$$

$$= \frac{1}{2} \cdot \frac{1}{\left(1-\left(-\frac{x}{2}\right)\right)} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^n \quad \left| \begin{array}{c} \text{if & 6 only if } \\ \left|-\frac{x}{2}\right| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} + \dots$$

To find interval of convergence:





Find a power series representation for  $\frac{1}{x+2}$ .

$$\frac{1}{2+x} = \frac{1}{2\left(1+\frac{x}{2}\right)}$$

$$= \frac{1}{2} \cdot \frac{1}{\left(1-\left(-\frac{x}{2}\right)\right)} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^{n} \quad \left| \begin{array}{c} \text{if & & only if } \\ \left|-\frac{x}{2}\right| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}} x^{n}$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^{2}}{8} - \frac{x^{3}}{16} + \dots$$

To find interval of convergence:

$$\left|-\frac{x}{2}\right| < 1$$

Find a power series representation for  $\frac{1}{x+2}$ .

$$\frac{1}{2+x} = \frac{1}{2\left(1+\frac{x}{2}\right)}$$

$$= \frac{1}{2} \cdot \frac{1}{\left(1-\left(-\frac{x}{2}\right)\right)} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^n \quad \left| \begin{array}{c} \text{if \& only if } \\ \left|-\frac{x}{2}\right| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} + \dots$$

To find interval of convergence:

$$\begin{vmatrix} -\frac{x}{2} \\ |x| & < & 2 \end{vmatrix}$$

Find a power series representation for  $\frac{1}{x+2}$ .

$$\frac{1}{2+x} = \frac{1}{2\left(1+\frac{x}{2}\right)}$$

$$= \frac{1}{2} \cdot \frac{1}{\left(1-\left(-\frac{x}{2}\right)\right)} = \frac{1}{2} \sum_{n=0}^{\infty} \left(-\frac{x}{2}\right)^{n} \quad \left| \begin{array}{c} \text{if & anly if } \\ \left|-\frac{x}{2}\right| < 1 \end{array} \right|$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}} x^{n}$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^{2}}{8} - \frac{x^{3}}{16} + \dots$$

To find interval of convergence:

$$\left| -\frac{x}{2} \right| < 1$$

$$|x| < 2$$

Therefore the interval of convergence is  $x \in (-2, 2)$ .

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$
$$= x^3$$
?

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$

$$= x^3 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

if & only if 
$$|x| < 2$$

Find a power series representation for  $\frac{x^3}{x+2}$ .

$$\frac{x^{3}}{x+2} = x^{3} \cdot \frac{1}{x+2}$$

$$= x^{3} \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}} x^{n}$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}}?$$

if & only if |x| < 2

$$\frac{x^{3}}{x+2} = x^{3} \cdot \frac{1}{x+2}$$

$$= x^{3} \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}} x^{n}$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{2^{n+1}} x^{n+3}$$

$$| \text{ if & only if } |x| < 2$$

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$

$$= x^3 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^{n+3}$$

$$= \frac{x^3}{2} - \frac{x^4}{4} + \frac{x^5}{8} - \frac{x^6}{16} + \cdots$$

Find a power series representation for  $\frac{x^3}{x+2}$ .

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$

$$= x^3 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^{n+3}$$

$$= \frac{x^3}{2} - \frac{x^4}{4} + \frac{x^5}{8} - \frac{x^6}{16} + \cdots$$

• Another way to write this is  $\frac{x^3}{x+2} = \sum_{n=3}^{\infty} \frac{(-1)^{n-1}}{2^{n-2}} x^n$ .

$$\frac{x^3}{x+2} = x^3 \cdot \frac{1}{x+2}$$

$$= x^3 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^n$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{n+1}} x^{n+3}$$

$$= \frac{x^3}{2} - \frac{x^4}{4} + \frac{x^5}{8} - \frac{x^6}{16} + \cdots$$

- Another way to write this is  $\frac{x^3}{x+2} = \sum_{n=3}^{\infty} \frac{(-1)^{n-1}}{2^{n-2}} x^n$ .
- The interval of convergence is again  $x \in (-2, 2)$ .