

# Precalculus

## Factoring quadratic polynomials

Todor Milev

2019

# Outline

## 1 Factoring quadratics

# Outline

1 Factoring quadratics

2 Vietas' formulas

## Definition ((Partial) Factorization)

To (partially) factor a polynomial means to rewrite it as a product of two polynomials of smaller degree.

## Definition ((Partial) Factorization)

To (partially) factor a polynomial means to rewrite it as a product of two polynomials of smaller degree.

## Example

$$\begin{aligned}x^2 - 1 &= (x - 1)(x + 1) \\ x^3 + 8 &= (x + 2)(x^2 - 2x + 4)\end{aligned}$$

## Theorem

*The quadratic  $ax^2 + bx + c$  factors as follows.*

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

*where  $x_1$  and  $x_2$  are the roots of the quadratic, given by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The quadratic  $ax^2 + bx + c$  factors as follows.*

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

*where  $x_1$  and  $x_2$  are the roots of the quadratic, given by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The quadratic  $ax^2 + bx + c$  factors as follows.*

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

*where  $x_1$  and  $x_2$  are the roots of the quadratic, given by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = ( \quad ? \quad )( \quad ? \quad )$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$3x^2 + 8x - 11 = (3x + p)(x + q)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$



## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + \color{red}{3xq} + \color{red}{px} + pq \\ &= 3x^2 + x(\color{red}{3q} + \color{red}{p}) + pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$8 = 3q + p$$

$$-11 = pq$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + ?)(x + ?)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= ? \\ q &= ? \end{aligned}$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$3x^2 + 8x - 11 = (3x + 11)(x - 1)$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$



$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- If there is a factorization using integers, it should be of the form

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + p)(x + q) \\ &= 3x^2 + 3xq + px + pq \\ &= 3x^2 + x(3q + p) + pq \end{aligned}$$

(Vieta's formulas) This means that :

$$\begin{aligned} 8 &= 3q + p \\ -11 &= pq \end{aligned}$$

$p, q$  must be divisors of 11:  $\pm 1, \pm 11$

$$\begin{aligned} p &= 11 \\ q &= -1 \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3}$$



$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} = \begin{cases} \frac{-8 + 14}{6} \\ \frac{-8 - 14}{6} \end{cases} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} = \begin{cases} \frac{-8 + 14}{6} = \frac{6}{6} = 1 \\ \frac{-8 - 14}{6} \end{cases} \end{aligned}$$



$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} = \begin{cases} \frac{-8 + 14}{6} = \frac{6}{6} = 1 \\ \frac{-8 - 14}{6} = -\frac{22}{6} = -\frac{11}{3} \end{cases} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} = \begin{cases} \frac{-8 + 14}{6} = \frac{6}{6} = 1 \\ \frac{-8 - 14}{6} = -\frac{22}{6} = -\frac{11}{3} \end{cases} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \text{ where } x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

## Example

Factor the polynomial. If possible, guess the factorization.

$$\begin{aligned} 3x^2 + 8x - 11 &= (3x + 11)(x - 1) \\ &= 3\left(x - \left(-\frac{11}{3}\right)\right)(x - 1) \end{aligned}$$

- What if we can't guess the factorization?
- Use the formulas for  $x_1, x_2$ .

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-8 \pm \sqrt{8^2 - 4 \cdot 3 \cdot (-11)}}{2 \cdot 3} \\ &= \frac{-8 \pm \sqrt{64 + 132}}{6} = \frac{-8 \pm \sqrt{196}}{6} \\ &= \frac{-8 \pm 14}{6} = \begin{cases} \frac{-8 + 14}{6} = \frac{6}{6} = 1 \\ \frac{-8 - 14}{6} = -\frac{22}{6} = -\frac{11}{3} \end{cases} \end{aligned}$$

## Proposition (Vieta's formulas)

*Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:*

$$a(x - x_1)(x - x_2) = ax^2 + bx + c$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic function with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \end{aligned}$$



## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - \textcolor{red}{a}x x_2 - \textcolor{red}{a}x_1 \textcolor{red}{x} + a(-x_1)(-x_2) &= ax^2 + bx + c \\ ax^2 - \textcolor{red}{a}(x_2 + x_1)\textcolor{red}{x} + ax_1 x_2 &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

*Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:*

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

*Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:*

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

*Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:*

$$\begin{aligned} a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned}a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \\ax_1x_2 &= c\end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned}a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \\ax_1x_2 &= c \\x_1x_2 &= \frac{c}{a}\end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned}a(x - x_1)(x - x_2) &= ax^2 + bx + c \\ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \\ax_1x_2 &= c \\x_1x_2 &= \frac{c}{a} \\-a(x_1 + x_2) &= b\end{aligned}$$

## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned}
 a(x - x_1)(x - x_2) &= ax^2 + bx + c \\
 ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\
 ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \\
 ax_1x_2 &= c \\
 x_1x_2 &= \frac{c}{a} \\
 -a(x_1 + x_2) &= b \\
 x_1 + x_2 &= -\frac{b}{a}
 \end{aligned}$$



## Proposition (Vieta's formulas)

Let  $ax^2 + bx + c$  be a quadratic functions with zeros  $x_1$  and  $x_2$ . Then:

$$\begin{aligned}
 a(x - x_1)(x - x_2) &= ax^2 + bx + c \\
 ax^2 - axx_2 - ax_1x + a(-x_1)(-x_2) &= ax^2 + bx + c \\
 ax^2 - a(x_2 + x_1)x + ax_1x_2 &= ax^2 + bx + c \\
 ax_1x_2 &= c \\
 x_1x_2 &= \frac{c}{a} \\
 -a(x_1 + x_2) &= b \\
 x_1 + x_2 &= -\frac{b}{a}
 \end{aligned}$$

The last two formulas are called Vieta's formulas (after François Viète (1540-1603), Latinized name: Franciscus Vieta).

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$x_1 + x_2 = -\frac{b}{a}$$

$$x_1 x_2 = \frac{c}{a}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$x_1 + x_2 = -\frac{b}{a}$$

$$x_1 x_2 = \frac{c}{a}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .
- Therefore the pair  $x_1, x_2$  is  $\pm 1, \pm 6$  or  $\pm 2, \pm 3$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .
- Therefore the pair  $x_1, x_2$  is  $\pm 1, \pm 6$  or  $\pm 2, \pm 3$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .
- Therefore the pair  $x_1, x_2$  is  $\pm 1, \pm 6$  or  $\pm 2, \pm 3$ .
- The sum of the two roots:  $x_1 + x_2 = -5$

$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .
- Therefore the pair  $x_1, x_2$  is  $\pm 1, \pm 6$  or  $\pm 2, \pm 3$ .
- The sum of the two roots:  $x_1 + x_2 = -5$



$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

$$\begin{aligned}x_1 + x_2 &= -\frac{b}{a} \\ x_1 x_2 &= \frac{c}{a}\end{aligned}$$

Vieta's formulas

## Example

Factor the quadratic.

$$x^2 + 5x + 6 = (x + 2)(x + 3)$$

- The product of the two roots:  $x_1 x_2 = 6$ .
- The divisors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$ .
- Therefore the pair  $x_1, x_2$  is  $\pm 1, \pm 6$  or  $\pm 2, \pm 3$ .
- The sum of the two roots:  $x_1 + x_2 = -5$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

- $(x - 1)(x - 1)$
- $(x + 1)(x + 1)$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

- $(x - 1)(x - 1)$   
 $(x + 1)(x + 1)$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\begin{cases} x_1 x_2 = \frac{c}{a} \\ x_1 + x_2 = -\frac{b}{a} \end{cases}$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-3 \pm \sqrt{5}}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-3 \pm \sqrt{5}}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic.

$$x^2 + 3x + 1 = \left( x - \left( \frac{-3 + \sqrt{5}}{2} \right) \right) \left( x - \left( \frac{-3 - \sqrt{5}}{2} \right) \right)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-3 \pm \sqrt{5}}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .



$$ax^2 + bx + c = a(x - x_1)(x - x_2), \quad \left| \begin{array}{lcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

- $(x - 1)(x - 1)$   
 $(x + 1)(x + 1)$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .

- $(x - 1)(x - 1)$   
 $(x + 1)(x + 1)$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \quad \left| \begin{array}{lcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $\begin{array}{l} (x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1 \\ (x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1 \end{array}$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2), \quad \left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?) (x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $\begin{array}{l} (x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1 \\ (x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1 \end{array}$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$



$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-1 \pm \sqrt{-3}}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{lcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm \sqrt{3}i}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = (x + ?)(x + ?)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm \sqrt{3}i}{2} \end{aligned}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$

$$\left| \begin{array}{rcl} x_1 x_2 & = & \frac{c}{a} \\ x_1 + x_2 & = & -\frac{b}{a} \end{array} \right.$$

## Example

Factor the quadratic, using complex numbers if needed.

$$x^2 + x + 1 = \left( x - \left( \frac{-1 + \sqrt{3}i}{2} \right) \right) \left( x - \left( \frac{-1 - \sqrt{3}i}{2} \right) \right)$$

- The product of the two roots:  $x_1 x_2 = 1$ .
- Integer options:  $x_1 = 1, x_2 = 1$  and  $x_1 = -1, x_2 = -1$ .
- $(x - 1)(x - 1) = (x - 1)^2 = x^2 - 2x + 1$   
 $(x + 1)(x + 1) = (x + 1)^2 = x^2 + 2x + 1$  both don't work.
- $\Rightarrow$  No easy factorization; must use quadratic formula.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-1 \pm \sqrt{1^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} \\ &= \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm \sqrt{3}i}{2} \end{aligned}$$