

# Calculus I

## Derivatives: linearity, product and quotient rules

Todor Milev

2019

## Theorem (The Power Rule (General Version))

*If  $n$  is any real number, then*

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

### Example (Power Rule, negative exponent)

Differentiate  $y = \frac{1}{x}$ .

### Example (Power Rule, negative exponent)

Differentiate  $y = \frac{1}{x}$ .

$$y = x^{-1}.$$

## Example (Power Rule, negative exponent)

Differentiate  $y = \frac{1}{x}$ .

$$y = x^{-1}.$$

Power Rule:  $\frac{dy}{dx} = ?$

### Example (Power Rule, negative exponent)

Differentiate  $y = \frac{1}{x}$ .

$$y = x^{-1}.$$

Power Rule:  $\frac{dy}{dx} = (-1)x^{-2}$

### Example (Power Rule, negative exponent)

Differentiate  $y = \frac{1}{x}$ .


$$y = x^{-1}.$$

Power Rule:  $\frac{dy}{dx} = (-1)x^{-2}$

$$= -\frac{1}{x^2}.$$

## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .






## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .




## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .


$$f'(x) = \frac{1}{3}x^{\frac{1}{3}-1}$$



## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .




$$\begin{aligned} f'(x) &= \frac{1}{3}x^{\frac{1}{3}-1} \\ &= \frac{1}{3}x^{-\frac{2}{3}} \end{aligned}$$

## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .




$$\begin{aligned} f'(x) &= \frac{1}{3}x^{\frac{1}{3}-1} \\ &= \frac{1}{3}x^{-\frac{2}{3}} \\ &= \frac{1}{3\sqrt[3]{x^2}}. \end{aligned}$$

## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .




$$\begin{aligned} f'(x) &= \frac{1}{3}x^{\frac{1}{3}-1} \\ &= \frac{1}{3}x^{-\frac{2}{3}} \\ &= \frac{1}{3\sqrt[3]{x^2}}. \\ f'(1) &= \frac{1}{3}. \end{aligned}$$

## Example (Calculating the tangent line using the Power Rule)

Find an equation for the tangent line to the cubic  $y = \sqrt[3]{x}$  at the point  $P = (1, 1)$ .

Here  $a = 1$  and  $f(x) = \sqrt[3]{x} = x^{\frac{1}{3}}$ .



$$\begin{aligned}f'(x) &= \frac{1}{3}x^{\frac{1}{3}-1} \\&= \frac{1}{3}x^{-\frac{2}{3}} \\&= \frac{1}{3\sqrt[3]{x^2}}. \\f'(1) &= \frac{1}{3}.\end{aligned}$$

Point-slope form:  $y - 1 = \frac{1}{3}(x - 1)$ , or  
 $y = \frac{1}{3}x + \frac{2}{3}$ .

### Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

### Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

$$y = x^{\frac{5}{6}}.$$



## Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

$$y = x^{\frac{5}{6}}.$$

Power Rule:  $\frac{dy}{dx} = ?$

## Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

$$y = x^{\frac{5}{6}}.$$

Power Rule:  $\frac{dy}{dx} = \frac{5x^{-\frac{1}{6}}}{6}$

## Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

$$y = x^{\frac{5}{6}}.$$

Power Rule:  $\frac{dy}{dx} = \frac{5x^{-\frac{1}{6}}}{6}$   
 $= ?$

## Example (Power Rule, fractional exponent)

Differentiate  $y = \sqrt[6]{x^5}$ .

$$y = x^{\frac{5}{6}}.$$

Power Rule:  $\frac{dy}{dx} = \frac{5x^{-\frac{1}{6}}}{6}$   
 $= \frac{5}{6\sqrt[6]{x}}.$

## Theorem (The Constant Multiple Rule)

*If  $c$  is a constant and  $f$  is a differentiable function, then*

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

Proof.



## Theorem (The Constant Multiple Rule)

*If  $c$  is a constant and  $f$  is a differentiable function, then*

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .



## Theorem (The Constant Multiple Rule)

*If  $c$  is a constant and  $f$  is a differentiable function, then*

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\text{Then } g'(x) = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$$



## Theorem (The Constant Multiple Rule)

If  $c$  is a constant and  $f$  is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\begin{aligned}\text{Then } g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h}\end{aligned}$$





## Theorem (The Constant Multiple Rule)

If  $c$  is a constant and  $f$  is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\begin{aligned}\text{Then } g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{c(f(x+h) - f(x))}{h}\end{aligned}$$



## Theorem (The Constant Multiple Rule)

If  $c$  is a constant and  $f$  is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\begin{aligned}\text{Then } g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{c(f(x+h) - f(x))}{h} \\ \text{Limit Law 3: } &= c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}\end{aligned}$$



## Theorem (The Constant Multiple Rule)

If  $c$  is a constant and  $f$  is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\begin{aligned}\text{Then } g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{c(f(x+h) - f(x))}{h}\end{aligned}$$

$$\begin{aligned}\text{Limit Law 3: } &= c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= c?\end{aligned}$$



## Theorem (The Constant Multiple Rule)

If  $c$  is a constant and  $f$  is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x).$$

### Proof.

Let  $g(x) = cf(x)$ .

$$\begin{aligned}\text{Then } g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{c(f(x+h) - f(x))}{h}\end{aligned}$$

$$\begin{aligned}\text{Limit Law 3: } &= c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= c f'(x).\end{aligned}$$



## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5) .$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5) .$$

$$\frac{dy}{dx} = \frac{d}{dx} \left[ \left(\frac{2}{7}\right) (x^5) \right]$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5).$$

$$\frac{dy}{dx} = \frac{d}{dx} \left[ \left(\frac{2}{7}\right) (x^5) \right]$$

$$\text{Constant Multiple Rule: } = \left(\frac{2}{7}\right) \frac{d}{dx} (x^5)$$



## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5).$$

$$\frac{dy}{dx} = \frac{d}{dx} \left[ \left(\frac{2}{7}\right) (x^5) \right]$$

Constant Multiple Rule:  $= \left(\frac{2}{7}\right) \frac{d}{dx} (x^5)$

$$= \left(\frac{2}{7}\right) (\text{?})$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5).$$

$$\frac{dy}{dx} = \frac{d}{dx} \left[ \left(\frac{2}{7}\right) (x^5) \right]$$

Constant Multiple Rule:  $= \left(\frac{2}{7}\right) \frac{d}{dx} (x^5)$

$$= \left(\frac{2}{7}\right) (5x^4)$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $y = \frac{2x^5}{7}$ .

$$y = \left(\frac{2}{7}\right) (x^5).$$

$$\frac{dy}{dx} = \frac{d}{dx} \left[ \left(\frac{2}{7}\right) (x^5) \right]$$

$$\begin{aligned}\text{Constant Multiple Rule: } &= \left(\frac{2}{7}\right) \frac{d}{dx} (x^5) \\ &= \left(\frac{2}{7}\right) (5x^4) \\ &= \frac{10x^4}{7}.\end{aligned}$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

$$\frac{du}{dx} = \frac{d}{dx} [(-1)(x)]$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

$$\frac{du}{dx} = \frac{d}{dx} [(-1)(x)]$$

$$\text{Constant Multiple Rule: } = (-1) \frac{d}{dx} (x)$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

$$\frac{du}{dx} = \frac{d}{dx} [(-1)(x)]$$

Constant Multiple Rule:  $= (-1) \frac{d}{dx} (x)$

$$= (-1) (?)$$



## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

$$\frac{du}{dx} = \frac{d}{dx} [(-1)(x)]$$

$$\begin{aligned}\text{Constant Multiple Rule: } &= (-1) \frac{d}{dx} (x) \\ &= (-1) (1)\end{aligned}$$

## Example (Constant Multiple Rule, Power Rule)

Find the derivative of  $u = -x$ .

$$u = (-1)(x).$$

$$\frac{du}{dx} = \frac{d}{dx} [(-1)(x)]$$

$$\begin{aligned}\text{Constant Multiple Rule: } &= (-1) \frac{d}{dx} (x) \\ &= (-1)(1) \\ &= -1.\end{aligned}$$

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

$$\frac{dt}{dx} = \frac{d}{dx} \left[ (2\pi) (x^{-4}) \right]$$

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

$$\frac{dt}{dx} = \frac{d}{dx} \left[ (2\pi) (x^{-4}) \right]$$

$$\text{Constant Multiple Rule: } = (2\pi) \frac{d}{dx} (x^{-4})$$

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

$$\frac{dt}{dx} = \frac{d}{dx} \left[ (2\pi) (x^{-4}) \right]$$

Constant Multiple Rule:  $= (2\pi) \frac{d}{dx} (x^{-4})$

$$= (2\pi) ( ? )$$

## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

$$\frac{dt}{dx} = \frac{d}{dx} \left[ (2\pi) (x^{-4}) \right]$$

$$\begin{aligned} \text{Constant Multiple Rule: } &= (2\pi) \frac{d}{dx} (x^{-4}) \\ &= (2\pi) (-4x^{-5}) \end{aligned}$$



## Example (Constant Multiple Rule, Power Rule, Negative Exponent)

Find the derivative of  $t = \frac{2\pi}{x^4}$ .

$$t = (2\pi) (x^{-4}) .$$

$$\frac{dt}{dx} = \frac{d}{dx} \left[ (2\pi) (x^{-4}) \right]$$

$$\begin{aligned} \text{Constant Multiple Rule: } &= (2\pi) \frac{d}{dx} (x^{-4}) \\ &= (2\pi) (-4x^{-5}) \\ &= -\frac{8\pi}{x^5} . \end{aligned}$$

## Theorem (The Sum Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

Proof.



## Theorem (The Sum Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

## Proof.

Let  $F(x) = f(x) + g(x)$ .



## Theorem (The Sum Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

## Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\text{Then } F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h}\end{aligned}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$





## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

## Proof.

Let  $F(x) = f(x) + g(x)$ .

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$

$$\text{Limit Law 1: } = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

$$\text{Let } F(x) = f(x) + g(x).$$

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$

$$\begin{aligned}\text{Limit Law 1: } &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= f'(x) + g'(x).\end{aligned}$$



## Theorem (The Sum Rule)

If  $f$  and  $g$  are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x).$$

### Proof.

$$\text{Let } F(x) = f(x) + g(x).$$

$$\begin{aligned}\text{Then } F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[ \frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right]\end{aligned}$$

$$\begin{aligned}\text{Limit Law 1: } &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= f'(x) + g'(x).\end{aligned}$$



The Sum Rule can be extended to any number of summands. For instance, using the theorem twice, we get

$$(f + g + h)' = [(f + g) + h]' = (f + g)' + h' = f' + g' + h'.$$

The Sum Rule can be extended to any number of summands. For instance, using the theorem twice, we get

$$(f + g + h)' = [(f + g) + h]' = (f + g)' + h' = f' + g' + h'.$$

By writing  $f - g$  as  $f + (-1)g$  and applying the Sum Rule and the Constant Multiple Rule, we get

### Theorem (The Difference Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x).$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the Power Rule can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\text{Then } \frac{dy}{dx} =$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the Power Rule can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\text{Then } \frac{dy}{dx} = \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right)$$

The Constant Multiple Rule, the **Sum Rule**, the **Difference Rule**, and the Power Rule can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5)\end{aligned}$$



The **Constant Multiple Rule**, the Sum Rule, the Difference Rule, and the Power Rule can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5)\end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (\text{?}) + 2\sqrt{3} (\text{?}) - 4 (\text{?}) + \frac{1}{8} (\text{?}) - (\text{?}) \end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} \left( ? \right) - 4 \left( ? \right) + \frac{1}{8} (?) - (?)\end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} \left( ? \right) - 4 \left( ? \right) + \frac{1}{8} (?) - (?) \end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (?) + \frac{1}{8} (?) - (?) \end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 \left( ? \right) + \frac{1}{8} (?) - (?) \end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (?) - (?)\end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (?) - (?) \end{aligned}$$



The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned} \text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (1) - (?) \end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (1) - (?)\end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the **Power Rule** can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (1) - (0)\end{aligned}$$

The Constant Multiple Rule, the Sum Rule, the Difference Rule, and the Power Rule can be combined to differentiate any polynomial.

### Example (Derivative of a Polynomial)

$$\text{If } y = x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5,$$

$$\begin{aligned}\text{Then } \frac{dy}{dx} &= \frac{d}{dx} \left( x^{16} + 2\sqrt{3}x^7 - 4x^3 + \frac{x}{8} - 5 \right) \\ &= \frac{d}{dx} (x^{16}) + \frac{d}{dx} (2\sqrt{3}x^7) - \frac{d}{dx} (4x^3) + \frac{d}{dx} \left( \frac{x}{8} \right) - \frac{d}{dx} (5) \\ &= \frac{d}{dx} (x^{16}) + 2\sqrt{3} \frac{d}{dx} (x^7) - 4 \frac{d}{dx} (x^3) + \frac{1}{8} \frac{d}{dx} (x) - \frac{d}{dx} (5) \\ &= (16x^{15}) + 2\sqrt{3} (7x^6) - 4 (3x^2) + \frac{1}{8} (1) - (0) \\ &= 16x^{15} + 14\sqrt{3}x^6 - 12x^2 + \frac{1}{8}.\end{aligned}$$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}.$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}.$

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3 \frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3? - ?$$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3 \frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - ?$$



## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - ?$$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}.$

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}.$

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3 \frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3 \frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Power Rule:  $= 3 \left( ? \right) - \left( ? \right)$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3 \frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Power Rule:  $= 3 \left( -\frac{1}{2} x^{-\frac{3}{2}} \right) - \left( ? \right)$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3 \frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Power Rule:  $= 3 \left( -\frac{1}{2} x^{-\frac{3}{2}} \right) - \left( ? \right)$

## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3 \frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Power Rule:  $= 3 \left( -\frac{1}{2} x^{-\frac{3}{2}} \right) - \left( -\frac{2}{3} x^{-\frac{5}{3}} \right)$



## Example (Difference Rule, Negative Fractional Exponents)

Differentiate  $v = \frac{3\sqrt{x} - \sqrt[3]{x}}{x}$ .

$$v = 3\frac{\sqrt{x}}{x} - \frac{\sqrt[3]{x}}{x}$$

$$v = 3x^{-\frac{1}{2}} - x^{-\frac{2}{3}}.$$

Difference Rule:  $\frac{dv}{dx} = \frac{d}{dx} \left( 3x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Constant Multiple Rule:  $= 3\frac{d}{dx} \left( x^{-\frac{1}{2}} \right) - \frac{d}{dx} \left( x^{-\frac{2}{3}} \right)$

Power Rule:  $= 3 \left( -\frac{1}{2}x^{-\frac{3}{2}} \right) - \left( -\frac{2}{3}x^{-\frac{5}{3}} \right)$

$$= \frac{2}{3}x^{-\frac{5}{3}} - \frac{3}{2}x^{-\frac{3}{2}}.$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h}$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \end{aligned}$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x (a^h - 1)}{h}\end{aligned}$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x (a^h - 1)}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - a^0}{h}\end{aligned}$$

# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x (a^h - 1)}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - a^0}{h}\end{aligned}$$



# Derivatives of Exponential Functions

Compute the derivative of  $f(x) = a^x$  using the definition:

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\&= \lim_{h \rightarrow 0} \frac{a^x (a^h - 1)}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} \\&= a^x \lim_{h \rightarrow 0} \frac{a^h - a^0}{h} \\&= a^x f'(0).\end{aligned}$$

We have shown that, if  $f(x) = a^x$  is differentiable at 0, then it is differentiable everywhere, and

$$f'(x) = f'(0)a^x.$$

We have shown that, if  $f(x) = a^x$  is differentiable at 0, then it is differentiable everywhere, and

$$f'(x) = f'(0)a^x.$$

We leave the following theorem without proof.

## Theorem

*Let  $a$  be a positive number and let  $f(x) = a^x$ . Then the limit*

$$f'(0) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$$

*exists.*

We have shown that, if  $f(x) = a^x$  is differentiable at 0, then it is differentiable everywhere, and

$$f'(x) = f'(0)a^x.$$

We leave the following theorem without proof.

### Theorem

*Let  $a$  be a positive number and let  $f(x) = a^x$ . Then the limit*

$$f'(0) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$$

*exists.*



We will later show that

$$f'(0) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h} = \ln(a).$$

Here,  $\ln$  is the natural logarithm function.

If  $f(x) = a^x$ , then  $f'(x) = f'(0)a^x$ .

The formula above is simplest when  $f'(0) = 1$ . Since  $\lim_{h \rightarrow 0} \frac{2^h - 1}{h} \approx 0.69$  and  $\lim_{h \rightarrow 0} \frac{3^h - 1}{h} \approx 1.10$ , we expect there is a number  $a$  between 2 and 3 such that  $\lim_{h \rightarrow 0} \frac{a^h - 1}{h} = 1$ .





If  $f(x) = a^x$ , then  $f'(x) = f'(0)a^x$ .

The formula above is simplest when  $f'(0) = 1$ . Since  $\lim_{h \rightarrow 0} \frac{2^h - 1}{h} \approx 0.69$  and  $\lim_{h \rightarrow 0} \frac{3^h - 1}{h} \approx 1.10$ , we expect there is a number  $a$  between 2 and 3 such that  $\lim_{h \rightarrow 0} \frac{a^h - 1}{h} = 1$ .

### Definition (e)

$e$  is the number such that  $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$ .






If  $f(x) = a^x$ , then  $f'(x) = f'(0)a^x$ .

The formula above is simplest when  $f'(0) = 1$ . Since  $\lim_{h \rightarrow 0} \frac{2^h - 1}{h} \approx 0.69$  and  $\lim_{h \rightarrow 0} \frac{3^h - 1}{h} \approx 1.10$ , we expect there is a number  $a$  between 2 and 3 such that  $\lim_{h \rightarrow 0} \frac{a^h - 1}{h} = 1$ .

### Definition (e)

$e$  is the number such that  $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$ .



## Definition (Natural Exponential Function)

$e^x$  is called the natural exponential function. Its derivative is

$$\frac{d}{dx} (e^x) = e^x.$$



## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

$$\frac{dy}{dx} = \frac{d}{dx}(e^x) + \frac{d}{dx}(x^7)$$

## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^x) + \frac{d}{dx}(x^7) \\ &= ? + ?\end{aligned}$$

## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^x) + \frac{d}{dx}(x^7) \\ &= e^x + ?\end{aligned}$$

## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^x) + \frac{d}{dx}(x^7) \\ &= e^x + ?\end{aligned}$$

## Example (Derivative of a Polynomial and the Natural Exponential Function)

Differentiate  $y = e^x + x^7$ .

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(e^x) + \frac{d}{dx}(x^7) \\ &= e^x + 7x^6.\end{aligned}$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) =$$

$$(fg)(x) = f(x)g(x) =$$

$$g'(x) =$$

$$(fg)'(x) =$$

$$f'(x)g'(x) =$$



We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = ?$$

$$(fg)(x) = f(x)g(x) =$$

$$g'(x) =$$

$$(fg)'(x) =$$

$$f'(x)g'(x) =$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) =$$

$$g'(x) =$$

$$(fg)'(x) =$$

$$f'(x)g'(x) =$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = ?$$

$$f'(x)g'(x) =$$

$$(fg)(x) = f(x)g(x) =$$

$$(fg)'(x) =$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = 2x.$$

$$f'(x)g'(x) =$$

$$(fg)(x) = f(x)g(x) =$$

$$(fg)'(x) =$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) =$$

$$g'(x) = 2x.$$

$$(fg)'(x) =$$

$$f'(x)g'(x) = ?$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) =$$

$$g'(x) = 2x.$$

$$(fg)'(x) =$$

$$f'(x)g'(x) = 2x.$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = 2x.$$

$$f'(x)g'(x) = 2x.$$

$$(fg)(x) = f(x)g(x) = ?$$

$$(fg)'(x) =$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = 2x.$$

$$f'(x)g'(x) = 2x.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$(fg)'(x) =$$



We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = 2x.$$

$$f'(x)g'(x) = 2x.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$(fg)'(x) = ?$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$g'(x) = 2x.$$

$$f'(x)g'(x) = 2x.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$(fg)'(x) = 3x^2.$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$g'(x) = 2x.$$

$$(fg)'(x) = 3x^2.$$

$$f'(x)g'(x) = 2x.$$

Therefore

$$f'(x)g'(x) \neq (fg)'(x) \quad .$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$g'(x) = 2x.$$

$$(fg)'(x) = 3x^2.$$

$$f'(x)g'(x) = 2x.$$

Therefore

$$f'(x)g'(x) \neq (fg)'(x) \quad .$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$g'(x) = 2x.$$

$$(fg)'(x) = 3x^2.$$

$$f'(x)g'(x) = 2x.$$

Therefore

$$f'(x)g'(x) \neq (fg)'(x) \quad .$$

We need a formula for the derivative of the product of two functions. One might guess that the derivative of a product is the product of the derivatives; however, this is wrong.

### Example (Not the Product Rule)

Let  $f(x) = x$  and  $g(x) = x^2$ .

$$f'(x) = 1.$$

$$(fg)(x) = f(x)g(x) = x^3.$$

$$g'(x) = 2x.$$

$$(fg)'(x) = 3x^2.$$

$$f'(x)g'(x) = 2x.$$

Therefore

$$f'(x)g'(x) \neq (fg)'(x) \quad .$$

The correct formula is called the Product Rule.

## Theorem (The Product Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

Proof.

## Theorem (The Product Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$F'(x) =$$



## Theorem (The Product Rule)

*If  $f$  and  $g$  are both differentiable, then*

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$F'(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - \cancel{f(x+h)g(x)} + \cancel{f(x+h)g(x)} - f(x)g(x)}{h} \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\textcolor{red}{f(x+h)}g(x+h) - \textcolor{red}{f(x+h)}g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( \textcolor{red}{f(x+h)} \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)\color{red}{g(x+h)} - f(x+h)\color{red}{g(x)} + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{\color{red}{g(x+h)} - \color{red}{g(x)}}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \end{aligned}$$



## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \textcolor{red}{?} \quad ? \quad + \quad ? \quad ? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both **differentiable**, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)? \quad + ? \quad ? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x) \text{?} \quad + \text{?} \quad ? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both **differentiable**, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

### Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + ? \quad ? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

### Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + ? \quad ? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + g(x)? \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + g(x)? \end{aligned}$$



## Theorem (The Product Rule)

If  $f$  and  $g$  are both *differentiable*, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

### Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + g(x)f'(x). \end{aligned}$$

## Theorem (The Product Rule)

If  $f$  and  $g$  are both differentiable, then

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x).$$

## Proof.

Let  $F(x) = f(x)g(x)$ . Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \left( f(x+h) \frac{g(x+h) - g(x)}{h} + g(x) \frac{f(x+h) - f(x)}{h} \right) \\ &= \lim_{h \rightarrow 0} f(x+h) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &\quad + \lim_{h \rightarrow 0} g(x) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = f(x)g'(x) + g(x)f'(x). \quad \square \end{aligned}$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

Product Rule: 
$$f'(x) = \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x)$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

Product Rule: 
$$f'(x) = \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x)$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

$$\begin{aligned}\text{Product Rule: } f'(x) &= \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x) \\ &= (\text{?}) (e^x) + (x^3) (\text{?})\end{aligned}$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

$$\begin{aligned}\text{Product Rule: } f'(x) &= \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x) \\ &= \left( ? \right) (e^x) + (x^3) (e^x)\end{aligned}$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

$$\begin{aligned}\text{Product Rule: } f'(x) &= \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x) \\ &= (?) (e^x) + (x^3) (e^x)\end{aligned}$$



## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

$$\begin{aligned}\text{Product Rule: } f'(x) &= \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x) \\ &= (3x^2) (e^x) + (x^3) (e^x)\end{aligned}$$

## Example (Product Rule, polynomial times the Natural Exponential Function)

Differentiate  $f(x) = x^3 e^x$ .

$$\begin{aligned}\text{Product Rule: } f'(x) &= \frac{d}{dx} (x^3) (e^x) + (x^3) \frac{d}{dx} (e^x) \\ &= (3x^2) (e^x) + (x^3) (e^x) \\ &= e^x (x^3 + 3x^2) .\end{aligned}$$

## Theorem (The Quotient Rule)

*If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then*

$$\frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} \quad \left| \text{ (Leibniz notation) } \right.$$

## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\begin{array}{lcl} \frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) & = & \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} \quad \left( \text{Leibniz notation} \right) \\ \left( \frac{f(x)}{g(x)} \right)' & = & \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} \quad \left( \text{' notation} \right) \end{array}$$

## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\begin{aligned} \frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) &= \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} && \text{(Leibniz notation)} \\ \left( \frac{f(x)}{g(x)} \right)' &= \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2} && \text{' notation} \end{aligned}$$

## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\begin{aligned} \frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) &= \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} & (\text{Leibniz notation}) \\ \left( \frac{f(x)}{g(x)} \right)' &= \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} & ' \text{ notation} \end{aligned}$$

## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) = \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2}$$

*(Leibniz notation)*

$$\left( \frac{f(x)}{g(x)} \right)' = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

*' notation*

$$\left( \frac{f}{g} \right)' = \frac{f'g - fg'}{g^2}$$

*abbreviated*

## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\begin{array}{lcl} \frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) & = & \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} & \text{(Leibniz notation)} \\ \left( \frac{f(x)}{g(x)} \right)' & = & \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} & \text{' notation} \\ \left( \frac{f}{g} \right)' & = & \frac{f'g - fg'}{g^2} & \text{abbreviated} \end{array}$$

- The proof of the Quotient Rule is similar to the proof of the Product Rule.



## Theorem (The Quotient Rule)

If  $f$  and  $g$  are differentiable and  $g(x) \neq 0$ , then

$$\begin{array}{lcl} \frac{d}{dx} \left( \frac{f(x)}{g(x)} \right) & = & \frac{\frac{d}{dx} (f(x)) g(x) - f(x) \frac{d}{dx} (g(x))}{(g(x))^2} & \text{(Leibniz notation)} \\ \left( \frac{f(x)}{g(x)} \right)' & = & \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} & \text{' notation} \\ \left( \frac{f}{g} \right)' & = & \frac{f'g - fg'}{g^2} & \text{abbreviated} \end{array}$$

- The proof of the Quotient Rule is similar to the proof of the Product Rule.
- There is an alternative algebraic proof via the Product Rule, the Power Rule and the (not yet studied) Chain Rule.

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}.$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\frac{dy}{dx} = \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\frac{dy}{dx} = \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(\text{?})(-x^6 + 2) - (x^5 + 2x)(\text{?})}{(-x^6 + 2)^2}\end{aligned}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2)(-x^6 + 2) - (x^5 + 2x)(?) }{(-x^6 + 2)^2}\end{aligned}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned} \frac{dy}{dx} &= \frac{\frac{d}{dx} (x^5 + 2x) (-x^6 + 2) - (x^5 + 2x) \frac{d}{dx} (-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2) (-x^6 + 2) - (x^5 + 2x) (\text{?})}{(-x^6 + 2)^2} \end{aligned}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2)(-x^6 + 2) - (x^5 + 2x)(-6x^5)}{(-x^6 + 2)^2}\end{aligned}$$



## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\ &= \frac{(5x^4 + 2)(-x^6 + 2) - (x^5 + 2x)(-6x^5)}{(-x^6 + 2)^2} \\ &= \frac{(-5x^{10} - 2x^6 + 10x^4 + 4) - (-6x^{10} - 12x^6)}{(-x^6 + 2)^2}\end{aligned}$$

## Example (Quotient Rule, rational function)

Differentiate  $y = \frac{x^5 + 2x}{-x^6 + 2}$ .

Quotient Rule:

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{d}{dx}(x^5 + 2x)(-x^6 + 2) - (x^5 + 2x)\frac{d}{dx}(-x^6 + 2)}{(-x^6 + 2)^2} \\&= \frac{(5x^4 + 2)(-x^6 + 2) - (x^5 + 2x)(-6x^5)}{(-x^6 + 2)^2} \\&= \frac{(-5x^{10} - 2x^6 + 10x^4 + 4) - (-6x^{10} - 12x^6)}{(-x^6 + 2)^2} \\&= \frac{x^{10} + 10x^6 + 10x^4 + 4}{(-x^6 + 2)^2}.\end{aligned}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\frac{d}{dx} \left( \frac{1}{2x-1} \right)$$

Product rule

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\frac{d}{dx} \left( \frac{1}{2x-1} \right) = ?$$

Product rule

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\frac{d}{dx} \left( \frac{1}{2x-1} \right) = \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} \quad \left| \text{Product rule} \right.$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\frac{d}{dx} \left( \frac{1}{2x-1} \right) = \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} \quad \Bigg| \quad \text{Product rule}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\frac{d}{dx} \left( \frac{1}{2x-1} \right) = \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} \quad \Bigg| \quad \text{Product rule}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \text{Product rule} \right. \\ &= \frac{? \cdot (2x-1) - ?}{(2x-1)^2}\end{aligned}$$



$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \text{Product rule} \right. \\ &= \frac{0 \cdot (2x-1) - ?}{(2x-1)^2}\end{aligned}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \text{Product rule} \right. \\ &= \frac{0 \cdot (2x-1) - ?}{(2x-1)^2}\end{aligned}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \text{Product rule} \right. \\ &= \frac{0 \cdot (2x-1) - 2}{(2x-1)^2}\end{aligned}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \text{Product rule} \right. \\ &= \frac{0 \cdot (2x-1) - 2}{(2x-1)^2} \\ &= \frac{-2}{(2x-1)^2}\end{aligned}$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

## Example

Compute the derivative. Use the quotient rule.

$$\begin{aligned}\frac{d}{dx} \left( \frac{1}{2x-1} \right) &= \frac{(1)'(2x-1) - 1 \cdot (2x-1)'}{(2x-1)^2} && \left| \begin{array}{l} \text{Product rule} \end{array} \right. \\ &= \frac{0 \cdot (2x-1) - 2}{(2x-1)^2} \\ &= \frac{-2}{(2x-1)^2}\end{aligned}$$