

Calculus II

Trigonometric integrals

Todor Milev

2019

Outline

- 1 Trigonometric Integrals
 - Integrating rational trigonometric integrals
 - Ad hoc methods for trigonometric integrals

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - Apply the substitution $\theta = 2 \arctan t$ to transform to integral of rational function.

Integrals of the form $\int R(\cos \theta, \sin \theta) d\theta$, R

Let R be an arbitrary rational function in two variables (quotient of polynomials in two variables).

Question

Can we integrate $\int R(\cos \theta, \sin \theta) d\theta$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - Apply the substitution $\theta = 2 \arctan t$ to transform to integral of rational function.
 - Solve as previously studied.

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = ?$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z}{(\cos^2 z + \sin^2 z)}$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}}$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z)$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} .$$

$$\cos(2z) = ?$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad .$$

$$\cos(2z) = \cos^2 z - \sin^2 z$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad . \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z)}{(\cos^2 z + \sin^2 z)}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \quad . \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$.

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta, \cos \theta$?

$$\sin \theta =$$

$$\cos \theta =$$

Recall the expression of $\sin(2z), \cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t)$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)}$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} . \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} . \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2}$$

$$\cos \theta =$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2}$$

$$\cos \theta = \cos(2 \arctan t)$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned} \sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot \end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\sin \theta = \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2}$$

$$\cos \theta = \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)}$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\sin(2z) = 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z}.$$

$$\cos(2z) = \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z}.$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

Recall the expression of $\sin(2z)$, $\cos(2z)$ via $\tan z$:

$$\begin{aligned}\sin(2z) &= 2 \sin z \cos z = \frac{2 \sin z \cos z \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{2 \tan z}{1 + \tan^2 z} \cdot \\ \cos(2z) &= \cos^2 z - \sin^2 z = \frac{(\cos^2 z - \sin^2 z) \frac{1}{\cos^2 z}}{(\cos^2 z + \sin^2 z) \frac{1}{\cos^2 z}} = \frac{1 - \tan^2 z}{1 + \tan^2 z} \cdot\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta, \cos \theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2}\end{aligned}$$

$d\theta$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t)\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = ? \quad dt\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= ?\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= \tan\left(\frac{\theta}{2}\right)\end{aligned}$$

The rationalizing substitution $\theta = 2 \arctan t$

Let R - rational function in two variables. $\int R(\cos \theta, \sin \theta) d\theta$ can be integrated via the substitution $\theta = 2 \arctan t$. How does this transform $\sin \theta$, $\cos \theta$? How does this transform $d\theta$? How is t expressed via θ ?

$$\begin{aligned}\sin \theta &= \sin(2 \arctan t) = \frac{2 \tan(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{2t}{1 + t^2} \\ \cos \theta &= \cos(2 \arctan t) = \frac{1 - \tan^2(\arctan t)}{1 + \tan^2(\arctan t)} = \frac{1 - t^2}{1 + t^2} \\ d\theta &= 2d(\arctan t) = \frac{2}{1 + t^2} dt \\ t &= \tan\left(\frac{\theta}{2}\right)\end{aligned}$$

Theorem

The substitution given above transforms $\int R(\cos \theta, \sin \theta) d\theta$ to an integral of a rational function of t .

Example

$$\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} = \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} = \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} = \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + 4} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + 4} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + 4} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(\textcolor{red}{1} + t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + \textcolor{red}{5} \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + \textcolor{red}{4}} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + 4} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\ &= \int \frac{2dt}{6t^2 + 4t + 4} \\ &= \int \frac{dt}{3t^2 + 2t + 2} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{2}{3} \right)}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 \text{(complete square)} \quad &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{1}{9} - \frac{1}{9} + \frac{2}{3} \right)}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{1}{9} - \frac{1}{9} + \frac{2}{3} \right)} \\
 &= \frac{1}{3} \int \frac{dt}{\left(t + \frac{1}{3} \right)^2 + \frac{5}{9}}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{1}{9} - \frac{1}{9} + \frac{2}{3} \right)} \\
 &= \frac{1}{3} \int \frac{dt}{\left(t + \frac{1}{3} \right)^2 + \frac{5}{9}}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{1}{9} - \frac{1}{9} + \frac{2}{3} \right)} \\
 &= \frac{1}{3} \int \frac{dt}{\left(t + \frac{1}{3} \right)^2 + \frac{5}{9}} \\
 &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \int \frac{2dt}{(1+t^2) \left(2 \frac{2t}{t^2+1} - \frac{(1-t^2)}{1+t^2} + 5 \right)} \\
 &= \int \frac{2dt}{6t^2 + 4t + 4} \\
 &= \int \frac{dt}{3t^2 + 2t + 2} \\
 &= \int \frac{dt}{3 \left(t^2 + 2t \frac{1}{3} + \frac{1}{9} - \frac{1}{9} + \frac{2}{3} \right)} \\
 &= \frac{1}{3} \int \frac{dt}{\left(t + \frac{1}{3} \right)^2 + \frac{5}{9}} \\
 &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)}
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} = \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)} \\ &= \frac{3}{5} \int \frac{d \left(t + \frac{1}{3} \right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)^2 + 1 \right)} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)} \\ &= \frac{3}{5} \int \frac{d \left(t + \frac{1}{3} \right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)^2 + 1 \right)} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)} \\ &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d \left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)^2 + 1 \right)} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3} \right)^2 + 1 \right)} \\ &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d \left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right)^2 + 1 \right)} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$, $z = \frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)$.

$$\begin{aligned} \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3}\right)^2 + 1 \right)} \\ &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)^2 + 1\right)} \\ &= \frac{\sqrt{5}}{5} \int \frac{dz}{z^2 + 1} \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$, $z = \frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)$.

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3}\right)^2 + 1 \right)} \\
 &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)^2 + 1\right)} \\
 &= \frac{\sqrt{5}}{5} \int \frac{dz}{z^2 + 1} \\
 &= \frac{\sqrt{5}}{5} \arctan z + C
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$, $z = \frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)$.

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3}\right)^2 + 1 \right)} \\
 &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)^2 + 1\right)} \\
 &= \frac{\sqrt{5}}{5} \int \frac{dz}{z^2 + 1} \\
 &= \frac{\sqrt{5}}{5} \arctan z + C \\
 &= \frac{\sqrt{5}}{5} \arctan \left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right) \right) + C
 \end{aligned}$$

Example

Let $\theta = 2 \arctan t$, $\cos \theta = \frac{1-t^2}{1+t^2}$, $\sin \theta = \frac{2t}{1+t^2}$, $z = \frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)$.

$$\begin{aligned}
 \int \frac{d\theta}{2 \sin \theta - \cos \theta + 5} &= \frac{1}{3} \int \frac{dt}{\frac{5}{9} \left(\frac{9}{5} \left(t + \frac{1}{3}\right)^2 + 1 \right)} \\
 &= \frac{3}{5} \int \frac{\frac{\sqrt{5}}{3} d\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)}{\left(\left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3}\right)\right)^2 + 1\right)} \\
 &= \frac{\sqrt{5}}{5} \int \frac{dz}{z^2 + 1} \\
 &= \frac{\sqrt{5}}{5} \arctan z + C \\
 &= \frac{\sqrt{5}}{5} \arctan \left(\frac{3}{\sqrt{5}} \left(t + \frac{1}{3} \right) \right) + C \\
 &= \frac{\sqrt{5}}{5} \arctan \left(\frac{3}{\sqrt{5}} \left(\tan \left(\frac{\theta}{2} \right) + \frac{1}{3} \right) \right) + C
 \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\int \sec \theta d\theta$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$.

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \\ &= \ln \left| \frac{1+t}{1-t} \right| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\begin{aligned}
 \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\
 &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\
 &= -\ln |1-t| + \ln |1+t| + C \\
 &= \ln \left| \frac{1+t}{1-t} \right| + C \\
 &= \ln \left| \frac{1 + \tan\left(\frac{\theta}{2}\right)}{1 - \tan\left(\frac{\theta}{2}\right)} \right| + C
 \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\begin{aligned} \int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{1}{\left(\frac{1-t^2}{1+t^2}\right)} \frac{2}{(1+t^2)} dt \\ &= \int \frac{2}{1-t^2} dt = \int \left(\frac{1}{1-t} + \frac{1}{1+t} \right) dt \quad \left| \text{part. fractions} \right. \\ &= -\ln |1-t| + \ln |1+t| + C \\ &= \ln \left| \frac{1+t}{1-t} \right| + C \\ &= \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta =$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\tan \theta + \sec \theta = \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})) (\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2}))} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})) (\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2}))} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})) (\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2}))} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

Set $\theta = 2 \arctan t$, $\cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}$, $d\theta = 2 \frac{1}{1 + t^2} dt$.

$$\int \sec \theta d\theta = \ln \left| \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \right| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$

The integral $\int \sec \theta d\theta$ appears often in practice. A quicker solution will be shown later, but first we show the standard method.

Example

$$\text{Set } \theta = 2 \arctan t, \cos \theta = \frac{1 - \tan^2(\frac{\theta}{2})}{1 + \tan^2(\frac{\theta}{2})} = \frac{1 - t^2}{1 + t^2}, d\theta = 2 \frac{1}{1 + t^2} dt.$$

$$\int \sec \theta d\theta = \ln |\tan \theta + \sec \theta| + C$$

This is a perfectly good answer, however there's a simplification:

$$\begin{aligned} \tan \theta + \sec \theta &= \frac{\sin \theta + 1}{\cos \theta} = \frac{2 \sin(\frac{\theta}{2}) \cos(\frac{\theta}{2}) + \sin^2(\frac{\theta}{2}) + \cos^2(\frac{\theta}{2})}{\cos^2(\frac{\theta}{2}) - \sin^2(\frac{\theta}{2})} \\ &= \frac{(\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2}))^2}{(\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2}))(\cancel{\cos(\frac{\theta}{2}) + \sin(\frac{\theta}{2})})} \\ &= \frac{\sin(\frac{\theta}{2}) + \cos(\frac{\theta}{2})}{\cos(\frac{\theta}{2}) - \sin(\frac{\theta}{2})} = \frac{1 + \tan(\frac{\theta}{2})}{1 - \tan(\frac{\theta}{2})} \end{aligned}$$

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.
- This integration technique results in rather long computations.

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.
- This integration technique results in rather long computations.
- Particular integral types may be computable with quicker ad hoc techniques.

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.
- This integration technique results in rather long computations.
- Particular integral types may be computable with quicker ad hoc techniques.
- We illustrate such techniques on examples.

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.
- This integration technique results in rather long computations.
- Particular integral types may be computable with quicker ad hoc techniques.
- We illustrate such techniques on examples.
- Examples to which our ad hoc techniques apply arise from integrals needed outside of the subject of Calculus II, so these techniques are important.

Trigonometric Integrals - quick ad hoc techniques

- As we saw, every rational trigonometric expression can be integrated with the substitution $\theta = 2 \arctan t$.
- This integration technique results in rather long computations.
- Particular integral types may be computable with quicker ad hoc techniques.
- We illustrate such techniques on examples.
- Examples to which our ad hoc techniques apply arise from integrals needed outside of the subject of Calculus II, so these techniques are important.
- The trigonometric integral we saw, $\int \frac{d\theta}{2 \sin \theta - \cos \theta + 5}$, will not work with any of following ad-hoc techniques, so the general method is important as well.

Example

$$\int \sin^3 x dx$$

Example

$$\int \sin^3 x dx = \int \sin^2 x \sin x dx$$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\ &= \int \sin^2 x d(?) \end{aligned}$$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\ &= \int \sin^2 x d(-\cos x)\end{aligned}$$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\ &= \int \sin^2 x d(-\cos x) \\ &= \int (-1) \left(? \right) d(\cos x)\end{aligned}$$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\ &= \int \sin^2 x d(-\cos x) \\ &= \int (-1) \left(\text{?} \right) d(\cos x)\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\ &= \int \sin^2 x d(-\cos x) \\ &= \int (-1) (1 - \cos^2 x) d(\cos x)\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\&= \int \sin^2 x d(-\cos x) \\&= \int (-1) (1 - \cos^2 x) d(\cos x) \\&= \int (\cos^2 x - 1) d(\cos x)\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\&= \int \sin^2 x d(-\cos x) \\&= \int (-1) (1 - \cos^2 x) d(\cos x) \\&= \int (\cos^2 x - 1) d(\cos x) \\&= \int (u^2 - 1) du\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Set $u = \cos x$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\&= \int \sin^2 x d(-\cos x) \\&= \int (-1) (1 - \cos^2 x) d(\cos x) \\&= \int (\cos^2 x - 1) d(\cos x) \\&= \int (u^2 - 1) du \\&= \frac{u^3}{3} - u + C\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Set $u = \cos x$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\&= \int \sin^2 x d(-\cos x) \\&= \int (-1) (1 - \cos^2 x) d(\cos x) \\&= \int (\cos^2 x - 1) d(\cos x) \\&= \int (u^2 - 1) du \\&= \frac{u^3}{3} - u + C\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Set $u = \cos x$

Example

$$\begin{aligned}\int \sin^3 x dx &= \int \sin^2 x \sin x dx \\&= \int \sin^2 x d(-\cos x) \\&= \int (-1) (1 - \cos^2 x) d(\cos x) \\&= \int (\cos^2 x - 1) d(\cos x) \\&= \int (u^2 - 1) du \\&= \frac{u^3}{3} - u + C \\&= \frac{1}{3} \cos^3 x - \cos x + C .\end{aligned}$$

Can we rewrite
 $\sin^2 x$ via $\cos x$?

Set $u = \cos x$

Example

$$\int \cos^5 x \sin^2 x dx$$

Example

$$\int \cos^5 x \sin^2 x dx = \int \cos^4 x \sin^2 x \cos x dx$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\ &= \int \cos^4 x \sin^2 x d(?) \end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\ &= \int \cos^4 x \sin^2 x d(\sin x)\end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\ &= \int \cos^4 x \sin^2 x d(\sin x)\end{aligned}$$

Can we rewrite
 $\cos^4 x$ via $\sin x$?

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\ &= \int \cos^4 x \sin^2 x d(\sin x) \\ &= \int (\cos^2 x)^2 \sin^2 x d(\sin x)\end{aligned}$$

Can we rewrite
 $\cos^4 x$ via $\sin x$?

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x)\end{aligned}$$

Can we rewrite
 $\cos^4 x$ via $\sin x$?

Example

$$\int \cos^5 x \sin^2 x dx = \int \cos^4 x \sin^2 x \cos x dx$$

$$= \int \cos^4 x \sin^2 x d(\sin x)$$

Can we rewrite
 $\cos^4 x$ via $\sin x$?

$$= \int (\cos^2 x)^2 \sin^2 x d(\sin x)$$

$$= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x)$$

Set $u = \sin x$

$$= \int (1 - u^2)^2 u^2 du$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du\end{aligned}$$

Example

$$\int \cos^5 x \sin^2 x dx = \int \cos^4 x \sin^2 x \cos x dx$$

$$= \int \cos^4 x \sin^2 x d(\sin x)$$

Can we rewrite
 $\cos^4 x$ via $\sin x$?

$$= \int (\cos^2 x)^2 \sin^2 x d(\sin x)$$

$$= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) \quad \left| \text{Set } u = \sin x \right.$$

$$= \int (1 - u^2)^2 u^2 du$$

$$= \int (1 - 2u^2 + u^4) u^2 du$$

$$= \int (u^2 - 2u^4 + u^6) du$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du \\&= \int (u^2 - 2u^4 + u^6) du \\&= ?\end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du \\&= \int (u^2 - 2u^4 + u^6) du \\&= \frac{u^3}{3} - 2\frac{u^5}{5} + \frac{u^7}{7} + C\end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du \\&= \int (u^2 - 2u^4 + u^6) du \\&= \frac{u^3}{3} - 2\frac{u^5}{5} + \frac{u^7}{7} + C\end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du \\&= \int (u^2 - 2u^4 + u^6) du \\&= \frac{u^3}{3} - 2\frac{u^5}{5} + \frac{u^7}{7} + C\end{aligned}$$

Example

$$\begin{aligned}\int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\&= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\&= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\&= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\&= \int (1 - u^2)^2 u^2 du \\&= \int (1 - 2u^2 + u^4) u^2 du \\&= \int (u^2 - 2u^4 + u^6) du \\&= \frac{u^3}{3} - 2\frac{u^5}{5} + \frac{u^7}{7} + C\end{aligned}$$

Example

$$\begin{aligned}
 \int \cos^5 x \sin^2 x dx &= \int \cos^4 x \sin^2 x \cos x dx \\
 &= \int \cos^4 x \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \cos^4 x \text{ via } \sin x? \end{array} \right. \\
 &= \int (\cos^2 x)^2 \sin^2 x d(\sin x) \\
 &= \int (1 - \sin^2 x)^2 \sin^2 x d(\sin x) && \left| \begin{array}{l} \text{Set } u = \sin x \end{array} \right. \\
 &= \int (1 - u^2)^2 u^2 du \\
 &= \int (1 - 2u^2 + u^4) u^2 du \\
 &= \int (u^2 - 2u^4 + u^6) du \\
 &= \frac{u^3}{3} - 2\frac{u^5}{5} + \frac{u^7}{7} + C \\
 &= \frac{\sin^3 x}{3} - 2\frac{\sin^5 x}{5} + \frac{\sin^7 x}{7} + C .
 \end{aligned}$$

$$\int \sin^m x \cos^n x dx$$

When n – odd:

$$\int \sin^m x \cos^n x dx$$

When m – odd:

$$\int \sin^m x \cos^n x dx = \int \sin^m x \cos^{n-1} x d(\sin x)$$

When n – odd:
 $\cos x dx$
 $= d(\sin x)$

$$\int \sin^m x \cos^n x dx$$

When m – odd:

$$\begin{aligned}\int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\ &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x)\end{aligned}$$

When n – odd:

$\cos x dx$

$= d(\sin x)$

Express $\cos x$
via $\sin x$

$$\int \sin^m x \cos^n x dx$$

When m – odd:

$$\begin{aligned}\int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\ &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x)\end{aligned}$$

When n – odd:

$\cos x dx$

$= d(\sin x)$

Express $\cos x$
via $\sin x$

$$\int \sin^m x \cos^n x dx$$

When m – odd:

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:
 $\cos x dx$
 $= d(\sin x)$

Express $\cos x$
 via $\sin x$

Set $\sin x = u$

$$\int \sin^m x \cos^n x dx$$

When m – odd:

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:
 $\cos x dx$
 $= d(\sin x)$

Express $\cos x$
 via $\sin x$

Set $\sin x = u$

$$\int \sin^m x \cos^n x dx = \int \sin^{m-1} x \cos^n x d(-\cos x)$$

When m – odd:
 $\sin x dx$
 $= d(-\cos x)$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

Set $\sin x = u$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x)
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\cos x$
via $\sin x$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

Set $\sin x = u$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x)
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\cos x$
via $\sin x$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \sin x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x)
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\cos x$
via $\sin x$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \sin x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x) \\
 &= - \int (1 - u^2)^{\frac{m-1}{2}} u^n du
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \cos x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \sin x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x) \\
 &= - \int (1 - u^2)^{\frac{m-1}{2}} u^n du
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \cos x = u$$

If both m, n – even,

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \sin x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x) \\
 &= - \int (1 - u^2)^{\frac{m-1}{2}} u^n du
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\sin x$
via $\cos x$

$$\text{Set } \cos x = u$$

If both m, n – even, use $\left| \begin{array}{l} \sin^2 x = \frac{1 - \cos(2x)}{2} \\ \cos^2 x = \frac{\cos(2x) + 1}{2} \end{array} \right.$ and substitute $s = 2x$ to lower trig powers. Repeat above considerations.

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^m x \cos^{n-1} x d(\sin x) \\
 &= \int \sin^m x (1 - \sin^2 x)^{\frac{n-1}{2}} d(\sin x) \\
 &= \int u^m (1 - u^2)^{\frac{n-1}{2}} du
 \end{aligned}$$

When n – odd:

$$\cos x dx$$

$$= d(\sin x)$$

Express $\cos x$
via $\sin x$

$$\text{Set } \sin x = u$$

$$\begin{aligned}
 \int \sin^m x \cos^n x dx &= \int \sin^{m-1} x \cos^n x d(-\cos x) \\
 &= - \int (1 - \cos^2 x)^{\frac{m-1}{2}} \cos^n x d(\cos x) \\
 &= - \int (1 - u^2)^{\frac{m-1}{2}} u^n du
 \end{aligned}$$

When m – odd:

$$\sin x dx$$

$$= d(-\cos x)$$

Express $\sin x$
via $\cos x$

$$\text{Set } \cos x = u$$

If both m, n – even, use $\left| \begin{array}{l} \sin^2 x = \frac{1 - \cos(2x)}{2} \\ \cos^2 x = \frac{\cos(2x) + 1}{2} \end{array} \right.$ and **substitute $s = 2x$** to lower trig powers. Repeat above considerations.

Example

$$\int_0^{\frac{\pi}{2}} \sin^2 x dx$$

Example

Example

$$\int_0^{\frac{\pi}{2}} \sin^2 x dx = \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx$$

express $\sin^2 x$
via $\cos(2x)$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}}\end{aligned}$$

express $\sin^2 x$
via $\cos(2x)$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}}\end{aligned}$$

express $\sin^2 x$
via $\cos(2x)$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}}\end{aligned}$$

express $\sin^2 x$
via $\cos(2x)$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\ &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) .\end{aligned}$$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\ &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) .\end{aligned}$$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\ &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = ? .\end{aligned}$$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\ &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.\end{aligned}$$

Example

Example

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\ &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\ &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.\end{aligned}$$


Example

$$\int_{t=0}^{t=1} \sqrt{1-t^2} \, dt$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example




$$\int_{t=0}^{t=1} \sqrt{1 - t^2} \, dt$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$.




$$\int_{t=0}^{t=1} \sqrt{1 - t^2} \, dt$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then
 $dt = d(\cos x) = ?$.



$$\int_{t=0}^{t=1} \sqrt{1 - t^2} \, dt$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$


Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$

. Then

$$dt = d(\cos x) = -\sin x \, dx.$$

$$\int_{t=0}^{t=1} \sqrt{1-t^2} \, dt$$



Example


$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then

$$dt = d(\cos x) = -\sin x \, dx.$$

$$\int_{t=0}^{t=1} \sqrt{1-t^2} \, dt = - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx$$



Example


$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then

$$dt = d(\cos x) = -\sin x \, dx.$$

$$\int_{t=0}^{t=1} \sqrt{1-t^2} \, dt = - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx$$




Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then
 $dt = d(\cos x) = -\sin x \, dx$.

$$\int_{t=0}^{t=1} \sqrt{1-t^2} \, dt = - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx$$




Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then
 $dt = d(\cos x) = -\sin x \, dx$.

$$\int_{t=1}^0 \sqrt{1-t^2} \, dt = - \int_{x=\frac{\pi}{2}}^0 \sqrt{1-\cos^2 x} \sin x \, dx$$




Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then
 $dt = d(\cos x) = -\sin x \, dx$.




$$\begin{aligned}
 \int_{t=0}^{t=1} \sqrt{1-t^2} \, dt &= - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx \\
 &= \int_{x=0}^{x=\frac{\pi}{2}} \sqrt{\sin^2 x} \sin x \, dx
 \end{aligned}$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}]$. Then
 $dt = d(\cos x) = -\sin x \, dx$.




$$\begin{aligned}
 \int_{t=0}^{t=1} \sqrt{1-t^2} \, dt &= - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx \\
 &= \int_{x=0}^{x=\frac{\pi}{2}} \sqrt{\sin^2 x} \sin x \, dx
 \end{aligned}$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}] \Rightarrow \sin x \geq 0$. Then
 $dt = d(\cos x) = -\sin x \, dx$.




$$\begin{aligned}
 \int_{t=0}^{t=1} \sqrt{1-t^2} \, dt &= - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx \\
 &= \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{\sin^2 x} \sin x \, dx \\
 &= \int_0^{\frac{\pi}{2}} \sin^2 x \, dx
 \end{aligned}$$

Example

$$\begin{aligned}
 \int_0^{\frac{\pi}{2}} \sin^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1 - \cos(2x)}{2} \right) dx && \left| \begin{array}{l} \text{express } \sin^2 x \\ \text{via } \cos(2x) \end{array} \right. \\
 &= \left[\frac{x}{2} - \frac{\sin(2x)}{4} \right]_0^{\frac{\pi}{2}} \\
 &= \left(\frac{\pi}{4} - \frac{\sin \pi}{4} \right) - \left(0 - \frac{\sin 0}{4} \right) = \frac{\pi}{4}.
 \end{aligned}$$

Example

Set $t = \cos x$, $x \in [0, \frac{\pi}{2}] \Rightarrow \sin x \geq 0$. Then
 $dt = d(\cos x) = -\sin x \, dx$.



$$\begin{aligned}
 \int_{t=0}^{t=1} \sqrt{1-t^2} \, dt &= - \int_{x=\frac{\pi}{2}}^{x=0} \sqrt{1-\cos^2 x} \sin x \, dx \\
 &= \int_{x=0}^{x=\frac{\pi}{2}} \sqrt{\sin^2 x} \sin x \, dx \\
 &= \int_0^{\frac{\pi}{2}} \sin^2 x \, dx = \frac{\pi}{4}.
 \end{aligned}$$

Example

$$\int \tan^8 x \sec^4 x dx$$

Example

$$\int \tan^8 x \sec^4 x dx = \int \tan^8 x \sec^2 x \sec^2 x dx$$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\ &= \int \tan^8 x \sec^2 x d(?)\end{aligned}$$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\ &= \int \tan^8 x \sec^2 x d(\tan x)\end{aligned}$$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\ &= \int \tan^8 x \sec^2 x d(\tan x) \\ &= \int \tan^8 x \left(? \right) d(\tan x)\end{aligned}$$

Can we rewrite $\sec^2 x$ via $\tan x$?

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\ &= \int \tan^8 x \sec^2 x d(\tan x) \\ &= \int \tan^8 x (1 + \tan^2 x) d(\tan x)\end{aligned}$$

Can we rewrite
 $\sec^2 x$ via $\tan x$?

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\ &= \int \tan^8 x \sec^2 x d(\tan x) \\ &= \int \tan^8 x (1 + \tan^2 x) d(\tan x) \\ &= \int u^8 (1 + u^2) du\end{aligned}$$

Can we rewrite $\sec^2 x$ via $\tan x$?
Set $u = \tan x$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\&= \int \tan^8 x \sec^2 x d(\tan x) \\&= \int \tan^8 x (1 + \tan^2 x) d(\tan x) \\&= \int u^8 (1 + u^2) du \\&= \int (u^8 + u^{10}) du\end{aligned}$$

Can we rewrite $\sec^2 x$ via $\tan x$?
Set $u = \tan x$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\&= \int \tan^8 x \sec^2 x d(\tan x) \\&= \int \tan^8 x (1 + \tan^2 x) d(\tan x) \\&= \int u^8 (1 + u^2) du \\&= \int (u^8 + u^{10}) du \\&= ?\end{aligned}$$

Can we rewrite $\sec^2 x$ via $\tan x$?
Set $u = \tan x$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\&= \int \tan^8 x \sec^2 x d(\tan x) && \left. \begin{array}{l} \text{Can we rewrite} \\ \sec^2 x \text{ via } \tan x? \end{array} \right| \\&= \int \tan^8 x (1 + \tan^2 x) d(\tan x) && \left| \text{Set } u = \tan x \right. \\&= \int u^8 (1 + u^2) du \\&= \int (u^8 + u^{10}) du \\&= \frac{u^9}{9} + \frac{u^{11}}{11} + C\end{aligned}$$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\&= \int \tan^8 x \sec^2 x d(\tan x) && \left. \begin{array}{l} \text{Can we rewrite} \\ \sec^2 x \text{ via } \tan x? \end{array} \right| \\&= \int \tan^8 x (1 + \tan^2 x) d(\tan x) && \left| \text{Set } u = \tan x \right. \\&= \int u^8 (1 + u^2) du \\&= \int (u^8 + u^{10}) du \\&= \frac{u^9}{9} + \frac{u^{11}}{11} + C\end{aligned}$$

Example

$$\begin{aligned}\int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\&= \int \tan^8 x \sec^2 x d(\tan x) \\&= \int \tan^8 x (1 + \tan^2 x) d(\tan x) \\&= \int u^8 (1 + u^2) du \\&= \int (u^8 + u^{10}) du \\&= \frac{u^9}{9} + \frac{u^{11}}{11} + C\end{aligned}$$

Can we rewrite $\sec^2 x$ via $\tan x$?
Set $u = \tan x$

Example

$$\begin{aligned}
 \int \tan^8 x \sec^4 x dx &= \int \tan^8 x \sec^2 x \sec^2 x dx \\
 &= \int \tan^8 x \sec^2 x d(\tan x) && \left. \begin{array}{l} \text{Can we rewrite} \\ \sec^2 x \text{ via } \tan x? \end{array} \right| \\
 &= \int \tan^8 x (1 + \tan^2 x) d(\tan x) && \text{Set } u = \tan x \\
 &= \int u^8 (1 + u^2) du \\
 &= \int (u^8 + u^{10}) du \\
 &= \frac{u^9}{9} + \frac{u^{11}}{11} + C \\
 &= \frac{\tan^9 x}{9} + \frac{\tan^{11} x}{11} + C .
 \end{aligned}$$

Example

$$\int \tan^5 x \sec^9 x dx$$

Example

$$\int \tan^5 x \sec^9 x dx = \int \tan^4 x \sec^8 x \tan x \sec x dx$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\ &= \int \tan^4 x \sec^8 x d(?) \end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\ &= \int \tan^4 x \sec^8 x d(\sec x)\end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\ &= \int \tan^4 x \sec^8 x d(\sec x)\end{aligned}$$

Can we rewrite
 $\tan^4 x$ via $\sec x$?

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\ &= \int \tan^4 x \sec^8 x d(\sec x) \\ &= \int (\tan^2 x)^2 \sec^8 x d(\sec x)\end{aligned}$$

Can we rewrite
 $\tan^4 x$ via $\sec x$?

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\ &= \int \tan^4 x \sec^8 x d(\sec x) \\ &= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\ &= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x)\end{aligned}$$

Can we rewrite
 $\tan^4 x$ via $\sec x$?

Example

$$\begin{aligned}
 \int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\
 &= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\
 &= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\
 &= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Set } u = \sec x \end{array} \right. \\
 &= \int (1 - u^2)^2 u^8 du
 \end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\&= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\&= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\&= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \text{Set } u = \sec x \right. \\&= \int (1 - u^2)^2 u^8 du \\&= \int (1 - 2u^2 + u^4) u^8 du\end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\&= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\&= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\&= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Set } u = \sec x \end{array} \right. \\&= \int (1 - u^2)^2 u^8 du \\&= \int (1 - 2u^2 + u^4) u^8 du \\&= \int (u^8 - 2u^{10} + u^{12}) du\end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\&= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\&= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\&= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \text{Set } u = \sec x \right. \\&= \int (1 - u^2)^2 u^8 du \\&= \int (1 - 2u^2 + u^4) u^8 du \\&= \int (u^8 - 2u^{10} + u^{12}) du \\&= ?\end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\&= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\&= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\&= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \text{Set } u = \sec x \right. \\&= \int (1 - u^2)^2 u^8 du \\&= \int (1 - 2u^2 + u^4) u^8 du \\&= \int (u^8 - 2u^{10} + u^{12}) du \\&= \frac{u^9}{9} - 2\frac{u^{11}}{11} + \frac{u^{13}}{13} + C\end{aligned}$$

Example

$$\begin{aligned}\int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\&= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\&= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\&= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Set } u = \sec x \end{array} \right. \\&= \int (1 - u^2)^2 u^8 du \\&= \int (1 - 2u^2 + u^4) u^8 du \\&= \int (u^8 - 2u^{10} + u^{12}) du \\&= \frac{u^9}{9} - 2\frac{u^{11}}{11} + \frac{u^{13}}{13} + C\end{aligned}$$

Example

$$\begin{aligned}
 \int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\
 &= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\
 &= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\
 &= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \text{Set } u = \sec x \right. \\
 &= \int (1 - u^2)^2 u^8 du \\
 &= \int (1 - 2u^2 + u^4) u^8 du \\
 &= \int (u^8 - 2u^{10} + u^{12}) du \\
 &= \frac{u^9}{9} - 2\frac{u^{11}}{11} + \frac{u^{13}}{13} + C
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\
 &= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\
 &= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\
 &= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \text{Set } u = \sec x \right. \\
 &= \int (1 - u^2)^2 u^8 du \\
 &= \int (1 - 2u^2 + u^4) u^8 du \\
 &= \int (u^8 - 2u^{10} + u^{12}) du \\
 &= \frac{u^9}{9} - 2\frac{u^{11}}{11} + \frac{u^{13}}{13} + C
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan^5 x \sec^9 x dx &= \int \tan^4 x \sec^8 x \tan x \sec x dx \\
 &= \int \tan^4 x \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Can we rewrite} \\ \tan^4 x \text{ via } \sec x? \end{array} \right. \\
 &= \int (\tan^2 x)^2 \sec^8 x d(\sec x) \\
 &= \int (\sec^2 x - 1)^2 \sec^8 x d(\sec x) && \left| \begin{array}{l} \text{Set } u = \sec x \end{array} \right. \\
 &= \int (1 - u^2)^2 u^8 du \\
 &= \int (1 - 2u^2 + u^4) u^8 du \\
 &= \int (u^8 - 2u^{10} + u^{12}) du \\
 &= \frac{u^9}{9} - 2\frac{u^{11}}{11} + \frac{u^{13}}{13} + C \\
 &= \frac{\sec^9 x}{9} - 2\frac{\sec^{11} x}{11} + \frac{\sec^{13} x}{13} + C .
 \end{aligned}$$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$$\int \tan^m x \sec^n x dx \quad \left| \begin{array}{l} n - \text{even}, n \geq 2 \end{array} \right.$$

$$\int \tan^m x \sec^n x dx \quad \left| \begin{array}{l} m - \text{odd}, n \geq 1 \end{array} \right.$$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$$\int \tan^m x \sec^n x dx = \int \tan^m x \sec^{n-2} x d(\tan x) \quad \left| \begin{array}{l} n - \text{even}, n \geq 2 \\ \sec^2 x dx \\ = d(\tan x) \end{array} \right.$$

$\int \tan^m x \sec^n x dx$	$m - \text{odd}, n \geq 1$
-----------------------------	----------------------------

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$$\begin{aligned} \int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \end{aligned} \quad \left| \begin{array}{l} n - \text{even}, n \geq 2 \\ \sec^2 x dx \\ = d(\tan x) \\ \text{Express } \sec x \\ \text{via } \tan x \end{array} \right.$$

$$\int \tan^m x \sec^n x dx \quad \left| \begin{array}{l} m - \text{odd}, n \geq 1 \end{array} \right.$$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$$\begin{aligned} \int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \end{aligned}$$

$n - \text{even}, n \geq 2$
 $\sec^2 x dx = d(\tan x)$
 Express $\sec x$
 via $\tan x$

$$\int \tan^m x \sec^n x dx$$

$m - \text{odd}, n \geq 1$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned} \int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du \end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx = d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
<hr style="border: 0; border-top: 1px solid black; margin: 10px 0;"/> $\int \tan^m x \sec^n x dx$	$m - \text{odd}, n \geq 1$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx = d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
<hr/> $\int \tan^m x \sec^n x dx = \int \tan^{m-1} x \sec^{n-1} x d(\sec x)$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx = d(\sec x)$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$$\begin{aligned}
 \int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\
 &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\
 &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du
 \end{aligned}$$

$n - \text{even}, n \geq 2$

$\sec^2 x dx$

$= d(\tan x)$

Express $\sec x$
via $\tan x$

Set $u = \tan x$

$$\begin{aligned}
 \int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\
 &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x)
 \end{aligned}$$

$m - \text{odd}, n \geq 1$

$\tan x \sec x dx$

$= d(\sec x)$

Express $\tan x$
via $\sec x$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx$ $= d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\ &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x)\end{aligned}$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx$ $= d(\sec x)$ Express $\tan x$ via $\sec x$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx$ $= d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\ &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x)\end{aligned}$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx$ $= d(\sec x)$ Express $\tan x$ via $\sec x$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx$ $= d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\ &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x) \\ &= \int (u^2 - 1)^{\frac{m-1}{2}} u^n du\end{aligned}$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx$ $= d(\sec x)$ Express $\tan x$ via $\sec x$ Set $u = \sec x$

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx = d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
--	---

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\ &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x) \\ &= \int (u^2 - 1)^{\frac{m-1}{2}} u^n du\end{aligned}$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx = d(\sec x)$ Express $\tan x$ via $\sec x$ Set $u = \sec x$
--	---

Outside of the above cases we either use more tricks or resort to the general method $x = 2 \arctan t$.

Partial strategy for fast evaluation of $\int \tan^m x \sec^n x dx$

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^m x \sec^{n-2} x d(\tan x) \\ &= \int \tan^m x (1 + \tan^2 x)^{\frac{n-2}{2}} d(\tan x) \\ &= \int u^m (1 + u^2)^{\frac{n-2}{2}} du\end{aligned}$	$n - \text{even}, n \geq 2$ $\sec^2 x dx$ $= d(\tan x)$ Express $\sec x$ via $\tan x$ Set $u = \tan x$
--	---

$\begin{aligned}\int \tan^m x \sec^n x dx &= \int \tan^{m-1} x \sec^{n-1} x d(\sec x) \\ &= \int (\sec^2 x - 1)^{\frac{m-1}{2}} \sec^{n-1} x d(\sec x) \\ &= \int (u^2 - 1)^{\frac{m-1}{2}} u^n du\end{aligned}$	$m - \text{odd}, n \geq 1$ $\tan x \sec x dx$ $= d(\sec x)$ Express $\tan x$ via $\sec x$ Set $u = \sec x$
--	---

Outside of the above cases we either use **more tricks** or resort to **the general method** $x = 2 \arctan t$.

Example

$$\int \tan x dx$$

Example

Example

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx$$

Example

Example

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(?)$$

Example

Example

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x)$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\ &= - \int \frac{du}{u}\end{aligned}$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\ &= - \int \frac{du}{u}\end{aligned}$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \left| \text{Set } u = \cos x \right. \\ &= - \int \frac{du}{u} = -\ln |u| + C\end{aligned}$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) && \left| \text{Set } u = \cos x \right. \\ &= - \int \frac{du}{u} = -\ln |u| + C \\ &= -\ln |\cos x| + C\end{aligned}$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) && \left| \text{Set } u = \cos x \right. \\ &= - \int \frac{du}{u} = -\ln |u| + C \\ &= -\ln |\text{red } x| + C = \ln |\text{red } x| + C\end{aligned}$$

Example

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\ &= - \int \frac{du}{u} = -\ln |u| + C \\ &= -\ln |\cos x| + C = \ln |\sec x| + C\end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$.

Example

$$\int \sec x dx$$

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\ &= - \int \frac{du}{u} = -\ln |u| + C \\ &= -\ln |\cos x| + C = \ln |\sec x| + C\end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\int \sec x dx = \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx$$

Example

$$\begin{aligned}\int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\ &= - \int \frac{du}{u} = -\ln |u| + C \\ &= -\ln |\cos x| + C = \ln |\sec x| + C\end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}\int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\ &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx\end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \left| \text{Set } u = \cos x \right. \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \text{ Set } u = \cos x \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\
 &= \int \frac{d(\tan x + \sec x)}{\sec x + \tan x}
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) \quad \Bigg| \quad \text{Set } u = \cos x \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \textcolor{red}{\sec x \tan x}}{\sec x + \tan x} dx \\
 &= \int \frac{\textcolor{red}{d}(\tan x + \textcolor{red}{\sec x})}{\sec x + \tan x}
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) && \left| \text{Set } u = \cos x \right. \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\
 &= \int \frac{d(\tan x + \sec x)}{\sec x + \tan x} && \left| \text{Set } u = \sec x + \tan x \right. \\
 &= \int \frac{du}{u}
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) && \left| \text{Set } u = \cos x \right. \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\
 &= \int \frac{d(\tan x + \sec x)}{\sec x + \tan x} && \left| \text{Set } u = \sec x + \tan x \right. \\
 &= \int \frac{du}{u} = \ln |u| + C
 \end{aligned}$$

Example

$$\begin{aligned}
 \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = \int \frac{1}{\cos x} d(-\cos x) && \left| \text{Set } u = \cos x \right. \\
 &= - \int \frac{du}{u} = -\ln |u| + C \\
 &= -\ln |\cos x| + C = \ln |\sec x| + C
 \end{aligned}$$

The following can be/was computed via $x = 2 \arctan t$. Alternatively:

Example

$$\begin{aligned}
 \int \sec x dx &= \int \sec x \frac{(\sec x + \tan x)}{(\sec x + \tan x)} dx \\
 &= \int \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x} dx \\
 &= \int \frac{d(\tan x + \sec x)}{\sec x + \tan x} && \left| \text{Set } u = \sec x + \tan x \right. \\
 &= \int \frac{du}{u} = \ln |u| + C \\
 &= \ln |\sec x + \tan x| + C.
 \end{aligned}$$

Example

$$\int \tan^3 x dx$$

Example

$$\int \tan^3 x dx = \int \tan x \tan^2 x dx$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\ &= \int \tan x (\sec^2 x - 1) dx\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\ &= \int \tan x (\sec^2 x - 1) dx \\ &= \int \tan x \sec^2 x dx - \int \tan x dx\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\ &= \int \tan x (\sec^2 x - 1) dx \\ &= \int \tan x \sec^2 x dx - \int \tan x dx\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(?) - ?\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - ?\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - ?\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x|\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x| \quad \left| \text{Set } u = \tan x \right. \\&= \int u du + \ln \left| \frac{1}{\sec x} \right|\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x| \quad \left| \text{Set } u = \tan x \right. \\&= \int u du + \ln \left| \frac{1}{\sec x} \right|\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x| \quad \left| \text{Set } u = \tan x \right. \\&= \int u du + \ln \left| \frac{1}{\sec x} \right| \\&= \frac{u^2}{2} + \ln |\cos x| + C\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x| \quad \left| \text{Set } u = \tan x \right. \\&= \int u du + \ln \left| \frac{1}{\sec x} \right| \\&= \frac{u^2}{2} + \ln |\cos x| + C\end{aligned}$$

Example

$$\begin{aligned}\int \tan^3 x dx &= \int \tan x \tan^2 x dx \\&= \int \tan x (\sec^2 x - 1) dx \\&= \int \tan x \sec^2 x dx - \int \tan x dx \\&= \int \tan x d(\tan x) - \ln |\sec x| \quad \left| \text{Set } u = \tan x \right. \\&= \int u du + \ln \left| \frac{1}{\sec x} \right| \\&= \frac{u^2}{2} + \ln |\cos x| + C \\&= \frac{\tan^2 x}{2} + \ln |\cos x| + C\end{aligned}$$

Example

$$\int \sec^3 x dx$$

Example

$$\int \sec^3 x dx = \int \sec x \sec^2 x dx$$

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\ &= \int \sec x d(?)\end{aligned}$$

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\ &= \int \sec x d(\tan x)\end{aligned}$$

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\ &= \int \sec x d(\tan x) \\ &= \sec x \tan x - \int \tan x d(\sec x)\end{aligned}$$

Integrate
by parts

Example

$$\int \sec^3 x dx = \int \sec x \sec^2 x dx$$

$$= \int \sec x d(\tan x)$$

$$= \sec x \tan x - \int \tan x d(\sec x)$$

Integrate
by parts

Example

$$\int \sec^3 x dx = \int \sec x \sec^2 x dx$$

$$= \int \sec x d(\tan x)$$

$$= \sec x \tan x - \int \tan x d(\sec x)$$

$$= \sec x \tan x - \int \tan x \quad dx$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan x \sec x \tan x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan x \sec x \tan x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\ &= \int \sec x d(\tan x) \\ &= \sec x \tan x - \int \tan x d(\sec x) \\ &= \sec x \tan x - \int \tan^2 x \sec x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan^2 x \sec x dx \\&= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan^2 x \sec x dx \\&= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\&= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan^2 x \sec x dx \\&= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\&= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan^2 x \sec x dx \\&= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\&= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx\end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}
 \int \sec^3 x dx &= \int \sec x \sec^2 x dx \\
 &= \int \sec x d(\tan x) \\
 &= \sec x \tan x - \int \tan x d(\sec x) \\
 &= \sec x \tan x - \int \tan^2 x \sec x dx \\
 &= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\
 &= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx \\
 2 \int \sec^3 x dx &= \sec x \tan x + ? \quad + C
 \end{aligned}$$

Integrate
by parts

Example

$$\begin{aligned}\int \sec^3 x dx &= \int \sec x \sec^2 x dx \\&= \int \sec x d(\tan x) && \left| \begin{array}{l} \text{Integrate} \\ \text{by parts} \end{array} \right. \\&= \sec x \tan x - \int \tan x d(\sec x) \\&= \sec x \tan x - \int \tan^2 x \sec x dx \\&= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\&= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx \\2 \int \sec^3 x dx &= \sec x \tan x + \textcolor{red}{?} + C\end{aligned}$$

Example

$$\begin{aligned}
 \int \sec^3 x dx &= \int \sec x \sec^2 x dx \\
 &= \int \sec x d(\tan x) \\
 &= \sec x \tan x - \int \tan x d(\sec x) \\
 &= \sec x \tan x - \int \tan^2 x \sec x dx \\
 &= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx \\
 &= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx \\
 2 \int \sec^3 x dx &= \sec x \tan x + \ln |\sec x + \tan x| + C
 \end{aligned}$$

Integrate
by parts

Example

$$\int \sec^3 x dx = \int \sec x \sec^2 x dx$$

$$= \int \sec x d(\tan x)$$

Integrate
by parts

$$= \sec x \tan x - \int \tan x d(\sec x)$$

$$= \sec x \tan x - \int \tan^2 x \sec x dx$$

$$= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx$$

$$= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx$$

$$2 \int \sec^3 x dx = \sec x \tan x + \ln |\sec x + \tan x| + C$$

$$\int \sec^3 x dx = \frac{1}{2} (\sec x \tan x + \ln |\sec x + \tan x|) + K.$$

To evaluate integrals of the form

$$\textcircled{1} \int \sin(mx) \cos(nx) dx$$

$$\textcircled{2} \int \sin(mx) \sin(nx) dx$$

$$\textcircled{3} \int \cos(mx) \cos(nx) dx$$

use the corresponding identity:

$$\textcircled{1} \sin A \cos B = \frac{1}{2} [\sin(A - B) + \sin(A + B)]$$

$$\textcircled{2} \sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)]$$

$$\textcircled{3} \cos A \cos B = \frac{1}{2} [\cos(A - B) + \cos(A + B)]$$

Example

$$\int \sin(4x) \cos(5x) dx$$

Example

$$\int \sin(4x) \cos(5x) dx = \int \frac{1}{2} [\sin(4x - 5x) + \sin(4x + 5x)] dx$$

Example

$$\begin{aligned}\int \sin(4x) \cos(5x) dx &= \int \frac{1}{2} [\sin(4x - 5x) + \sin(4x + 5x)] dx \\ &= \frac{1}{2} \int (\sin(-x) + \sin(9x)) dx\end{aligned}$$

Example

$$\begin{aligned}\int \sin(4x) \cos(5x) dx &= \int \frac{1}{2} [\sin(4x - 5x) + \sin(4x + 5x)] dx \\ &= \frac{1}{2} \int (\sin(-x) + \sin(9x)) dx \\ &= \frac{1}{2} \int (-\sin x + \sin(9x)) dx\end{aligned}$$

Example

$$\begin{aligned}\int \sin(4x) \cos(5x) dx &= \int \frac{1}{2} [\sin(4x - 5x) + \sin(4x + 5x)] dx \\ &= \frac{1}{2} \int (\sin(-x) + \sin(9x)) dx \\ &= \frac{1}{2} \int (-\sin x + \sin(9x)) dx \\ &= \frac{1}{2} \left(\cos x - \frac{1}{9} \cos(9x) \right) + C\end{aligned}$$