

Calculus II


Definite integrals of the form $\int_p^q \sqrt{ax^2 + c} dx,$
 $a, c > 0$

Todor Milev

2019


Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Example


Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



why do we call
 $y = \sqrt{x^2 + 1}$ hyperbola?

Example


Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola?

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2\end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned}\sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1\end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$(y - x)(y + x) = 1$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{1}{2} (y - x) (y + x) = \frac{1}{2}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$


$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) = \frac{1}{2}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$.

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$.

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}.$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned}
 \sqrt{x^2 + 1} &= y \\
 x^2 + 1 &= y^2 \\
 y^2 - x^2 &= 1 \\
 \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\
 uv &= \frac{1}{2} \\
 v &= \frac{1}{2u},
 \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



distance b-n (x, y) and line $u = 0$ equals


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



distance b-n (x, y) and line

$u = 0$ equals

$$\sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2}$$

We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\sqrt{x^2 + 1} = y$$

$$x^2 + 1 = y^2$$

$$y^2 - x^2 = 1$$

$$\frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) = \frac{1}{2}$$


$$uv = \frac{1}{2}$$

$$v = \frac{1}{2u}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



distance b-n (x, y) and line

$u = 0$ equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} \end{aligned}$$


We studied $v = \frac{1}{2}u$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



distance b-n (x, y) and line

$u = 0$ equals

$$\begin{aligned} & \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \sqrt{\frac{1}{2}(y-x)^2} = \pm \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) \end{aligned}$$


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals

$$\begin{aligned} & \pm \sqrt{\left(x - \frac{(x+y)}{2}\right)^2 + \left(y - \frac{(x+y)}{2}\right)^2} \\ &= \pm \sqrt{\frac{1}{2}(y-x)^2} = \frac{\sqrt{2}}{2}(y-x) = u. \end{aligned}$$


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y-x) \frac{\sqrt{2}}{2}(y+x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y-x) \\ v = \frac{\sqrt{2}}{2}(y+x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals u .


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals u . Similarly compute that signed distance b-n (x, y) and the line $v = 0$ equals v .


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals u . Similarly compute that **signed distance b-n (x, y) and the line $v = 0$ equals v .**


We studied $v = \frac{1}{u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



Signed distance b-n (x, y) and line $u = 0$ equals u . Similarly compute that signed distance b-n (x, y) and the line $v = 0$ equals v .
 $\Rightarrow y^2 - x^2 = 1$ is the **hyperbola**
 $v = \frac{1/2}{u}$ in the (u, v) -plane.


We studied $v = \frac{1}{2u}$ is called a hyperbola: why do we call $y = \sqrt{x^2 + 1}$ hyperbola? Compute:

$$\begin{aligned} \sqrt{x^2 + 1} &= y \\ x^2 + 1 &= y^2 \\ y^2 - x^2 &= 1 \\ \frac{\sqrt{2}}{2}(y - x) \frac{\sqrt{2}}{2}(y + x) &= \frac{1}{2} \\ uv &= \frac{1}{2} \\ v &= \frac{1}{2u}, \end{aligned}$$

where $\begin{cases} u = \frac{\sqrt{2}}{2}(y - x) \\ v = \frac{\sqrt{2}}{2}(y + x) \end{cases}$. Consider an arbitrary point (x, y) .

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\int_{?}^{?} 2\sqrt{x^2 + 1} dx$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= \left[x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}}
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned} & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\ &= 2 \left[x\sqrt{x^2 + 1} \right. \\ & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned} & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\ &= 2 \left[x\sqrt{x^2 + 1} \right. \\ & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}}
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned} & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\ &= 2 \left[x\sqrt{x^2 + 1} \right. \\ & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right)
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_0^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} \right. \\
 & \quad \left. + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right)
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2})
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




- Recall: integral can be solved via $x = \tan \theta$.

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




- Recall: integral can be solved via $x = \tan \theta$.
- Geometric interpretation of θ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




- Recall: integral can be solved via $x = \tan \theta$.
- Geometric interpretation of θ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.




- Recall: integral can be solved via $x = \tan \theta$.
- Geometric interpretation of θ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2 \ln (3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$

Example

Find the area locked b-n the hyperbolas $y = \pm\sqrt{x^2 + 1}$ and $x = \pm 2\sqrt{2}$.



- Recall: integral can be solved via $x = \tan \theta$.
- Geometric interpretation of θ ?

The area in question is:

$$\begin{aligned}
 & \int_{-2\sqrt{2}}^{2\sqrt{2}} 2\sqrt{x^2 + 1} dx \\
 &= 2 \left[x\sqrt{x^2 + 1} + \ln \left(\sqrt{x^2 + 1} + x \right) \right]_{-2\sqrt{2}}^{2\sqrt{2}} \\
 &= 2 \left(2\sqrt{2}\sqrt{(2\sqrt{2})^2 + 1} + \ln \left(\sqrt{(2\sqrt{2})^2 + 1} + 2\sqrt{2} \right) \right) \\
 &= 12\sqrt{2} + 2\ln(3 + 2\sqrt{2}) \\
 &\approx 20.496
 \end{aligned}$$