Calculus II Trigonometry review

Todor Milev

2019

Outline

- Review of trigonometry
 - The Trigonometric Functions
 - Trigonometric Identities
 - Trigonometric Identities and Complex Numbers
 - Graphs of the Trigonometric Functions

Outline

- Review of trigonometry
 - The Trigonometric Functions
 - Trigonometric Identities
 - Trigonometric Identities and Complex Numbers
 - Graphs of the Trigonometric Functions

Inverse Trigonometric Functions

License to use and redistribute


These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work.

as long as you reasonably acknowledge the original project.


- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
 https://creativecommons.org/licenses/by/3.0/us/
 and the links therein.

Fix unit circle, center O, coordinates (0,0).






Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$.



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A.



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


 $\sin \theta$

 $\cos \theta$

 $\tan \theta$

 $\cot\theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


 $\sin \theta$

 $\cos \theta$

 $\tan \theta$

 $\cot\theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}}$$

 $\cos \theta$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

 $\cos \theta$

 $\tan\theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

 $\cos \theta$

 $\tan \theta$

 $\cot\theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1}$$

 $\cos \theta$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

 $\cos \theta$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

 $\cos \theta$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$




Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$
 $\cos \theta = \frac{\text{adj}}{\text{hyp}}$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$




Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$
 $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$




Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$
 $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$




Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$
 $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1}$

 $\tan\theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

 $\tan \theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

 $\tan\theta$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1}$$

 $\cot \theta$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

 $\sec \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

∠OED = **?**



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

sec o



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

 $csc\theta$

/OED = ?



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

 $\sec \theta$

 $csc\theta$

/OED = ?



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

 $\sec \theta$

 $csc\theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

Todor Milev

Trigonometry review

 $csc\theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

 $\csc \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

 $\csc \theta$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

Todor Milev

Trigonometry review



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

 $=\theta$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

Todor Milev



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

Todor Milev

Trigonometry review



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

 $\csc \theta$

Todor Miley

Trigonometry review



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

 $=\theta$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$


$$\sec \theta$$

Todor Milev

Trigonometry review

 $csc\theta$

2019



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|}$$

$$\sec \theta$$

Todor Milev

Trigonometry review



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|}$$

 $\sec \theta$



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

 $=\theta$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let *OB* intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1}$$

$$\sec \theta$$



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

Todor Milev Trigonometry review



$$\beta = 180^{\circ} - 90^{\circ} - \theta$$

$$= 90^{\circ} - \theta$$

$$\angle OED = 180^{\circ} - 90^{\circ} - \beta$$

$$= 90^{\circ} - (90^{\circ} - \theta)$$

$$= \theta$$

Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.


$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\csc \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1}$$

$$\csc \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}}$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$



Fix unit circle, center O, coordinates (0,0). Let $\angle DOB = \theta$. Let OB intersect the circle at point A. Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1}$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$


$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.


 By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.
- For example, $\frac{\sin \theta}{\sin \theta} = 1$ is considered a valid trigonometric identity, although, when considered as a function, the left hand side is not defined for $\theta \neq 0$.



$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$


•
$$\csc \theta = \frac{1}{\sin \theta}$$

•
$$\sec \theta = \frac{1}{\cos \theta}$$


$$\cot \theta = \frac{1}{\tan \theta}$$

•
$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

• $\cot \theta = \frac{\cos \theta}{\sin \theta}$




$$\begin{aligned} \sin\theta &= \frac{y}{r} & \csc\theta &= \frac{r}{y} \\ \cos\theta &= \frac{x}{r} & \sec\theta &= \frac{r}{x} \\ \tan\theta &= \frac{y}{x} & \cot\theta &= \frac{x}{y} \end{aligned}$$



$$\begin{aligned} \sin\theta &= \frac{y}{r} & \csc\theta &= \frac{r}{y} \\ \cos\theta &= \frac{x}{r} & \sec\theta &= \frac{r}{x} \\ \tan\theta &= \frac{y}{x} & \cot\theta &= \frac{x}{y} \end{aligned}$$

$$\sin^2\theta + \cos^2\theta$$




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

$$\sin^2 \theta + \cos^2 \theta$$
$$= \frac{y^2}{r^2} + \frac{x^2}{r^2}$$



$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$


$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

$$\sin^2 \theta + \cos^2 \theta$$

$$= \frac{y^2}{r^2} + \frac{x^2}{r^2}$$

$$= \frac{y^2 + x^2}{r^2}$$




$$\begin{aligned} \sin\theta &= \frac{y}{r} & \csc\theta &= \frac{r}{y} \\ \cos\theta &= \frac{x}{r} & \sec\theta &= \frac{r}{x} \\ \tan\theta &= \frac{y}{x} & \cot\theta &= \frac{x}{y} \end{aligned}$$

$$\sin^2 \theta + \cos^2 \theta$$

$$= \frac{y^2}{r^2} + \frac{x^2}{r^2}$$

$$= \frac{y^2 + x^2}{r^2}$$

$$= \frac{r^2}{r^2}$$



$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$


$$\sin^2 \theta + \cos^2 \theta$$

$$= \frac{y^2}{r^2} + \frac{x^2}{r^2}$$

$$= \frac{y^2 + x^2}{r^2}$$

$$= \frac{r^2}{r^2}$$

$$= 1$$



$$\begin{aligned} \sin\theta &= \frac{y}{r} & \csc\theta &= \frac{r}{y} \\ \cos\theta &= \frac{x}{r} & \sec\theta &= \frac{r}{x} \\ \tan\theta &= \frac{y}{x} & \cot\theta &= \frac{x}{y} \end{aligned}$$


$$\sin^2 \theta + \cos^2 \theta$$

$$= \frac{y^2}{r^2} + \frac{x^2}{r^2}$$

$$= \frac{y^2 + x^2}{r^2}$$


$$= \frac{r^2}{r^2}$$

Therefore $\sin^2 \theta + \cos^2 \theta = 1$.



$$\begin{array}{ll} \sin\theta = \frac{y}{r} & \csc\theta = \frac{r}{y} \\ \cos\theta = \frac{x}{\xi} & \sec\theta = \frac{r}{x} \\ \tan\theta = \frac{y}{x} & \cot\theta = \frac{x}{y} \end{array}$$


Example $(\tan^2 \theta + 1 = \sec^2 \theta)$



$$\begin{array}{ll} \sin\theta = \frac{y}{r} & \csc\theta = \frac{r}{y} \\ \cos\theta = \frac{x}{t} & \sec\theta = \frac{r}{x} \\ \tan\theta = \frac{y}{x} & \cot\theta = \frac{x}{y} \end{array}$$

Example (tan² θ + 1 = sec² θ)

$$\sin^2\theta + \cos^2\theta = 1$$




$$\begin{array}{ll} \sin\theta = \frac{y}{r} & \csc\theta = \frac{r}{y} \\ \cos\theta = \frac{x}{r} & \sec\theta = \frac{r}{x} \\ \tan\theta = \frac{y}{x} & \cot\theta = \frac{x}{y} \end{array}$$

Example (tan² θ + 1 = sec² θ)

$$\sin^{2}\theta + \cos^{2}\theta = 1$$

$$\frac{\sin^{2}\theta}{\cos^{2}\theta} + \frac{\cos^{2}\theta}{\cos^{2}\theta} = \frac{1}{\cos^{2}\theta}$$



$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{l} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

Example (tan² θ + 1 = sec² θ)

$$\sin^{2}\theta + \cos^{2}\theta = 1$$

$$\frac{\sin^{2}\theta}{\cos^{2}\theta} + \frac{\cos^{2}\theta}{\cos^{2}\theta} = \frac{1}{\cos^{2}\theta}$$

$$\tan^{2}\theta + 1 = \sec^{2}\theta$$

2019

The remaining identities are consequences of the addition formulas:

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

Substitute -y for y, and use the fact that sin(-y) = -sin y and cos(-y) = cos y:

$$sin(x - y) = sin x cos y - cos x sin y$$

 $cos(x - y) = cos x cos y + sin x sin y$

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

2019

The remaining identities are consequences of the addition formulas:

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

To get the double angle formulas, substitute x for y:

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

To get the double angle formulas, substitute *x* for *y*:

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using $\cos^2 x = 1 - \sin^2 x$ and $\sin^2 x = 1 - \cos^2 x$:

$$cos(2x) = 2cos^2 x - 1$$

$$cos(2x) = 1 - 2sin^2 x$$

2019

The remaining identities are consequences of the addition formulas:

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

To get the double angle formulas, substitute *x* for *y*:

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using $\cos^2 x = 1 - \sin^2 x$ and $\sin^2 x = 1 - \cos^2 x$:

$$cos(2x) = 2cos^2 x - 1$$

$$cos(2x) = 1 - 2sin^2 x$$

To get the half-angle formulas, solve these equations for $\cos^2 x$ and $\sin^2 x$ respectively.

$$\cos^2 x = \frac{1 + \cos(2x)}{2}, \qquad \sin^2 x = \frac{1 - \cos(2x)}{2}$$

Todor Milev Trigonometry review

2019

The remaining identities are consequences of the addition formulas:

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

$$sin(x + y) = sin x cos y + cos x sin y$$

 $cos(x + y) = cos x cos y - sin x sin y$

Divide the first equation by the second, and then cancel $\cos x \cos y$ from the top and bottom:

$$tan(x + y) = \frac{tan x + tan y}{1 - tan x tan y}$$

$$sin(x + y) = sin x cos y + cos x sin y$$


 $cos(x + y) = cos x cos y - sin x sin y$

Divide the first equation by the second, and then cancel $\cos x \cos y$ from the top and bottom:

$$tan(x + y) = \frac{tan x + tan y}{1 - tan x tan y}$$

Do the same for the subtraction formulas:

$$tan(x - y) = \frac{tan x - tan y}{1 + tan x tan y}$$




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

 Positive angles are obtained by rotating counterclockwise.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{l} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{f} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.
- $\bullet \sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin\theta.$




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{f} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.
- $\bullet \sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin\theta.$




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.
- $\bullet \sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin\theta.$
- sin is an odd function.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$


$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then (x, -y) is on the terminal arm of $-\theta$.
- $\bullet \sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin\theta.$
- sin is an odd function.
- cos is an even function.




$$\begin{array}{ll} \sin\theta = \frac{y}{r} & \csc\theta = \frac{r}{y} \\ \cos\theta = \frac{x}{r} & \sec\theta = \frac{r}{x} \\ \tan\theta = \frac{y}{x} & \cot\theta = \frac{x}{y} \end{array}$$



$$\sin \theta = \frac{y}{\cos \theta} = \frac{1}{2}$$

$$\begin{array}{ll} \sin\theta = \frac{y}{r} & \csc\theta = \frac{r}{y} \\ \cos\theta = \frac{x}{r} & \sec\theta = \frac{r}{x} \\ \tan\theta = \frac{y}{x} & \cot\theta = \frac{x}{y} \end{array}$$

• 2π represents a full rotation.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r.




$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r.
- $\sin(\theta + 2\pi) = \sin \theta$.



$$\sin \theta = \frac{y}{r} \quad \csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r} \quad \sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r.
- $\sin(\theta + 2\pi) = \sin \theta$.
- We say sin and cos are 2π -periodic.

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
 .

The number i is called the imaginary unit.

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number i is called the imaginary unit.

$$\pm \sqrt{-1} = i.$$

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\pm \sqrt{-1} = i$.

The set of complex numbers \mathbb{C} is defined as the set

$$\{a+bi|a,b-\text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

$$\sqrt{-1} = i$$
.

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
 .

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

$$\sqrt{-1}=i$$
.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i$$

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

$$\sqrt{-1}=i$$
.

Complex addition/subtraction

$$(\mathbf{a} + b\mathbf{i}) \pm (\mathbf{c} + d\mathbf{i}) = (\mathbf{a} \pm \mathbf{c}) + (\mathbf{b} \pm \mathbf{d})\mathbf{i}$$

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
 .

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

$$\sqrt{-1}=i$$
.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i$$

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

The set of complex numbers ${\mathbb C}$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number i is a number for which

$$i^2 = -1$$
 .

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
 .

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

The set of complex numbers $\ensuremath{\mathbb{C}}$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a+bi)(c+di) = ac+adi+bci+bdi^2 = ac+adi+bci-bdi$$

The set of complex numbers $\ensuremath{\mathbb{C}}$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a+bi)(c+di) = ac + adi + bci + bdi^2 = ac + adi + bci - bd$$

= $(ac - bd) + i(ad + bc)$

2019

Definition (Complex numbers)

The set of complex numbers $\mathbb C$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a+bi)(c+di) = ac+adi+bci+bdi^2 = ac+adi+bci-bdi$$

= $(ac-bd)+i(ad+bc)$

2019

Definition (Complex numbers)

The set of complex numbers $\ensuremath{\mathbb{C}}$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

$$(a+bi)(c+di) = ac+adi+bci+bdi^2 = ac+adi+bci-bdi$$

= $(ac-bd)+i(ad+bc)$

The set of complex numbers $\ensuremath{\mathbb{C}}$ is defined as the set

$$\{a + bi | a, b - \text{real numbers}\},\$$

where the number *i* is a number for which

$$i^2 = -1$$
.

The number *i* is called the imaginary unit. By definition, $\sqrt{-1} = i$.

Complex addition/subtraction

$$(a+bi)\pm(c+di)=(a\pm c)+(b\pm d)i \quad .$$

Complex multiplication

$$(a+bi)(c+di) = ac+adi+bci+bdi^2 = ac+adi+bci-bd$$

= $(ac-bd)+i(ad+bc)$

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + \frac{(-1)^n x^{2n+1}}{(2n+1)!} + \dots$$
$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + \frac{(-1)^n x^{2n}}{(2n)!} + \dots$$

$$e^{x} = 1 + x + \frac{x^{2}}{2!} + \frac{x^{3}}{3!} + \dots + \frac{x^{n}}{n!} + \dots$$

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x \qquad -\frac{x^3}{3!} \qquad +\frac{x^5}{5!} - \dots$$

Rearrange.

15/40

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$cos x = 1 -\frac{x^2}{2!} +\frac{x^4}{4!} + \dots$$

$$e^{ix} = 1 + ix + \frac{(ix)^2}{2!} + \frac{(ix)^3}{3!} + \frac{(ix)^4}{4!} + \frac{(ix)^5}{5!} + \dots$$

Rearrange. Plug-in z = ix.

15/40

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x \qquad -\frac{x^3}{3!} \qquad +\frac{x^5}{5!} - \dots$$

$$\frac{\cos x = 1 \quad -\frac{x^2}{2!} \quad +\frac{x^4}{4!} \quad +\dots}{e^{ix} = 1 \quad +ix \quad -\frac{x^2}{2!} \quad -i\frac{x^3}{3!} \quad +\frac{x^4}{4!} \quad +i\frac{x^5}{5!} \quad -\dots}$$

Rearrange. Plug-in z = ix. Use $i^2 = -1$.

15/40

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i\sin x = ix$$
 $-i\frac{x^3}{3!}$ $+i\frac{x^5}{5!}$ $-\dots$

$$\frac{\cos x = 1 \quad -\frac{x^2}{2!} \quad +\frac{x^4}{4!} \quad + \dots}{e^{ix} = 1 \quad +ix \quad -\frac{x^2}{2!} \quad -i\frac{x^3}{3!} \quad +\frac{x^4}{4!} \quad +i\frac{x^5}{5!} \quad - \dots}$$

Rearrange. Plug-in z = ix. Use $i^2 = -1$. Multiply $\sin x$ by i.

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \cdots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i\sin x = ix \qquad -i\frac{x^3}{3!} \qquad +i\frac{x^5}{5!} - \dots$$

$$cos x = 1 -\frac{x^2}{2!} +\frac{x^4}{4!} + \dots$$

$$e^{ix} = 1 +ix -\frac{x^2}{2!} -i\frac{x^3}{3!} +\frac{x^4}{4!} +i\frac{x^5}{5!} - \dots$$

Rearrange. Plug-in z = ix. Use $i^2 = -1$. Multiply $\sin x$ by i. Add to get $e^{ix} = \cos x + i \sin x$.

- $e^{ix} = \cos x + i \sin x$
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$
- $e^0 = 1$

(Euler's Formula). (exponentiation rule: valid for \mathbb{C}).

(exponentiation rule).

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

All trigonometric formulas can be easily derived using the above formulas.

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

$$e^{i(x+y)}$$
 —

2019

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

$$e^{i(x+y)} = \cos(x+y) + i\sin(x+y)$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula). • $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $ullet e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

$$e^{i(x+y)} = \cos(x+y) + i\sin(x+y)$$

 $e^{ix}e^{iy} = \cos(x+y) + i\sin(x+y)$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

Proof.

$$e^{i(x+y)} = \cos(x+y) + i\sin(x+y)$$

$$e^{ix}e^{iy} = \cos(x+y) + i\sin(x+y)$$

$$(\cos x + i\sin x)(\cos y + i\sin y) = \cos(x+y) + i\sin(x+y)$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(x + y) = sin x cos y + sin y cos x$$

 $cos(x + y) = cos x cos y - sin x sin y$

Proof.

$$e^{i(x+y)} = \cos(x+y) + i\sin(x+y)$$

$$e^{ix}e^{iy} = \cos(x+y) + i\sin(x+y)$$

$$(\cos x + i\sin x)(\cos y + i\sin y) = \cos(x+y) + i\sin(x+y)$$

$$\cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) = \cos(x+y) + i\sin(x+y)$$

2019

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\frac{\sin(x+y)}{\cos(x+y)} = \frac{\sin x \cos y + \sin y \cos x}{\cos x \cos y - \sin x \sin y}$$

Proof.

```
e^{i(x+y)} = \cos(x+y) + i\sin(x+y)
e^{ix}e^{iy} = \cos(x+y) + i\sin(x+y)
(\cos x + i\sin x)(\cos y + i\sin y) = \cos(x+y) + i\sin(x+y)
\cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) = \cos(x+y) + i\sin(x+y)
```

Compare coefficient in front of *i* and

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin(x + y) = \sin x \cos y + \sin y \cos x$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y.$$

Proof.

```
e^{i(x+y)} = \cos(x+y) + i\sin(x+y)
e^{ix}e^{iy} = \cos(x+y) + i\sin(x+y)
(\cos x + i\sin x)(\cos y + i\sin y) = \cos(x+y) + i\sin(x+y)
\cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) = \cos(x+y) + i\sin(x+y)
```

Compare coefficient in front of i and remaining terms to get the desired equalities.

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

 $e^{ix} = \cos x + i \sin x$

(Euler's Formula).

• $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$

(exponentiation rule: valid for \mathbb{C}). (exponentiation rule).

- $e^0 = 1$
- $\bullet \sin(-X) = -\sin X, \cos(-X) = \cos X$

(easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

$$1 = e^0$$

 $e^{ix} = \cos x + i \sin x$

(Euler's Formula).

 $\bullet e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$

- (exponentiation rule: valid for \mathbb{C}). (exponentiation rule).

(easy to remember).

Example

• $e^0 = 1$

$$\sin^2 x + \cos^2 x = 1$$

$$\begin{array}{rcl}
1 & = & e^0 \\
 & = & e^{ix-ix} =
\end{array}$$

 $e^{ix} = \cos x + i \sin x$

(Euler's Formula). (exponentiation rule: valid for \mathbb{C}).

• $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$

(exponentiation rule).

 $\bullet \sin(-x) = -\sin x, \cos(-x) = \cos x$

(easy to remember).

Example

• $e^0 = 1$

$$\sin^2 x + \cos^2 x = 1$$

$$\begin{array}{rcl}
1 & = & e^0 \\
 & = & e^{ix-ix} = e^{ix}e^{-ix} =
\end{array}$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

1 =
$$e^0$$

= $e^{ix-ix} = e^{ix}e^{-ix} = (\cos x + i\sin x)(\cos(-x) + i\sin(-x))$

2019

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

$$1 = e^{0}
= e^{ix-ix} = e^{ix}e^{-ix} = (\cos x + i\sin x)(\cos(-x) + i\sin(-x))
= (\cos x + i\sin x)(\cos x - i\sin x) =$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

$$1 = e^{0}
= e^{ix-ix} = e^{ix}e^{-ix} = (\cos x + i\sin x)(\cos(-x) + i\sin(-x))
= (\cos x + i\sin x)(\cos x - i\sin x) = \cos^{2} x - i^{2}\sin^{2} x$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$1 = e^{0}
= e^{ix-ix} = e^{ix}e^{-ix} = (\cos x + i\sin x)(\cos(-x) + i\sin(-x))
= (\cos x + i\sin x)(\cos x - i\sin x) = \cos^{2} x - i^{2}\sin^{2} x
= \cos^{2} x + \sin^{2} x$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(2x) = 2 sin x cos x$$

$$cos(2x) = cos^2 x - sin^2 x$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
 - $e^{ix}e^{iy}=e^{ix+iy}=e^{i(x+y)}$ (exponentiation rule: valid for $\mathbb C$).
 - $e^0 = 1$ (exponentiation rule).
 - $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(2x) = 2 sin x cos x$$

$$cos(2x) = cos^2 x - sin^2 x .$$

Proof.

$$e^{i(2x)}$$

 $e^{ix} = \cos x + i \sin x$

(Euler's Formula).

• $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$

- (exponentiation rule: valid for \mathbb{C}). (exponentiation rule).
- $\bullet \sin(-x) = -\sin x, \cos(-x) = \cos x$

(easy to remember).

Example

• $e^0 = 1$

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula). • $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

 $e^{ix}e^{ix} = \cos(2x) + i\sin(2x)$

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(2x) = 2 sin x cos x$$

$$cos(2x) = cos^2 x - sin^2 x$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

$$e^{ix}e^{ix} = \cos(2x) + i\sin(2x)$$

$$(\cos x + i\sin x)(\cos x + i\sin x) = \cos(2x) + i\sin(2x)$$

Todor Milev Trigonometry review 2019

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule)
- $e^0 = 1$ (exponentiation rule). • $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\sin(2x) = 2\sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

$$e^{ix}e^{ix} = \cos(2x) + i\sin(2x)$$

$$(\cos x + i\sin x)^2 = (\cos x + i\sin x)(\cos x + i\sin x) = \cos(2x) + i\sin(2x)$$

$$\cos^2 x - \sin^2 x + i(2\sin x\cos x) = \cos(2x) + i\sin(2x)$$

Todor Milev Trigonometry review 2019

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\frac{\sin(2x)}{\cos(2x)} = \frac{2\sin x \cos x}{\cos^2 x - \sin^2 x}.$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

$$e^{ix}e^{ix} = \cos(2x) + i\sin(2x)$$

$$(\cos x + i\sin x)^2 = (\cos x + i\sin x)(\cos x + i\sin x) = \cos(2x) + i\sin(2x)$$

$$\cos^2 x - \sin^2 x + i(2\sin x\cos x) = \cos(2x) + i\sin(2x)$$

Compare coefficient in front of i and

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix}e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}). • $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$sin(2x) = 2 sin x cos x$$

$$cos(2x) = cos^2 x - sin^2 x .$$

Proof.

$$e^{i(2x)} = \cos(2x) + i\sin(2x)$$

$$e^{ix}e^{ix} = \cos(2x) + i\sin(2x)$$

$$(\cos x + i\sin x)^2 = (\cos x + i\sin x)(\cos x + i\sin x) = \cos(2x) + i\sin(2x)$$

$$\cos^2 x - \sin^2 x + i(2\sin x\cos x) = \cos(2x) + i\sin(2x)$$

Compare coefficient in front of i and remaining terms to get the desired equalities.

Todor Milev Trigonometry review 2019

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

Todor Milev Trigonometry review 2019

2019

• Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.

Example

Express sin(3x) and cos(3x) via cos x and sin x.

$$\cos(3x) + i\sin(3x) \\ = e^{3ix}$$

Example

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

$$\cos(\frac{3x}{3x}) + i\sin(\frac{3x}{3x}) = e^{3ix}$$

Example

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

$$\cos(3x) + i\sin(3x)$$
$$= e^{3ix}$$

Example

Express sin(3x) and cos(3x) via cos x and sin x.

$$\cos(3x) + i\sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3$$

Example

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

$$\cos(3x) + i\sin(3x)$$
$$= e^{3ix}$$

$$= \left(e^{ix}\right)^3 = (\cos x + i \sin x)^3$$

Euler's f-la

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = ?$

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

$$cos(3x) + i sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3 = (cos x + i sin x)^3$$

Euler's f-la

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
= e^{3ix}
= $(e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
= $\cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
= e^{3ix}
= $(e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
= $\cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
= e^{3ix}
= $(e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
= $\cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3 = (\cos x + i\sin x)^3$$

$$= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$$

$$= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$$
Euler's f-la

2019

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

Example

$$\cos(3x) + i\sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3 = (\cos x + i\sin x)^3$$

$$= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$$

$$= \cos^3 x + 3i\cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x$$
Euler's f-la

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$

$$= e^{3ix}$$

$$= (e^{ix})^3 = (\cos x + i\sin x)^3$$

$$= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$$

$$= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$$
Euler's f-la

2019

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

Example

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
 $= e^{3ix}$
 $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
 $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
 $= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
 $= \cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$

2019

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

Example

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
 $= e^{3ix}$
 $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
 $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
 $= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
 $= \cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
 $= e^{3ix}$
 $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
 $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
 $= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
 $= \cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$
 $= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$. $\cos(3x) + i\sin(3x)$ | Euler's f-la $= e^{3ix}$ $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$

$$= \cos^3 x + 3i\cos^2 x \sin x + 3i^2 \cos x \sin^2 x + i^3 \sin^3 x$$

$$=\cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$$

$$= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$$

Use
$$i^2 = -1$$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
 $= e^{3ix}$
 $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
 $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
 $= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
 $= \cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$
 $= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
= e^{3ix}
= $(e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
= $\cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
= $\cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
= $\cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$
= $(\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.

Express $\sin(3x)$ and $\cos(3x)$ via $\cos x$ and $\sin x$.

$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
= e^{3ix}
= $(e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
= $\cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
= $\cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
= $\cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$
= $(\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$

The real parts of the starting and final expression must be equal; therefore:

$$\cos(3x) = \cos^3 x - 3\cos x \sin^2 x$$

- Recall Euler's formula: $e^{i\alpha} = \cos \alpha + i \sin \alpha$.
- Recall the formula: $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$.


Express sin(3x) and cos(3x) via cos x and sin x.


$$\cos(3x) + i\sin(3x)$$
 | Euler's f-la
 $= e^{3ix}$
 $= (e^{ix})^3 = (\cos x + i\sin x)^3$ | Euler's f-la
 $= \cos^3 x + 3\cos^2 x (i\sin x) + 3\cos x (i\sin x)^2 + (i\sin x)^3$
 $= \cos^3 x + 3i\cos^2 x \sin x + 3i^2\cos x \sin^2 x + i^3\sin^3 x$
 $= \cos^3 x + 3i\cos^2 x \sin x - 3\cos x \sin^2 x - i\sin^3 x$ | Use $i^2 = -1$
 $= (\cos^3 x - 3\cos x \sin^2 x) + i(3\cos^2 x \sin x - \sin^3 x)$

The real parts of the starting and final expression must be equal; likewise the imaginary parts must be equal; therefore:


$$\cos(3x) = \cos^3 x - 3\cos x \sin^2 x$$

$$\sin(3x) = 3\cos^2 x \sin x - \sin^3 x$$






• $\sin x$ has zeroes at $n\pi$ for all integers n.




- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.




- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- -1 ≤ $\sin x$ ≤ 1.


2019



- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- $-1 \le \sin x \le 1.$
- $-1 < \cos x < 1$.




- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- $-1 < \sin x < 1$.
- $-1 < \cos x < 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right



- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- -1 ≤ $\sin x$ ≤ 1.
- $-1 < \cos x < 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right


Todor Milev Trigonometry review 2019



- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- $-1 < \sin x < 1$.
- $-1 < \cos x < 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right

Todor Milev Trigonometry review 2019


Graphs of the Trigonometric Functions




- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- -1 ≤ $\sin x$ ≤ 1.
- $-1 < \cos x < 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right we get the graph of $\sin x$.


Todor Milev Trigonometry review 2019

Graphs of the Trigonometric Functions





- $\sin x$ has zeroes at $n\pi$ for all integers n.
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n.
- -1 ≤ $\sin x$ ≤ 1.
- \bullet -1 < cos *x* < 1.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right we get the graph of $\sin x$. This is a consequence of $\cos \left(x \frac{\pi}{2}\right) = \sin x$.








$$= \tan x$$
 $y = \cot x$















If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis, we get the graph of $\cot x$.





If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis, we get the graph of $\cot x$. This follows from $\tan \left(x \pm \frac{\pi}{2}\right) = -\cot x$.














• sin x isn't one-to-one.




- sin x isn't one-to-one.
- It is if we restrict the domain to $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.




- sin x isn't one-to-one.
- It is if we restrict the domain to $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.
- Then it has an inverse function.
- We call it arcsin or sin⁻¹.




- sin x isn't one-to-one.
- It is if we restrict the domain to $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.
- Then it has an inverse function.
- We call it arcsin or sin⁻¹.





- sin x isn't one-to-one.
- It is if we restrict the domain to $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.
- Then it has an inverse function.
- We call it arcsin or sin⁻¹.
- $\arcsin x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.



Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

• arcsin y = the appropriate angle whose sine equals y.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

• $\arcsin y = the appropriate angle whose sine equals y$.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

$$\bullet \sin\left(\frac{?}{?}\right) = \frac{1}{2}.$$

• arcsin y = the appropriate angle whose sine equals y.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

$$\bullet \sin\left(\mathbf{?}\right) = \frac{1}{2}.$$

• arcsin y = the appropriate angle whose sine equals y.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

•
$$\sin\left(\mathbf{?}\right) = \frac{1}{2}$$
.

• arcsin y = the appropriate angle whose sine equals y.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

•
$$\sin\left(\frac{?}{2}\right) = \frac{1}{2}$$
.

• arcsin y = the appropriate angle whose sine equals y.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

•
$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$
.

- arcsin y = the appropriate angle whose sine equals y.
- Important: the output angle must lie in the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

- $\bullet \, \sin\left(\frac{\pi}{6}\right) = \frac{1}{2}.$
- $\bullet -\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}.$

- arcsin y = the appropriate angle whose sine equals y.
- Important: the output angle must lie in the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

Find
$$\arcsin\left(\frac{1}{2}\right)$$
.

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$.
- $\bullet -\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}.$
- Therefore $\arcsin\left(\frac{1}{2}\right) = \frac{\pi}{6}$.


Find
$$\tan \left(\arcsin \left(\frac{1}{3}\right)\right)$$
.

Find $\tan \left(\arcsin \left(\frac{1}{3}\right)\right)$.

• Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.


Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.




Find $\tan \left(\arcsin \left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled.




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$




Find $\tan \left(\arcsin \left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin \left(\frac{1}{3}\right)$.




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin(\frac{1}{3})$.
- Length of adjacent side = ?




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin \left(\frac{1}{3}\right)$.
- Length of adjacent side = $\sqrt{3^2 1^2}$




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin(\frac{1}{3})$.
- Length of adjacent side = $\sqrt{3^2 1^2} = \sqrt{8} = 2\sqrt{2}$.




Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin(\frac{1}{3})$.
- Length of adjacent side = $\sqrt{3^2 1^2} = \sqrt{8} = 2\sqrt{2}$.
- Then tan $\left(\arcsin\left(\frac{1}{3}\right)\right) = ?$



Find $\tan \left(\arcsin\left(\frac{1}{3}\right)\right)$.

- Let $\theta = \arcsin\left(\frac{1}{3}\right)$, so $\sin \theta = \frac{1}{3}$.
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle θ be as labeled. Then $\sin \theta = \frac{1}{3}$ and so $\theta = \arcsin \left(\frac{1}{3}\right)$.
- Length of adjacent side = $\sqrt{3^2 1^2} = \sqrt{8} = 2\sqrt{2}$.
- Then $\tan \left(\arcsin \left(\frac{1}{3}\right)\right) = \frac{1}{2\sqrt{2}}$.



Find arcsin(sin(1.5)).

Find arcsin(sin(1.5)).

• $\frac{\pi}{2} \approx$?

Find arcsin(sin(1.5)).


• $\frac{\pi}{2} \approx 1.57$.


Find $\arcsin(\sin(1.5))$.

- $\frac{\pi}{2} \approx 1.57$.
- Therefore $-\frac{\pi}{2} \le 1.5 \le \frac{\pi}{2}$.

Find $\arcsin(\sin(1.5))$.


- $\frac{\pi}{2} \approx 1.57$.
- Therefore $-\frac{\pi}{2} \le 1.5 \le \frac{\pi}{2}$.
- Therefore $\arcsin(\sin 1.5) = 1.5$.







Find arcsin(sin 2).


• 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.







- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.







- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.







- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.







- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.







- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.






- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.






- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.


$$a = ?$$






- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.

$$a = \pi - 2$$
.







Find arcsin(sin 2).

- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.

$$a = \pi - 2$$
.


Therefore $\arcsin(\sin 2) = \arcsin(\sin a)$






- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.

$$a = \pi - 2$$
.
Therefore $\arcsin(\sin 2) = \arcsin(\sin a)$
 $= a$







Find arcsin(sin 2).

- 2 is not between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$.
- We need the angle a between $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ for which $\sin 2 = \sin a$.

$$a = \pi - 2.$$

Therefore $\arcsin(\sin 2) = \arcsin(\sin a)$ = $a = \pi - 2$.





$$\frac{\mathsf{d}}{\mathsf{d}x}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: ? =?

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' =$?

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' =$?

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' = 1$

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' = 1$

$$y' = \frac{1}{\cos y}$$

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' = 1$

$$y' = \frac{1}{\cos y}$$
$$= \frac{1}{\pm \sqrt{1 - \sin^2 y}}$$

Todor Milev

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.

Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' = 1$

$$y' = \frac{1}{\cos y}$$

$$= \frac{1}{\pm \sqrt{1 - \sin^2 y}}$$

But
$$\cos y > 0$$
: $= \frac{1}{\sqrt{1 - \sin^2 y}}$

$$\frac{\mathsf{d}}{\mathsf{d}x}\left(\arcsin x\right) = \frac{1}{\sqrt{1-x^2}}, \qquad -1 < x < 1.$$

Proof.

Let
$$y = \arcsin x$$
.


Then
$$\sin y = x$$
 and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.

Differentiate implicitly: $\cos y \cdot y' = 1$


Highlight cos
$$y \cdot y' = 1$$

$$y' = \frac{1}{\cos y}$$


$$= \frac{1}{\pm \sqrt{1 - \sin^2 y}}$$
But $\cos y > 0$:
$$= \frac{1}{\sqrt{1 - \sin^2 y}} = \frac{1}{\sqrt{1 - x^2}}.$$




- Domain: ?
- Range: ?
- 3 $\arcsin x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- arcsin(sin x) = x for $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$.
- $\sin(\arcsin x) = x$ for $-1 \le x \le 1$.




- Domain: ?
- Range: ?
- o arcsin $x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- arcsin(sin x) = x for $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$.
- $\sin(\arcsin x) = x \text{ for }$ $-1 \le x \le 1.$




- Domain: [-1,1].
- Range: ?
- arcsin $x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- arcsin(sin x) = x for $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$.
- $\sin(\arcsin x) = x \text{ for }$ $-1 \le x \le 1.$




- Domain: [-1,1].
- Range: ?
- arcsin $x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- arcsin(sin x) = x for $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$.
- $\sin(\arcsin x) = x \text{ for }$ $-1 \le x \le 1.$




- **●** Domain: [-1,1].
- **2** Range: $[-\frac{\pi}{2}, \frac{\pi}{2}]$.
- arcsin $x = y \Leftrightarrow \sin y = x$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- arcsin(sin X) = X for $-\frac{\pi}{2} \le X \le \frac{\pi}{2}$.
- $\sin(\arcsin x) = x \text{ for }$ $-1 \le x \le 1.$




• Same for cos x.




- Same for cos x.
- Restrict the domain to $[0, \pi]$.





- Same for cos x.
- Restrict the domain to $[0, \pi]$.
- The inverse is called arccos or cos⁻¹.







- Restrict the domain to $[0, \pi]$.
- The inverse is called arccos or cos⁻¹.









- Restrict the domain to $[0, \pi]$.
- The inverse is called arccos or cos⁻¹.
- $\operatorname{arccos}(x) = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.







- Domain:
- Range:
- arccos $x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$
- $d(\operatorname{arccos} x) = -\frac{1}{\sqrt{1-x^2}}.$




- Domain: ?
- Range:
- arccos $x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$




- **●** Domain: [−1,1].
- Range:
- arccos $x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$
- $d(\operatorname{arccos} x) = -\frac{1}{\sqrt{1-x^2}}.$



- Domain: [-1,1].
- Range: ?
- arccos $x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$



- Domain: [-1,1].
- **2** Range: $[0, \pi]$.
- 3 $\operatorname{arccos} x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$



- Domain: [-1,1].
- **2** Range: $[0, \pi]$.
- 3 $\arccos x = y \Leftrightarrow \cos y = x$ and $0 \le y \le \pi$.
- arccos(cos x) = x for $0 \le x \le \pi$.
- $\begin{array}{l} \mathbf{5} & \cos(\arccos x) = x \text{ for} \\ -1 \leq x \leq 1. \end{array}$
- (The proof is similar to the proof of the formula for the derivative of $\frac{d}{dx}(arccos x) = -\frac{1}{\sqrt{1-x^2}}$.

Rewrite $\sin(2\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$.

sin(2 arccos(x))

Rewrite $\sin(2\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$.

sin(2 arccos(x))

Rewrite $\sin(2\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

sin(2 arccos(x))

$$sin(2 \frac{arccos(x)}{arccos(x)}) = sin(2y)$$

Set
$$y = \arccos x$$

$$\sin(2\arccos(x)) = \sin(2y)$$
= ?

Set
$$y = \arccos x$$

Express via $\sin y, \cos y$

$$sin(2 arccos(x)) = \frac{sin(2y)}{2 cos y sin y}$$

Set
$$y = \arccos x$$

Express via $\sin y$, $\cos y$

$$\sin(2\arccos(x)) = \sin(2y)$$

$$= 2\cos y \sin y$$

$$= 2\cos y \left(\pm\sqrt{1-\cos^2 y}\right)$$
Set $y = \arccos x$
Express via $\sin y$, $\cos y$
Express $\sin y$ via $\cos y$

$$\sin(2 \arccos(x)) = \sin(2y)$$

$$= 2 \cos y \sin y$$

$$= 2 \cos y \left(\pm \sqrt{1 - \cos^2 y}\right)$$

$$= 2 \cos y \sqrt{1 - \cos^2 y}$$

Set
$$y = \arccos x$$

Express via $\sin y$, $\cos y$
Express $\sin y$ via $\cos y$
 $\sin y > 0$ because
 $0 < y < \pi$

$$sin(2 \operatorname{arccos}(x)) = \sin(2y)$$

$$= 2 \cos y \sin y$$

$$= 2 \cos y \left(\pm \sqrt{1 - \cos^2 y} \right)$$

$$= 2 \cos y \sqrt{1 - \cos^2 y}$$

Set
$$y = \arccos x$$

Express via $\sin y$, $\cos y$
Express $\sin y$ via $\cos y$
 $\sin y > 0$ because
 $0 < y < \pi$

$$sin(2 \operatorname{arccos}(x)) = \sin(2y)$$

$$= 2 \cos y \sin y$$

$$= 2 \cos y \left(\pm \sqrt{1 - \cos^2 y}\right)$$

$$= 2 \cos y \sqrt{1 - \cos^2 y}$$

$$= 2x \sqrt{1 - x^2}$$

Set
$$y = \arccos x$$

Express via $\sin y$, $\cos y$
Express $\sin y$ via $\cos y$
 $\sin y > 0$ because
 $0 \le y \le \pi$
use $x = \cos y$

Rewrite $\sin(2\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$.

To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

$$sin(2 \operatorname{arccos}(x)) = \sin(2y)
= 2 \cos y \sin y
= 2 \cos y \left(\pm \sqrt{1 - \cos^2 y}\right) = 2 \cos y \sqrt{1 - \cos^2 y}$$

$$= 2 \cos y \sqrt{1 - \cos^2 y}$$

$$= 2x\sqrt{1 - x^2}$$
Use

Set $y = \arccos x$ Express via $\sin y$, $\cos y$ Express $\sin y$ via $\cos y$ $\sin y > 0$ because $0 \le y \le \pi$ use $x = \cos y$

Rewrite $cos(3 \operatorname{arccos}(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$.

cos(3 arccos(x))

Rewrite $cos(3 \operatorname{arccos}(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify arccos x we try to use cos(arccos x) = x.

cos(3 arccos(x))

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

cos(3 arccos(x))

$$cos(3 \operatorname{arccos}(x)) = cos(3y)$$

$$y = \arccos x$$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$
 | $y = \arccos x$

$$cos(3 \operatorname{arccos}(x)) = cos(3y) = \frac{cos(2y + y)}{2}$$
 | $y = \operatorname{arccos}(x)$ | Angle sum f-la

$$cos(3 \operatorname{arccos}(x)) = cos(3y) = cos(2y + y)$$

= $cos(2y) cos y - sin(2y) sin y$ | $y = \operatorname{arccos} x$
Angle sum f-la

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (?) \cos y$$

$$-? \sin y$$

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$-? \sin y$$

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- ? \sin y$$

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the \cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$y = \arccos x$$
Angle sum f-la
$$Express via$$

$$\sin y, \cos y$$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$-2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$y = \arccos x$$
Angle sum f-la
Express via
$$\sin y, \cos y$$

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

 $y = \arccos x$

Rewrite $\cos(3\arccos(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\arccos x$ we try to use $\cos(\arccos x) = x$. Therefore our aim is to rewrite the expression only using the cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

$$= \cos^3 y - 3(?)$$

$$\cos y$$

$$y = \arccos x$$
Angle sum f-late to the following sin y, cos y
$$\sin y + \cos y = \cos y$$
Express sin y via cos y
$$\cos y = \cos^3 y - 3(?)$$

Angle sum f-la

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

$$= \cos^3 y - 3(1 - \cos^2 y)\cos y$$

$$y = \arccos x$$
Angle sum f-la
Express via
$$\sin y, \cos y$$
Express $\sin y$
via $\cos y$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

$$= \cos^3 y - 3(1 - \cos^2 y)\cos y$$

$$= 4\cos^3 y - 3\cos y$$

$$= \cos^3 y - 3\cos y$$

$$= \cos^3 y - 3\cos y$$

$$= \cos^3 y - 3\cos y$$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

$$= \cos^3 y - 3(1 - \cos^2 y)\cos y$$

$$= 4\cos^3 y - 3\cos y$$

$$= \cos^3 y - 3\cos y$$

$$y = \arccos y$$
Angle sum f-la
Express via
$$\sin y, \cos y$$

$$\operatorname{Express } \sin y$$
via $\cos y$

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$

$$= \cos^3 y - 3(1 - \cos^2 y)\cos y$$

$$= 4\cos^3 y - 3\cos y$$

$$= 4x^3 - 3x$$

$$y = \arccos x$$
Angle sum f-la
Express via
$$\sin y, \cos y$$
Express $\sin y$
via $\cos y$

Rewrite $cos(3 \operatorname{arccos}(x))$ as an algebraic expression of x and $\sqrt{1-x^2}$. To simplify $\operatorname{arccos} x$ we try to use $cos(\operatorname{arccos} x) = x$. Therefore our aim

is to rewrite the expression only using the cos function.

$$\cos(3\arccos(x)) = \cos(3y) = \cos(2y + y)$$

$$= \cos(2y)\cos y - \sin(2y)\sin y$$

$$= (\cos^2 y - \sin^2 y)\cos y$$

$$- 2\sin y\cos y\sin y$$

$$= \cos^3 y - \sin^2 y\cos y - 2\sin^2 y\cos y$$

$$= \cos^3 y - 3\sin^2 y\cos y$$


$$= \cos^3 y - 3(1 - \cos^2 y)\cos y$$


$$= 4\cos^3 y - 3\cos y$$


$$= 4x^3 - 3x$$

$$| y = \arccos x$$
Angle sum f-la
Express via
$$\sin y, \cos y$$


$$\operatorname{Express } \sin y$$
via $\cos y$









- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: ?
- Range of arctan:




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan:




- tan x isn't one-to-one.
- Restrict the domain to (-π/2, π/2).
 The inverse is called tan⁻¹ or
- arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: ?




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- $\lim_{x\to\infty} \arctan x =$?
- $\lim_{x \to -\infty} \arctan x =$




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- $\lim_{x\to\infty} \arctan x = \frac{\pi}{2}$.
- $\lim_{x \to -\infty} \arctan x =$



- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- $\lim_{x\to\infty} \arctan x = \frac{\pi}{2}$.
- $\lim_{x \to -\infty} \arctan x =$?




- tan x isn't one-to-one.
- Restrict the domain to $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- The inverse is called tan⁻¹ or arctan.
- $\arctan x = y \Leftrightarrow \tan y = x$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
- Domain of arctan: $(-\infty, \infty)$.
- Range of arctan: $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
- $\bullet \lim_{x\to\infty}\arctan x=\frac{\pi}{2}.$
- $\lim_{x \to -\infty} \arctan x = -\frac{\pi}{2}$.




Simplify the expression cos(arctan x).


• Let $y = \arctan x$, so $\tan y = x$.




- Let $y = \arctan x$, so $\tan y = x$.
- Draw a right triangle with opposite *x* and adjacent 1.




- Let $y = \arctan x$, so $\tan y = x$.
- Draw a right triangle with opposite x and adjacent 1.
- Length of hypotenuse = ?




- Let $y = \arctan x$, so $\tan y = x$.
- Draw a right triangle with opposite x and adjacent 1.
- Length of hypotenuse = $\sqrt{1^2 + x^2}$.



- Let $y = \arctan x$, so $\tan y = x$.
- Draw a right triangle with opposite x and adjacent 1.
- Length of hypotenuse = $\sqrt{1^2 + x^2}$.
- Then cos(arctan x) = ?



- Let $y = \arctan x$, so $\tan y = x$.
- Draw a right triangle with opposite x and adjacent 1.
- Length of hypotenuse = $\sqrt{1^2 + x^2}$.
- Then $\cos(\arctan x) = \frac{1}{\sqrt{1+x^2}}$.



Evaluate

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right).$$

Evaluate

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right).$$

$$\frac{1}{x-2} \to \infty$$
 as $x \to 2^+$.

Evaluate

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right).$$

$$\frac{1}{x-2} \to \infty$$
 as $x \to 2^+$.

Therefore

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right)=\mathbf{?}$$

Evaluate

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right).$$

$$\frac{1}{x-2} \to \infty$$
 as $x \to 2^+$.

Therefore

$$\lim_{x\to 2^+}\arctan\left(\frac{1}{x-2}\right)=\frac{\pi}{2}.$$

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$



$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$\sec^2 y \cdot y' =$$
?

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$\sec^2 y \cdot y' =$$
?

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$\sec^2 y \cdot y' = 1$$

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$sec^2 y \cdot y' = 1$$

$$y' = \frac{1}{\sec^2 y}$$

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$sec^2 y \cdot y' = 1$$

$$y' = \frac{1}{\sec^2 y}$$
$$= \frac{1}{?}$$

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$sec^2 y \cdot y' = 1$$

$$y' = \frac{1}{\sec^2 y}$$
$$= \frac{1}{1 + \tan^2 y}$$

$$\frac{\mathsf{d}}{\mathsf{d}x}(\arctan x) = \frac{1}{1+x^2}.$$

Let
$$y = \arctan x$$
.

Then
$$\tan y = x$$
.

Differentiate implicitly:
$$sec^2 y \cdot y' = 1$$

$$y' = \frac{1}{\sec^2 y}$$
$$= \frac{1}{1 + \tan^2 y}$$
$$= \frac{1}{1 + x^2}.$$

The remaining inverse trigonometric functions aren't used as often:


$$y = \operatorname{arccsc} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \operatorname{csc} y = x \quad \text{ and } \quad y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right]$$

 $y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \operatorname{sec} y = x \quad \text{ and } \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right]$
 $y = \operatorname{arccot} x \quad (|x| \in \mathbb{R}) \quad \Leftrightarrow \quad \operatorname{cot} y = x \quad \text{ and } \quad y \in \left(0, \pi\right)$

The remaining inverse trigonometric functions aren't used as often:


$$y = \operatorname{arccsc} x$$
 $(|x| \ge 1)$ \Leftrightarrow $\operatorname{csc} y = x$ and $y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right]$
 $y = \operatorname{arcsec} x$ $(|x| \ge 1)$ \Leftrightarrow $\operatorname{sec} y = x$ and $y \in \left[0, \frac{\pi}{2}\right] \cup \left[\pi, \frac{3\pi}{2}\right]$
 $y = \operatorname{arccot} x$ $(|x| \in \mathbb{R})$ \Leftrightarrow $\operatorname{cot} y = x$ and $y \in \left(0, \pi\right)$

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in ?$$




$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$

• Plot sec x.




$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$ and

 $X \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right).$


$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in ?$$



- Plot sec x.
- Restrict domain to make one-to-one: Two common choices:


$$x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$$
 and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$




- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right]$.

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$




- Plot sec x
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ is good because the domain is easiest to remember: an interval without a point. **NOT** our choice.

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$




- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ is good because the domain is easiest to remember: an interval without a point. **NOT** our choice.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ is good because $\tan x$ is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$




- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ is good because the domain is easiest to remember: an interval without a point. **NOT** our choice.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ is good because $\tan x$ is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ is good because the domain is easiest to remember: an interval without a point. **NOT** our choice.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ is good because $\tan x$ is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

$$y = \operatorname{arcsec} x \quad (|x| \ge 1) \quad \Leftrightarrow \quad \sec y = x \quad \text{ and } \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot sec x.
- Restrict domain to make one-to-one: Two common choices: $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ and $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$.
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$ is good because the domain is easiest to remember: an interval without a point. **NOT** our choice.
- $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ is good because $\tan x$ is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

Table of derivatives of inverse trigonometric functions:

$$\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1 - x^2}} \qquad \frac{d}{dx}(\arccos x) = -\frac{1}{x\sqrt{x^2 - 1}}$$

$$\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1 - x^2}} \qquad \frac{d}{dx}(\arccos x) = \frac{1}{x\sqrt{x^2 - 1}}$$

$$\frac{d}{dx}(\arctan x) = \frac{1}{1 + x^2} \qquad \frac{d}{dx}(\operatorname{arccot} x) = -\frac{1}{1 + x^2}$$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = ?$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
 $= (?)$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
 $= \left(-u^{-2}\right) \left($

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
 $= \left(-u^{-2}\right) \left(?\right)$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
 $= \left(-u^{-2}\right) \left(\frac{1}{\sqrt{1-x^2}}\right)$

Differentiate
$$y = \frac{1}{\arcsin x}$$
.
Let $u = \arcsin x$.
Then $y = u^{-1}$.
Chain Rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
 $= \left(-u^{-2}\right) \left(\frac{1}{\sqrt{1-x^2}}\right)$
 $= -\frac{1}{(\arcsin x)^2 \sqrt{1-x^2}}$.

All of the inverse trigonometric derivatives also give rise to integration formulas. These two are the most important:

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + C.$$

$$\int \frac{1}{x^2 + 1} dx = \arctan x + C.$$