

# Precalculus

## Quadratic polynomials viewed as functions

Todor Milev

2019

## Definition

Let  $a, b, c$  be real numbers with  $a \neq 0$ . The function

$$f(x) = ax^2 + bx + c$$

is called a *quadratic function*.


## Definition

Let  $a, b, c$  be real numbers with  $a \neq 0$ . The function

$$f(x) = ax^2 + bx + c$$

is called a *quadratic function*.

- The graph of a quadratic function is called a parabola.



## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1$$

## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1 = 3 \left( x^2 - ? x \right) + 1$$

## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1 = 3 \left( x^2 - \frac{5}{3}x \right) + 1$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - \color{red}{2} \cdot \frac{5}{\color{red}{2} \cdot 3} x \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + ? \quad - ? \right) + 1 \end{aligned}$$



## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \textcolor{red}{?} - \textcolor{red}{?} \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \textcolor{red}{?} \quad \quad - \textcolor{red}{?} \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - ? \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - ? \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\ &= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \end{aligned}$$



## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\ &= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\ &= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \\ &= 3 \left( x - \frac{5}{6} \right)^2 + ? \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned}3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\&= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{13}{12}.\end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c = a \left( x^2 + \frac{b}{a}x \right) + c$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c = a \left( x^2 + \frac{b}{a}x \right) + c$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned} ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\ &= a \left( x^2 + \color{red}{2} \cdot \frac{b}{\color{red}{2}a}x \right) + c \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned} ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\ &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\ &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \quad \right)^2 - \left( \quad \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \quad \right)^2 \end{array} \right. \end{aligned}$$



## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned} ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\ &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\ &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \end{array} \right. \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A + B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( \color{red}{x}^2 + 2\frac{b}{2a}\color{red}{x} + \left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 \right) + c \\
 &= a \left( \left( \color{red}{x} + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left(\frac{b}{2a}\right)^2 \\ \text{use} \\ (\color{red}{A} + B)^2 = \\ \color{red}{A}^2 + 2\color{red}{A}B + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A + B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A + B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c && \left. \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right\} \\
 &= \textcolor{red}{a} \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= \textcolor{red}{a} \left( x + \frac{b}{2a} \right)^2 - \textcolor{red}{a} \cdot \frac{b^2}{4a^2} + c
 \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right. \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 - \cancel{a} \cdot \frac{b^2}{\cancel{4a^2}} + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a}.
 \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . **To complete the square means** to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c && \left. \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right\} \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 - a \cdot \frac{b^2}{4a^2} + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a}.
 \end{aligned}$$



Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$f(x) =$$

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$f(x) = a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \quad \left| \text{complete the square} \right.$$

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \end{aligned} \quad \left| \begin{array}{l} \text{complete the square} \end{array} \right.$$

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \end{aligned} \quad \left| \begin{array}{l} \text{complete the square} \end{array} \right.$$

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + \textcolor{red}{c} - \frac{b^2}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - \textcolor{red}{4ac}}{\textcolor{red}{4a}} \end{aligned} \quad \left| \text{complete the square} \right.$$

## Definition (Discriminant of quadratic function)

The quantity  $D = b^2 - 4ac$  is called the *discriminant* of the quadratic function  $ax^2 + bx + c$ .

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} && \left| \text{complete the square} \right. \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}. \end{aligned}$$

## Definition (Discriminant of quadratic function)

The quantity  $D = b^2 - 4ac$  is called the *discriminant* of the quadratic function  $ax^2 + bx + c$ .

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} && \left| \text{complete the square} \right. \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}. \end{aligned}$$

## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .

## Definition (Discriminant of quadratic function)

The quantity  $D = b^2 - 4ac$  is called the *discriminant* of the quadratic function  $ax^2 + bx + c$ .

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality

$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} && \left| \text{complete the square} \right. \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \\ &= a \left( x - \left( -\frac{\textcolor{red}{b}}{2a} \right) \right)^2 - \frac{D}{4a}. \end{aligned}$$

## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



## Definition (Discriminant of quadratic function)

The quantity  $D = b^2 - 4ac$  is called the *discriminant* of the quadratic function  $ax^2 + bx + c$ .

Let  $a \neq 0$  and let  $f(x) = ax^2 + bx + c$ . Then we have the equality


$$\begin{aligned} f(x) &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} && \left| \begin{array}{l} \text{complete the square} \end{array} \right. \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{b^2 - 4ac}{4a} \\ &= a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}. \end{aligned}$$

## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .


## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of  $y = x^2$  is a parabola; its shape is assumed known.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of  $y = x^2$  is a parabola; its shape is assumed known.
- The standard form shows how the graph of an arbitrary quadratic is obtained from the graph of  $y = x^2$ :

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of  $y = x^2$  is a parabola; its shape is assumed known.
- The standard form shows how the graph of an arbitrary quadratic is obtained from the graph of  $y = x^2$ :
  - $ax^2$  stretches  $y = x^2$  by factor of  $a$  and possibly reflects across the  $x$  axis.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of  $y = x^2$  is a parabola; its shape is assumed known.
- The standard form shows how the graph of an arbitrary quadratic is obtained from the graph of  $y = x^2$ :
  - $ax^2$  stretches  $y = x^2$  by factor of  $a$  and possibly reflects across the  $x$  axis.
  - $a(x - h)^2$  shifts  $y = ax^2$  by  $h$  units right.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of  $y = x^2$  is a parabola; its shape is assumed known.
- The standard form shows how the graph of an arbitrary quadratic is obtained from the graph of  $y = x^2$ :
  - $ax^2$  stretches  $y = x^2$  by factor of  $a$  and possibly reflects across the  $x$  axis.
  - $a(x - h)^2$  shifts  $y = ax^2$  by  $h$  units right.
  - $a(x - h)^2 + k$  shifts  $y = a(x - h)^2 + k$  by  $k$  units up.

## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .




- The graph of a quadratic function is a parabola.



## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.

## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .




- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.



## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.
- The point  $(h, k) = \left(-\frac{b}{2a}, -\frac{D}{4a}\right)$  is called the vertex of the parabola.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.
- The point  $(h, k) = \left(-\frac{b}{2a}, -\frac{D}{4a}\right)$  is called the vertex of the parabola.
- The parabola is symmetric with respect to **the line  $x = h = -\frac{b}{2a}$** , i.e., the vertical line through its vertex.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- The graph of a quadratic function is a parabola.
- When  $a > 0$  the parabola opens upwards.
- When  $a < 0$  the parabola opens downwards.
- When  $|a|$  increases, the parabola becomes steeper.
- The point  $(h, k) = \left(-\frac{b}{2a}, -\frac{D}{4a}\right)$  is called the vertex of the parabola.
- The parabola is **symmetric** with respect to the line  $x = h = -\frac{b}{2a}$ , i.e., the vertical line through its vertex.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- When we change  $h$  and  $k$  we move the vertex of the parabola without change in steepness.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- When we change  $h$  and  $k$  we move the vertex of the parabola without change in steepness.

## Definition


The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- When we change  $h$  and  $k$  we move the vertex of the parabola without change in steepness.


## Definition

The expression  $f(x) = a(x - h)^2 + k$ , where  $h = -\frac{b}{2a}$  and  $k = -\frac{D}{4a} = -\frac{b^2 - 4ac}{4a}$  is called the standard form of  $ax^2 + bx + c$ .



- When we change  $h$  and  $k$  we move the vertex of the parabola without change in steepness.
- Therefore when we change  $b$  and  $c$  we move the vertex of the parabola without change in steepness.


## Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .



## Example




Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

Standard form

## Example




Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

Standard form

$$a(x - ?)^2 + ? = y$$

## Example



Write an equation of a parabola with **vertex at (1, 2)** that passes through the point  $(2, \frac{3}{2})$ .


$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

Standard form

Vertex at (1, 2)

## Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .


$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

Standard form

Vertex at  $(1, 2)$

## Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .


$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

Standard form

Vertex at  $(1, 2)$

## Example



Write an equation of a parabola with vertex at (1, 2) that **passes through the point (2,  $\frac{3}{2}$ )**.

$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$


$$a(\mathbf{2} - 1)^2 + 2 = \mathbf{\frac{3}{2}}$$

Standard form

Vertex at (1, 2)

**Passes through (2,  $\frac{2}{3}$ )**

## Example



Write an equation of a parabola with vertex at (1, 2) that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

$$a(\textcolor{red}{x} - 1)^2 + 2 = y$$


$$a(\textcolor{red}{2} - 1)^2 + 2 = \frac{3}{2}$$

Standard form

Vertex at (1, 2)

Passes through  $(\textcolor{red}{2}, \frac{2}{3})$

## Example



Write an equation of a parabola with vertex at (1, 2) that passes through the point (2,  $\frac{3}{2}$ ).

$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


Standard form

Vertex at (1, 2)

Passes through (2,  $\frac{3}{2}$ )



## Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


$$a = \frac{\frac{3}{2}}{2} - 2$$

Standard form

Vertex at  $(1, 2)$

Passes through  $(2, \frac{2}{3})$

# Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


$$a = \frac{3}{2} - 2$$

Standard form

Vertex at  $(1, 2)$

Passes through  $(2, \frac{2}{3})$

# Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

$$a(x - 1)^2 + 2 = y$$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


$$a = \frac{3}{2} - 2 = -\frac{1}{2}$$

Standard form

Vertex at  $(1, 2)$

Passes through  $(2, \frac{2}{3})$

# Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

Standard form

$$a(x - 1)^2 + 2 = y$$

Vertex at  $(1, 2)$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


Passes through  $(2, \frac{2}{3})$

$$a = \frac{\frac{3}{2}}{2} - 2 = -\frac{1}{2}$$

$$y = -\frac{1}{2}(x - 1)^2 + 2$$

Final answer

# Example



Write an equation of a parabola with vertex at (1, 2) that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

Standard form

$$a(x - 1)^2 + 2 = y$$

Vertex at (1, 2)

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$


Passes through  $(2, \frac{2}{3})$

$$a = \frac{\frac{3}{2}}{2} - 2 = -\frac{1}{2}$$

$$y = -\frac{1}{2}(x - 1)^2 + 2$$

Final answer

# Example



Write an equation of a parabola with vertex at  $(1, 2)$  that passes through the point  $(2, \frac{3}{2})$ .

$$a(x - h)^2 + k = y$$

Standard form

$$a(x - 1)^2 + 2 = y$$

Vertex at  $(1, 2)$

$$a(2 - 1)^2 + 2 = \frac{3}{2}$$

Passes through  $(2, \frac{2}{3})$

$$a = \frac{\frac{3}{2}}{1} - 2 = -\frac{1}{2}$$

$$y = -\frac{1}{2}(x - 1)^2 + 2$$

Final answer

$$y = -\frac{1}{2}x^2 + x + \frac{3}{2}$$

Alternative answer

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$ax^2 + bx + c = 0$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$\begin{aligned} ax^2 + bx + c &= 0 & | \text{ complete the square} \\ a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} &= 0 \end{aligned}$$



## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$\begin{array}{l|l} ax^2 + bx + c = 0 & \text{complete the square} \\ a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0 & \text{where } D = b^2 - 4ac \end{array}$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$\begin{array}{l|l} ax^2 + bx + c = 0 & \text{complete the square} \\ \hline a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0 & \text{where } D = b^2 - 4ac \\ \hline a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0 & \end{array}$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$\begin{aligned} ax^2 + bx + c &= 0 & \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right. \\ a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} &= 0 \\ a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{\textcolor{red}{D}}{4a^2} \right) &= 0 \\ a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\textcolor{red}{\sqrt{D}}}{2a} \right)^2 \right) &= 0 \end{aligned}$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$\begin{array}{lcl}
 ax^2 + bx + c & = & 0 \\
 a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} & = & 0 \\
 a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) & = & 0 \\
 a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) & = & 0
 \end{array}
 \quad \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right.$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$ax^2 + bx + c = 0$$

*complete the square*

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

*where  $D = b^2 - 4ac$*

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0$$

*use  $A^2 - B^2$   
 $= (A - B)(A + B)$*

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$ax^2 + bx + c = 0$$

*complete the square*

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

*where  $D = b^2 - 4ac$*

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0$$

*use  $A^2 - B^2 = (A - B)(A + B)$*

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$ax^2 + bx + c = 0$$

*complete the square*

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

*where  $D = b^2 - 4ac$*

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0$$

*use  $A^2 - B^2 = (A - B)(A + B)$*

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0 \quad \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right.$$

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0 \quad \left| \begin{array}{l} \text{use } A^2 - B^2 \\ = (A - B)(A + B) \end{array} \right.$$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$



## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0 \quad \left| \begin{array}{l} \text{complete the square} \end{array} \right.$$

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0 \quad \left| \begin{array}{l} \text{where } D = b^2 - 4ac \end{array} \right.$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0 \quad \left| \begin{array}{l} \text{use } A^2 - B^2 \\ = (A - B)(A + B) \end{array} \right.$$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0 \quad \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right.$$

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0 \quad \left| \begin{array}{l} \text{use } A^2 - B^2 \\ = (A - B)(A + B) \end{array} \right.$$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

$$x = \frac{-b \pm \sqrt{D}}{2a}$$

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0 \quad \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right.$$

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0 \quad \left| \begin{array}{l} \text{use } A^2 - B^2 \\ = (A - B)(A + B) \end{array} \right.$$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

$$x = \frac{-b + \sqrt{D}}{2a}$$

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0 \quad \left| \begin{array}{l} \text{complete the square} \\ \text{where } D = b^2 - 4ac \end{array} \right.$$

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0 \quad \left| \begin{array}{l} \text{use } A^2 - B^2 \\ = (A - B)(A + B) \end{array} \right.$$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{D}}{2a}.$$

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0$$

complete the square

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

where  $D = b^2 - 4ac$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0$$

use  $A^2 - B^2$   
 $= (A - B)(A + B)$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{D}}{2a}$$

## Problem (Quadratic equation formula)

Solve the general quadratic equation

$$ax^2 + bx + c = 0$$

complete the square

$$a \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a} = 0$$

where  $D = b^2 - 4ac$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{D}{4a^2} \right) = 0$$

$$a \left( \left( x + \frac{b}{2a} \right)^2 - \left( \frac{\sqrt{D}}{2a} \right)^2 \right) = 0$$

$$a \left( x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} \right) \left( x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} \right) = 0$$

use  $A^2 - B^2 = (A - B)(A + B)$

$$x + \frac{b}{2a} - \frac{\sqrt{D}}{2a} = 0 \quad \text{or} \quad x + \frac{b}{2a} + \frac{\sqrt{D}}{2a} = 0$$

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{D}}{2a}$$

## Problem (Quadratic equation formula)

*Solve the general quadratic equation*

$$ax^2 + bx + c = 0$$

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or}$$

$$x = \frac{-b - \sqrt{D}}{2a}.$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{D}}{2a}$$



## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$*

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$*

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



## Theorem

*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$


*are given by:*

$$x = x_1 = \frac{-b + \sqrt{D}}{2a} \quad \text{or} \quad x = x_2 = \frac{-b - \sqrt{D}}{2a},$$

*where  $D = b^2 - 4ac$ , or equivalently by:*

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

# Theorem



*The solutions of the quadratic equation*


$$ax^2 + bx + c = 0$$

*are the numbers*

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}.$$

# Theorem



*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are the numbers*

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$


$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}.$$

- Abbreviated as

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{D}}{2a},$$

where  $D = b^2 - 4ac$ .

# Theorem



*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are the numbers*

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$


$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}.$$

- Abbreviated as

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{D}}{2a}, \quad \text{where } D = b^2 - 4ac.$$

- If  $D < 0$  then  $\sqrt{D}$  is not a real  $\Rightarrow$  quadratic has no real solutions.

# Theorem



*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are the numbers*

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$


$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}.$$

- Abbreviated as

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{D}}{2a}, \quad \text{where } D = b^2 - 4ac.$$

- If  $D < 0$  then  $\sqrt{D}$  is not a real  $\Rightarrow$  quadratic has no real solutions.
- If  $D = 0$  then  $x_1 = x_2$ , the equation has only one zero (with multiplicity two).

# Theorem



*The solutions of the quadratic equation*

$$ax^2 + bx + c = 0$$

*are the numbers*

$$x_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$


$$x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}.$$

- Abbreviated as

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{D}}{2a}, \quad \text{where } D = b^2 - 4ac.$$


- If  $D < 0$  then  $\sqrt{D}$  is not a real  $\Rightarrow$  quadratic has no real solutions.
- If  $D = 0$  then  $x_1 = x_2$ , the equation has only one zero (with multiplicity two). The zero is located at the vertex of the parabola.

## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

## Example




Find the x-intercepts of  $x^2 - 4x + 3$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$




## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}\end{aligned}$$


## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}\end{aligned}$$


## Example



Find the x-intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}\end{aligned}$$


## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\&= \frac{4 \pm \sqrt{4}}{2}\end{aligned}$$


## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\&= \frac{4 \pm \sqrt{4}}{2}\end{aligned}$$


## Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\&= \frac{4 \pm \sqrt{4}}{2} \\&= \frac{4 \pm 2}{2}\end{aligned}$$


# Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \left\{ \begin{array}{l} \frac{4 + 2}{2} \\ \frac{4 - 2}{2} \end{array} \right.
 \end{aligned}$$

# Example




Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \begin{cases} \frac{4+2}{2} = \frac{6}{2} \\ \frac{4-2}{2} \end{cases}
 \end{aligned}$$




# Example



Find the x-intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \begin{cases} \frac{4+2}{2} = \frac{6}{2} = 3 \\ \frac{4-2}{2} \end{cases}
 \end{aligned}$$


# Example



Find the x-intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \begin{cases} \frac{4+2}{2} = \frac{6}{2} = 3 \\ \frac{4-2}{2} = \frac{2}{2} = 1 \end{cases}
 \end{aligned}$$


# Example



Find the  $x$ -intercepts of  $x^2 - 4x + 3$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \begin{cases} \frac{4+2}{2} = \frac{6}{2} = 3 \\ \frac{4-2}{2} = \frac{2}{2} = 1 \end{cases}
 \end{aligned}$$


# Example



Find the x-intercepts of  $x^2 - 4x + 3$ .


$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1} \\
 &= \frac{4 \pm \sqrt{4}}{2} \\
 &= \frac{4 \pm 2}{2} \\
 &= \begin{cases} \frac{4+2}{2} = \frac{6}{2} = 3 \\ \frac{4-2}{2} = \frac{2}{2} = 1 \end{cases}
 \end{aligned}$$

## Example



Find the  $x$ -intercepts of  $\frac{x^2}{2} - x - 1$ .


## Example



Find the x-intercepts of  $\frac{x^2}{2} - x - 1$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


## Example



Find the x-intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}}
 \end{aligned}$$

## Example




Find the  $x$ -intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}}\end{aligned}$$




## Example



Find the  $x$ -intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}}\end{aligned}$$


## Example



Find the x-intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}}
 \end{aligned}$$


## Example



Find the  $x$ -intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}} \\
 &= 1 \pm \sqrt{3}
 \end{aligned}$$


## Example



Find the x-intercepts of  $\frac{x^2}{2} - x - 1$ .

$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}} \\
 &= 1 \pm \sqrt{3}
 \end{aligned}$$


## Example



Find the x-intercepts of  $\frac{x^2}{2} - x - 1$ .


$$\begin{aligned}
 x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\
 &= \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot \frac{1}{2} \cdot (-1)}}{2 \cdot \frac{1}{2}} \\
 &= 1 \pm \sqrt{3}
 \end{aligned}$$

## Example



Find the  $x$ -intercepts of  $x^2 - 2x + 3$ .


## Example



Find the  $x$ -intercepts of  $x^2 - 2x + 3$ .

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Example




Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1}\end{aligned}$$




## Example



Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1}\end{aligned}$$


## Example



Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1}\end{aligned}$$


## Example



Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1} \\&= \frac{2 \pm \sqrt{-8}}{2}\end{aligned}$$


## Example



Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1} \\&= \frac{2 \pm \sqrt{-8}}{2}\end{aligned}$$


## Example



Find the x-intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1} \\&= \frac{2 \pm \sqrt{-8}}{2} \\&\text{no real solutions}\end{aligned}$$

## Example



Find the  $x$ -intercepts of  $x^2 - 2x + 3$ .

$$\begin{aligned}x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-(-2) \pm \sqrt{(-2)^2 - 4 \cdot 1 \cdot (3)}}{2 \cdot 1} \\&= \frac{2 \pm \sqrt{-8}}{2} \\&\quad \text{no real solutions}\end{aligned}$$

**no  $x$  - intercepts**

## Proposition

*Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2 (x_1 - x_2)^2$ .*

## Proof.



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$a^2(x_1 - x_2)^2 = a^2 \left( \frac{-b + \sqrt{D}}{2a} - \frac{-b - \sqrt{D}}{2a} \right)$$





## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$a^2(x_1 - x_2)^2 = a^2 \left( \frac{-b + \sqrt{D}}{2a} - \frac{-b - \sqrt{D}}{2a} \right)$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$a^2(x_1 - x_2)^2 = a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right)$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned} a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\ &= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \end{aligned}$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned} a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\ &= a^2 \left( \frac{\cancel{2}\sqrt{D}}{\cancel{2}a} \right)^2 \end{aligned}$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned} a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\ &= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \\ &= a^2 \frac{D}{a^2} \end{aligned}$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned} a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\ &= a^2 \left( \frac{2\sqrt{D}}{2\cancel{a}} \right)^2 \\ &= a^2 \frac{D}{\cancel{a}^2} \end{aligned}$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned}a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\&= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \\&= \cancel{a^2} \frac{D}{\cancel{a^2}} \\&= D\end{aligned}$$



## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned}
 a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\
 &= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \\
 &= \cancel{a^2} \frac{D}{\cancel{a^2}} \\
 &= D, \quad \text{as desired.}
 \end{aligned}$$





## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2 (x_1 - x_2)^2$ .

## Proof.

$$\begin{aligned}
 a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\
 &= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \\
 &= \cancel{a^2} \frac{D}{\cancel{a^2}} \\
 &= D, \quad \text{as desired.}
 \end{aligned}$$



- Discriminant is zero  $\Leftrightarrow$  the quadratic has non-distinct roots

## Proposition

Let  $ax^2 + bx + c$ ,  $a \neq 0$  be a quadratic with discriminant  $D = b^2 - 4ac$  and roots  $x_1$  and  $x_2$ . Then  $D = a^2(x_1 - x_2)^2$ .

## Proof.


$$\begin{aligned}
 a^2(x_1 - x_2)^2 &= a^2 \left( \frac{\cancel{b} + \sqrt{D}}{2a} - \frac{\cancel{b} - \sqrt{D}}{2a} \right) \\
 &= a^2 \left( \frac{2\sqrt{D}}{2a} \right)^2 \\
 &= \cancel{a^2} \frac{D}{\cancel{a^2}} \\
 &= D, \quad \text{as desired.}
 \end{aligned}$$



- Discriminant is zero  $\Leftrightarrow$  the quadratic has non-distinct roots, hence the discriminant discriminates between the two roots.

## Example


Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.


- Quadratic roots:  $x_1, x_2 = ?$  .



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .




## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when  $b^2 - 4ac > 0$




## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when  $b^2 - 4ac > 0$




## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when
$$b^2 - 4ac > 0$$
$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$






## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when
$$b^2 - 4ac > 0$$
$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$




## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when
$$b^2 - 4ac > 0$$
$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.


- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.


- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when


$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when


$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when


$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12}) (k + \sqrt{12}) > 0$$





## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when


$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when



$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when



$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when



$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when


$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$


$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in ?$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$

$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$





## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$

$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$



## Example

Find the values of the parameter  $k$  for which the equation  $3x^2 - kx + 1$  has two real distinct roots.

- Quadratic roots:  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

- $\Rightarrow$  The roots  $x_1, x_2$  are real and distinct when

$$b^2 - 4ac > 0$$



$$(-k)^2 - 4 \cdot 3 \cdot 1 > 0$$

$$k^2 - 12 > 0$$


$$k^2 - \sqrt{12}^2 > 0$$

$$(k - \sqrt{12})(k + \sqrt{12}) > 0$$

$$k \in (-\infty, -\sqrt{12}) \cup (\sqrt{12}, \infty)$$




To plot a parabola by hand roughly, we need to do the following.



To plot a parabola by hand roughly, we need to do the following.


- Find the vertex of the parabola.






To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.




To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.
- Find the  $x$  intercept(s) if any.




To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.
- Find the  $x$  intercept(s) if any.
- Select (or re-select) axes scale so all important points found in the preceding items fit in the plot.




To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.
- Find the  $x$  intercept(s) if any.
- Select (or re-select) axes scale so all important points found in the preceding items fit in the plot.
- Plot the parabola freehand, making sure that the parabola passes through all special points you found in the preceding items.




To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.
- Find the  $x$  intercept(s) if any.
- Select (or re-select) axes scale so all important points found in the preceding items fit in the plot.
- Plot the parabola freehand, making sure that the parabola passes through all special points you found in the preceding items.
- If  $a > 0$  your parabola should open upwards, if  $a < 0$  your parabola should open downwards.




To plot a parabola by hand roughly, we need to do the following.

- Find the vertex of the parabola.
- Find the  $y$  intercept.
- Find the  $x$  intercept(s) if any.
- Select (or re-select) axes scale so all important points found in the preceding items fit in the plot.
- Plot the parabola freehand, making sure that the parabola passes through all special points you found in the preceding items.
- If  $a > 0$  your parabola should open upwards, if  $a < 0$  your parabola should open downwards.
- For  $|a| > 1$  we should aim to draw the graph steeper than  $a = x^2$ , for  $|a| < 1$  we should aim to draw the graph flatter than  $a = x^2$ .




## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The vertex of the parabola is given by:

$$x = ?$$

$$y = ?$$



## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2(-\frac{2}{3})} = \frac{21}{4}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$y = ?$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$y = ?$$

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$y = f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a}$$



## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} \end{aligned}$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} \end{aligned}$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} \end{aligned}$$

# Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$

# Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$

# Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right) \cancel{3}}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right) \cancel{3}}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$



## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right) \cancel{3}}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \end{aligned}$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right) \cancel{3}}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} \end{aligned}$$

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} \end{aligned}$$

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2\left(-\frac{2}{3}\right)} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} = \frac{171}{8}. \end{aligned}$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2(-\frac{2}{3})} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} = \frac{171}{8}. \end{aligned}$$

## Example



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2(-\frac{2}{3})} = \frac{21}{4}$$

$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4\left(-\frac{2}{3}\right)3}{4\left(-\frac{2}{3}\right)} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} = \frac{171}{8}. \end{aligned}$$

- The y-intercept is  $f(0) = ?$ .

## Example



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The vertex of the parabola is given by:

$$x = -\frac{b}{2a} = -\frac{7}{2(-\frac{2}{3})} = \frac{21}{4}$$


$$\begin{aligned} y &= f\left(-\frac{b}{2a}\right) = -\frac{D}{4a} = -\frac{(b^2 - 4ac)}{4a} \\ &= -\frac{7^2 - 4(-\frac{2}{3})}{4(-\frac{2}{3})} = \frac{49 + 8}{\frac{8}{3}} \\ &= \frac{3 \cdot 57}{8} = \frac{171}{8} \end{aligned}$$

- The y-intercept is  $f(0) = 3$ .

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of  
$$-\frac{2}{3}x^2 + 7x + 3 = 0$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$




## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$x = ?$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$


$$-2x^2 + 21x + 9 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \end{aligned}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$



## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$


$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \end{aligned}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$



## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)}$$

$$= \frac{-21 \pm \sqrt{441 + 72}}{-4}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \end{aligned}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$





## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \end{aligned}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$



## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \mp \sqrt{513}}{4} \end{aligned}$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \pm \sqrt{513}}{4} \end{aligned}$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$

$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \pm \sqrt{513}}{4} \\ &= \frac{21 \pm \sqrt{9 \cdot 57}}{4} \end{aligned}$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$-\frac{2}{3}x^2 + 7x + 3 = 0 \quad | \cdot 3$$


$$-2x^2 + 21x + 9 = 0$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \pm \sqrt{513}}{4} \\ &= \frac{21 \pm \sqrt{9 \cdot 57}}{4} \\ &= \frac{21 \pm \sqrt{9} \sqrt{57}}{4} \end{aligned}$$



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$

## Example



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3.$

- The  $x$  intercepts are given by the solutions of

$$\begin{aligned} -\frac{2}{3}x^2 + 7x + 3 &= 0 & | \cdot 3 \\ -2x^2 + 21x + 9 &= 0 \end{aligned}$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \pm \sqrt{513}}{4} \\ &= \frac{21 \pm \sqrt{9 \cdot 57}}{4} \\ &= \frac{21 \pm \sqrt{9} \sqrt{57}}{4} \\ &= \frac{21 \pm 3\sqrt{57}}{4} \end{aligned}$$

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- The  $x$  intercepts are given by the solutions of

$$\begin{aligned} -\frac{2}{3}x^2 + 7x + 3 &= 0 & | \cdot 3 \\ -2x^2 + 21x + 9 &= 0 \end{aligned}$$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-21 \pm \sqrt{21^2 - 4 \cdot (-2) \cdot 9}}{2 \cdot (-2)} \\ &= \frac{-21 \pm \sqrt{441 + 72}}{-4} \\ &= \frac{21 \pm \sqrt{513}}{4} \\ &= \frac{21 \pm \sqrt{9 \cdot 57}}{4} \\ &= \frac{21 \pm \sqrt{9} \sqrt{57}}{4} \\ &= \frac{21 \pm 3\sqrt{57}}{4} \end{aligned}$$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$   
 $x$ -intercepts at


$$x = \frac{21 - 3\sqrt{57}}{4},$$

$$x = \frac{21 + 3\sqrt{57}}{4}.$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4}$




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
y-intercept at  $y = 3$   
x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$



## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
y-intercept at  $y = 3$   
x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers ?




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
y-intercept at  $y = 3$   
x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .


- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers 21 and 22.



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
y-intercept at  $y = 3$   
x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .




- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4}$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .




- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4}$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4}$

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at


$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$



## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4}$


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$   
 $x$ -intercepts at

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



● Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
which is close to  $-1$ .


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example


Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 $y$ -intercept at  $y = 3$   
 $x$ -intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
  - $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1$ .
  - The parabola vertex is less than 22 units high and the parabola opens downwards.

## Example




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3.$

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5.$
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11.$
  - $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1.$
  - The parabola vertex is less than 22 units high and the parabola opens downwards.
  - Axes height of 22 units appears reasonable.

## Example




Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1$ .
- The parabola vertex is less than 22 units high and the parabola opens downwards.
- Axes height of 22 units appears reasonable.
- A grid of width 3 units appears reasonable.

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

## Example




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
**y-intercept at  $y = 3$**   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3.$

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5.$
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11.$
  - $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1.$
  - The parabola vertex is less than 22 units high and the parabola opens downwards.
  - Axes height of 22 units appears reasonable.
  - A grid of width 3 units appears reasonable.
  - Plot all relevant points.

## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1$ .
- The parabola vertex is less than 22 units high and the parabola opens downwards.
- Axes height of 22 units appears reasonable.
- A grid of width 3 units appears reasonable.
- Plot all relevant points.

Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$


**x-intercepts at**

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$



## Example



Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:

- $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
- $\frac{171}{8}$  is between the integers 21 and 22.
- $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$   
 which is close to  $\frac{44}{4} = 11$ .
- $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$   
 which is close to  $-1$ .
- The parabola vertex is less than 22 units high and the parabola opens downwards.
- Axes height of 22 units appears reasonable.
- A grid of width 3 units appears reasonable.
- Plot all relevant points.


Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$

**x-intercepts at**

$$x = \frac{21-3\sqrt{57}}{4},$$

$$x = \frac{21+3\sqrt{57}}{4}.$$

## Example




Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4}$ ,  
 $x = \frac{21+3\sqrt{57}}{4}$ .

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3$ .

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5$ .
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$  which is close to  $\frac{44}{4} = 11$ .
  - $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$  which is close to  $-1$ .
  - The parabola vertex is less than 22 units high and the parabola opens downwards.
  - Axes height of 22 units appears reasonable.
  - A grid of width 3 units appears reasonable.
  - Plot all relevant points.
  - Finally “connect the dots with a freehand drawing”.

## Example



Vertex at:  $(\frac{21}{4}, \frac{171}{8})$   
 y-intercept at  $y = 3$   
 x-intercepts at  
 $x = \frac{21-3\sqrt{57}}{4},$   
 $x = \frac{21+3\sqrt{57}}{4}.$

Plot roughly by hand the graph of  
 $f(x) = -\frac{2}{3}x^2 + 7x + 3.$

- Select scale to fit the picture:
  - $\frac{21}{4}$  is close to  $\frac{20}{4} = 5.$
  - $\frac{171}{8}$  is between the integers 21 and 22.
  - $\frac{21+3\sqrt{57}}{4}$  is close to  $\frac{21+3\sqrt{64}}{4} = \frac{21+24}{4} = \frac{45}{4}$  which is close to  $\frac{44}{4} = 11.$
  - $\frac{21-3\sqrt{57}}{4}$  is close to  $\frac{21-3\sqrt{64}}{4} = \frac{21-24}{4} = -\frac{3}{4}$  which is close to  $-1.$
  - The parabola vertex is less than 22 units high and the parabola opens downwards.
  - Axes height of 22 units appears reasonable.
  - A grid of width 3 units appears reasonable.
  - Plot all relevant points.
  - Finally “connect the dots with a freehand drawing”.

## Example

Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

## Example

Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.


## Example

Find the values of the parameter  $k$  for which

$x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive


## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up

## Example




Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it




## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.


## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = ?$


## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

## Example




Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  **the quadratic has no real solutions.**
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

# Example




Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

## Example




Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

## Example




Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$k^2 + 2k + 1 - 8k < 0$$



## Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .


- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$k^2 + 2k + 1 - 8k < 0$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$


$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = ?$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$k^2 + 2k + 1 - 8k < 0$$


$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$k^2 + 2k + 1 - 8k < 0$$

$$k^2 - 6k + 1 < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$\begin{aligned} k_1, k_2 &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2} \\ &= \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} \end{aligned}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$\begin{aligned} k_1, k_2 &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2} \\ &= \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} \end{aligned}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$\begin{aligned} k_1, k_2 &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2} \\ &= \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} \end{aligned}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$


$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$\begin{aligned} k_1, k_2 &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2} \\ &= \frac{2 \cdot 3 \pm \sqrt{4}\sqrt{8}}{2} = \frac{\cancel{2} (3 \pm \sqrt{8})}{\cancel{2}} = 3 \pm \sqrt{8} \end{aligned}$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$\begin{aligned}
 k_1, k_2 &= \frac{-(-6) \pm \sqrt{(-6)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{6 \pm \sqrt{32}}{2} \\
 &= \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{\cancel{2} (3 \pm \sqrt{8})}{\cancel{2}} = 3 \pm \sqrt{8}
 \end{aligned}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{\cancel{2} (3 \pm \sqrt{8})}{\cancel{2}} = 3 \pm \sqrt{8}$$

# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in ?$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in ?$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in ?$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .


$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in ?$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in (k_1, k_2) = (3 - \sqrt{8}, 3 + \sqrt{8})$$



# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$

$$(k - k_1)(k - k_2) < 0$$


$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in (k_1, k_2) = (3 - \sqrt{8}, 3 + \sqrt{8})$$





# Example



Find the values of the parameter  $k$  for which  $x^2 + (k + 1)x + 2k > 0$  holds for all real  $x$ .

- In order for the quadratic to be positive, its graph must lie entirely above the  $x$  axis.
- Leading coefficient is positive  $\Rightarrow$  graph opens up  $\Rightarrow$  is above  $x$  axis if it does not intersect it  $\Rightarrow$  the quadratic has no real solutions.
- The roots of a quadratic are  $x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ .

$$b^2 - 4ac < 0$$

$$(k + 1)^2 - 4 \cdot 1 \cdot 2k < 0$$


$$(k - k_1)(k - k_2) < 0$$

$$k_1, k_2 = \frac{2 \cdot 3 \pm \sqrt{4} \sqrt{8}}{2} = \frac{2(3 \pm \sqrt{8})}{2} = 3 \pm \sqrt{8}$$

$$k \in (k_1, k_2) = (3 - \sqrt{8}, 3 + \sqrt{8})$$




## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$3x^2 + 2x + 1 = 3 \left( x^2 + 2 \cdot \frac{1}{3}x + ? - ? \right) + 1$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$3x^2 + 2x + 1 = 3 \left( x^2 + 2 \cdot \frac{1}{3}x + ? - ? \right) + 1$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$3x^2 + 2x + 1 = 3 \left( x^2 + 2 \cdot \frac{1}{3}x + ? - ? \right) + 1$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$3x^2 + 2x + 1 = 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned} 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\ &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \end{aligned}$$

## Example




Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned} 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\ &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \\ &= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1 \end{aligned}$$




## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned} 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\ &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \\ &= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1 \\ &= 3 \left( x - \left( -\frac{1}{3} \right) \right)^2 + ? \end{aligned}$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned} 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\ &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \\ &= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1 \\ &= 3 \left( x - \left( -\frac{1}{3} \right) \right)^2 + ? \end{aligned}$$


## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned} 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\ &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \\ &= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1 \\ &= 3 \left( x - \left( -\frac{1}{3} \right) \right)^2 + \frac{2}{3} \end{aligned}$$

# Example




Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$\begin{aligned}
 3x^2 + 2x + 1 &= 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1 \\
 &= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1 \\
 &= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1 \\
 &= 3 \left( x - \left( -\frac{1}{3} \right) \right)^2 + \frac{2}{3}
 \end{aligned}$$

Minimum point =  $(?, ?)$

## Example



Find the minimum point on the curve  
 $y = 3x^2 + 2x + 1$  by completing the square.

$$3x^2 + 2x + 1 = 3 \left( x^2 + 2 \cdot \frac{1}{3}x + \frac{1}{9} - \frac{1}{9} \right) + 1$$

$$= 3 \left( \left( x + \frac{1}{3} \right)^2 - \frac{1}{9} \right) + 1$$

$$= 3 \left( x + \frac{1}{3} \right)^2 - \frac{1}{3} + 1$$

$$= 3 \left( x - \left( -\frac{1}{3} \right) \right)^2 + \frac{2}{3}$$

$$\text{Minimum point} = \left( -\frac{1}{3}, \frac{2}{3} \right)$$

# Maximum or minimum value of a quadratic function

- Let  $f(x) = ax^2 + bx + c$  - quadratic ( $a \neq 0$ ).
- Let  $D$  be the discriminant  $D = b^2 - 4ac$ .

$$f(x) = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a} \quad \left| \text{complete the square} \right.$$

# Maximum or minimum value of a quadratic function

- Let  $f(x) = ax^2 + bx + c$  - quadratic ( $a \neq 0$ ).
- Let  $D$  be the discriminant  $D = b^2 - 4ac$ .

$$f(x) = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a} \quad \left| \text{complete the square} \right.$$

- Therefore if  $a > 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a}$  .

# Maximum or minimum value of a quadratic function

- Let  $f(x) = ax^2 + bx + c$  - quadratic ( $a \neq 0$ ).
- Let  $D$  be the discriminant  $D = b^2 - 4ac$ .

$$f(x) = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a} \quad \left| \text{complete the square} \right.$$

- Therefore if  $a > 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a} \geq -\frac{D}{4a}$ .



# Maximum or minimum value of a quadratic function

- Let  $f(x) = ax^2 + bx + c$  - quadratic ( $a \neq 0$ ).
- Let  $D$  be the discriminant  $D = b^2 - 4ac$ .

$$f(x) = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a} \quad \left| \text{complete the square} \right.$$

- Therefore if  $a > 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a} \geq -\frac{D}{4a}$ .
- Similarly if  $a < 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a} \leq -\frac{D}{4a}$ .

# Maximum or minimum value of a quadratic function

- Let  $f(x) = ax^2 + bx + c$  - quadratic ( $a \neq 0$ ).
- Let  $D$  be the discriminant  $D = b^2 - 4ac$ .

$$f(x) = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a} \quad \left| \text{complete the square} \right.$$

- Therefore if  $a > 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a} \geq -\frac{D}{4a}$ .
- Similarly if  $a < 0$  then  $f(x) = a(\text{square}) - \frac{D}{4a} \leq -\frac{D}{4a}$ .

$$\text{Recall } f(x) = ax^2 + bx + c = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}.$$

## Proposition

Let  $f(x) = ax^2 + bx + c$ ,  $a \neq 0$  and let  $D = b^2 - 4ac$ .


- If  $a > 0$  then  $f(x)$  has no maximum and has minimum at  $x = -\frac{b}{2a}$ .
- If  $a < 0$  then  $f(x)$  has no minimum and has maximum at  $x = -\frac{b}{2a}$ .
- In both cases, the extremal value (either maximum or minimum) is  $f\left(-\frac{b}{2a}\right) = -\frac{b^2-4ac}{4a} = -\frac{D}{4a}$ .

$$\text{Recall } f(x) = ax^2 + bx + c = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}.$$

## Proposition

Let  $f(x) = ax^2 + bx + c$ ,  $a \neq 0$  and let  $D = b^2 - 4ac$ .

- If  $a > 0$  then  $f(x)$  has no maximum and has minimum at  $x = -\frac{b}{2a}$ .
- If  $a < 0$  then  $f(x)$  has no minimum and has maximum at  $x = -\frac{b}{2a}$ .
- In both cases, the extremal value (either maximum or minimum) is  $f\left(-\frac{b}{2a}\right) = -\frac{b^2-4ac}{4a} = -\frac{D}{4a}$ .




$$\text{Recall } f(x) = ax^2 + bx + c = a \left( x - \left( -\frac{b}{2a} \right) \right)^2 - \frac{D}{4a}.$$

## Proposition

Let  $f(x) = ax^2 + bx + c$ ,  $a \neq 0$  and let  $D = b^2 - 4ac$ .

- If  $a > 0$  then  $f(x)$  has no maximum and has minimum at  $x = -\frac{b}{2a}$ .
- If  $a < 0$  then  $f(x)$  has no minimum and has maximum at  $x = -\frac{b}{2a}$ .
- In both cases, the extremal value (either maximum or minimum) is  $f\left(-\frac{b}{2a}\right) = -\frac{b^2-4ac}{4a} = -\frac{D}{4a}$ .



## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

Maximizing:

$xz$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

Maximizing:

$$xz$$



## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$xz$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$xz = x(12 - x)$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.

$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$x = ?$$



## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$x = -\frac{b}{2a}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \\ z &= 12 - x = 12 - 6 \end{aligned}$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

Max. product  $= xz$



## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

$$\text{Max. product} = xz = 6 \cdot 6$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

$$\text{Max. product} = xz = 6 \cdot 6$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

$$\text{Max. product} = xz = 6 \cdot 6 = 36.$$

## Example

Let  $x, z$  be two numbers that add to 12. Choose  $x$  and  $z$  so that the product  $x \cdot z$  is maximal.



$$x + z = 12$$

$$z = 12 - x$$

Maximizing:

$$\begin{aligned} xz &= x(12 - x) \\ &= -x^2 + 12x \end{aligned}$$

Parabola opens down  $\Rightarrow$  has maximum, attained at:

$$\begin{aligned} x &= -\frac{b}{2a} \\ &= -\frac{12}{-2} = 6 \end{aligned}$$

$$z = 12 - x = 12 - 6 = 6$$

$$\text{Max. product} = xz = 6 \cdot 6 = 36.$$