

# Precalculus

## Trig cofunction identities and angle-sum formulas

Todor Milev

2019

# Outline

## 1 Cofunction identities

# Outline

- 1 Cofunction identities
- 2 Trigonometric Functions of Sums of Angles

# Outline

- 1 Cofunction identities
- 2 Trigonometric Functions of Sums of Angles
- 3 Double Angle Formulas

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.



# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{array}{llll} \sin\left(\frac{\pi}{2} - \alpha\right) & = & \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) & = & \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) & = & \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) & = & -\sin \alpha \end{array}$$

- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.
- The Quadrant I part of the proof serves as a visual aid for memorization.

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$


- The proof each formula is broken into 4 cases depending on which quadrant contains  $\alpha$ .
- This makes a total of 4 formulas  $\times$  4 cases per formula = 16 cases.
- We show only a few of the cases.
- The proof provides intuition why the formulas are true.
- The Quadrant I part of the proof serves as a visual aid for memorization.
- There is an algebraically simpler (but theoretically advanced) way to prove the above identities through the angle sum formulas, derived in turn from Euler's formula (studied later/in another course).

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = \frac{|PQ|}{|OP|}$$






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \quad \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \quad \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in quadrant I.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} & \left| \square ORPQ \right. \\ &= \frac{|OR|}{|OP|} \\ &= \cos \alpha & \left. \text{as desired} \right.\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.


$$\sin\left(\frac{\pi}{2} - \alpha\right) = -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square \text{ } OR'P'Q'$$


# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|}\end{aligned}$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We are showing  $\sin\left(\frac{\pi}{2} - \alpha\right) = \cos \alpha$  when  $\alpha$  is in Quadrant III. It follows  $\frac{\pi}{2} - \alpha$  is in Quadrant III.

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= -\frac{|P'R'|}{|OP'|} = -\frac{|OQ'|}{|OP'|} \quad \square OR'P'Q' \\ &= -\frac{|OR|}{|OP|} \\ &= \cos \alpha\end{aligned}$$

as desired




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) =$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\cos\left(\frac{\pi}{2} + \alpha\right) = -\frac{|OR'|}{|OP'|}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \Big| \quad \square ORPQ \\ &= -\frac{|P'Q'|}{|OP'|}\end{aligned}$$






# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \left| \square ORPQ \right. \\ &= -\frac{|P'Q'|}{|OP'|}\end{aligned}$$




# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \square ORPQ \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|}\end{aligned}$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.

$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} \quad \square ORPQ \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|} \\ &= -\sin \alpha.\end{aligned}$$


□

# Cofunction identities

## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

## Part of Proof.



We show  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$  when  $\alpha$  is in Quadrant I. It follows  $\frac{\pi}{2} + \alpha$  is in Quadrant II.


$$\begin{aligned}\cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|OR'|}{|OP'|} && \square ORPQ \\ &= -\frac{|P'Q'|}{|OP'|} \\ &= -\frac{|PR|}{|OP|} \\ &= -\sin \alpha. && \text{as desired } \square\end{aligned}$$

# Cofunction identities


## Proposition (Cofunction identities)

$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \cos \alpha & \sin\left(\frac{\pi}{2} + \alpha\right) &= \cos \alpha \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \sin \alpha & \cos\left(\frac{\pi}{2} + \alpha\right) &= -\sin \alpha\end{aligned}$$

To memorize the cofunction identities it suffices to memorize the Quadrant I case via the two diagrams below.




$$\begin{aligned}\sin\left(\frac{\pi}{2} - \alpha\right) &= \frac{|PQ|}{|OP|} \\ \cos\left(\frac{\pi}{2} - \alpha\right) &= \frac{|OQ|}{|OP|}\end{aligned}$$



$$\begin{aligned}\sin\left(\frac{\pi}{2} + \alpha\right) &= \frac{|OQ'|}{|OP|} \\ \cos\left(\frac{\pi}{2} + \alpha\right) &= -\frac{|PQ'|}{|OP|} = \frac{|PR|}{|OP|}\end{aligned}$$

## Definition (Similar triangles)


We say that a triangle  $\triangle ABC$  is similar to a triangle  $\triangle A'B'C'$  if the two triangles have equal angles.



## Definition (Similar triangles)

We say that a triangle  $\triangle ABC$  is similar to a triangle  $\triangle A'B'C'$  if the two triangles have equal angles.

- The equal angles are assumed given in the same order for both triangles, that is,  $\angle ABC = \angle A'B'C'$ ,  $\angle BCA = \angle B'C'A'$ ,  $\angle CAB = \angle C'A'B'$ .






## Definition (Similar triangles)

We say that a triangle  $\triangle ABC$  is similar to a triangle  $\triangle A'B'C'$  if the two triangles have equal angles.


- The equal angles are assumed given in the same order for both triangles, that is,  $\angle ABC = \angle A'B'C'$ ,  $\angle BCA = \angle B'C'A'$ ,  $\angle CAB = \angle C'A'B'$ .



## Definition (Similar triangles)

We say that a triangle  $\triangle ABC$  is similar to a triangle  $\triangle A'B'C'$  if the two triangles have equal angles.


- The equal angles are assumed given in the same order for both triangles, that is,  $\angle ABC = \angle A'B'C'$ ,  $\angle BCA = \angle B'C'A'$ ,  $\angle CAB = \angle C'A'B'$ .



## Definition (Similar triangles)

We say that a triangle  $\triangle ABC$  is similar to a triangle  $\triangle A'B'C'$  if the two triangles have equal angles.

- The equal angles are assumed given in the same order for both triangles, that is,  $\angle ABC = \angle A'B'C'$ ,  $\angle BCA = \angle B'C'A'$ ,  $\angle CAB = \angle C'A'B'$ .




The following statement is proved in the subject of Euclidean (planar) geometry.

### Theorem (Similar triangles have equal side ratios)

*Let  $\triangle ABC$  and  $\triangle A'B'C'$  be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is*

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.

### Theorem (Similar triangles have equal side ratios)

*Let  $\triangle ABC$  and  $\triangle A'B'C'$  be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is*

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.

### Theorem (Similar triangles have equal side ratios)

*Let  $\triangle ABC$  and  $\triangle A'B'C'$  be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is*

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




The following statement is proved in the subject of Euclidean (planar) geometry.

### Theorem (Similar triangles have equal side ratios)

*Let  $\triangle ABC$  and  $\triangle A'B'C'$  be two similar triangles. Then the ratios of the lengths of the sides of the two triangles are equal, that is*

$$\frac{|AB|}{|BC|} = \frac{|A'B'|}{|B'C'|} \quad \frac{|BC|}{|CA|} = \frac{|B'C'|}{|C'A'|} \quad \frac{|CA|}{|AB|} = \frac{|C'A'|}{|A'B'|}$$




$\sin(\alpha + \beta)$ ,  $\cos(\alpha + \beta)$  via  $\sin \alpha$ ,  $\sin \beta$ ,  $\cos \alpha$ ,  $\cos \beta$

$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$




$\sin(\alpha + \beta)$ ,  $\cos(\alpha + \beta)$  via  $\sin \alpha$ ,  $\sin \beta$ ,  $\cos \alpha$ ,  $\cos \beta$



$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = ?$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = \frac{|CD|}{|OC|}$$

$$\cos(\alpha + \beta) = ?$$


$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\sin(\alpha + \beta) = \frac{|CD|}{|OC|} = |CD|$$

$$\cos(\alpha + \beta) = ?$$

$\sin(\alpha + \beta), \cos(\alpha + \beta)$  via  $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\sin(\alpha + \beta) = \frac{|CD|}{|OC|} = |CD|$$

$$\cos(\alpha + \beta) = ?$$




# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ|\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$|QD| = |BA|$$

$$| \square DABQ$$

$$\begin{aligned} \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned} |QD| &= |BA| \\ &= \sin \alpha |OB| \end{aligned}$$

$$\left| \begin{array}{l} \square DABQ \\ \triangle OAB \end{array} \right|$$

$$\begin{aligned} \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}
 |QD| &= |BA| \\
 &= \sin \alpha |OB| \\
 &= \sin \alpha \cos \beta |OC|
 \end{aligned}
 \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right.$$

$$\begin{aligned}
 \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\
 &= |QD| + |CQ|
 \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ|\end{aligned}$$

$$\begin{aligned}|QD| &= |BA| \\ &= \sin \alpha |OB| \\ &= \sin \alpha \cos \beta |OC| \\ &= \sin \alpha \cos \beta\end{aligned} \quad \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right.$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}
 |QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\
 &= \sin \alpha |OB| \\
 &= \sin \alpha \cos \beta |OC| \\
 &= \sin \alpha \cos \beta
 \end{aligned}$$

$$\begin{aligned}
 \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\
 &= |QD| + |CQ| \\
 &= \sin \alpha \cos \beta + ?
 \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\ &= \sin \alpha |OB| \\ &= \sin \alpha \cos \beta |OC| \\ &= \sin \alpha \cos \beta \\ |CQ| &= \end{aligned}$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\ &= \sin \alpha |OB| \\ &= \sin \alpha \cos \beta |OC| \\ &= \sin \alpha \cos \beta \\ |CQ| &= \end{aligned}$$



# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\ &= \sin \alpha |OB| \\ &= \sin \alpha \cos \beta |OC| \\ &= \sin \alpha \cos \beta \\ |CQ| &= \end{aligned}$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\ &= \sin \alpha |OB| \\ &= \sin \alpha \cos \beta |OC| \\ &= \sin \alpha \cos \beta \\ |CQ| &= \end{aligned}$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}
 |QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\
 &= \sin \alpha |OB| \\
 &= \sin \alpha \cos \beta |OC| \\
 &= \sin \alpha \cos \beta \\
 |CQ| &= \cos \alpha |CB| & \left| \triangle CQB \right.
 \end{aligned}$$

$$\begin{aligned}
 \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\
 &= |QD| + |CQ| \\
 &= \sin \alpha \cos \beta + ?
 \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| && \square DABQ \\ &= \sin \alpha |OB| && \triangle OAB \\ &= \sin \alpha \cos \beta |OC| && \triangle OBC \\ &= \sin \alpha \cos \beta \\ |CQ| &= \cos \alpha |CB| && \triangle CQB \\ &= \cos \alpha \sin \beta |OC| && \triangle OBC\end{aligned}$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + ?\end{aligned}$$


$$\cos(\alpha + \beta) = ?$$

$$\begin{aligned}|QD| &= |BA| && \square DABQ \\ &= \sin \alpha |OB| && \triangle OAB \\ &= \sin \alpha \cos \beta |OC| && \triangle OBC \\ &= \sin \alpha \cos \beta \\ |CQ| &= \cos \alpha |CB| && \triangle CQB \\ &= \cos \alpha \sin \beta |OC| && \triangle OBC \\ &= \cos \alpha \sin \beta\end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

$$\begin{array}{l|l} |QD| = |BA| & \square DABQ \\ = \sin \alpha |OB| & \triangle OAB \\ = \sin \alpha \cos \beta |OC| & \triangle OBC \\ = \sin \alpha \cos \beta & \\ |CQ| = \cos \alpha |CB| & \triangle CQB \\ = \cos \alpha \sin \beta |OC| & \triangle OBC \\ = \cos \alpha \sin \beta & \end{array}$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$




$$\begin{aligned}
 |QD| &= |BA| & \left| \begin{array}{l} \square DABQ \\ \triangle OAB \\ \triangle OBC \end{array} \right. \\
 &= \sin \alpha |OB| \\
 &= \sin \alpha \cos \beta |OC| \\
 &= \sin \alpha \cos \beta
 \end{aligned}$$

$$\begin{aligned}
 |CQ| &= \cos \alpha |CB| & \left| \begin{array}{l} \triangle CQB \\ \triangle OBC \end{array} \right. \\
 &= \cos \alpha \sin \beta |OC| \\
 &= \cos \alpha \sin \beta
 \end{aligned}$$

$$\begin{aligned}
 \sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\
 &= |QD| + |CQ| \\
 &= \sin \alpha \cos \beta + \cos \alpha \sin \beta
 \end{aligned}$$

$$\cos(\alpha + \beta) = ?$$

# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + \cos \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}\cos(\alpha + \beta) &= \frac{|OD|}{|OC|} = |OD| \\ &= |OA| - |DA| \\ &= \cos \alpha \cos \beta - \sin \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}|QD| &= |BA| && \square DABQ \\ &= \sin \alpha |OB| && \triangle OAB \\ &= \sin \alpha \cos \beta |OC| && \triangle OBC \\ &= \sin \alpha \cos \beta\end{aligned}$$


$$\begin{aligned}|CQ| &= \cos \alpha |CB| && \triangle CQB \\ &= \cos \alpha \sin \beta |OC| && \triangle OBC \\ &= \cos \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}|OA| &= \cos \alpha |OB| && \triangle OAB \\ &= \cos \alpha \cos \beta |OC| && \triangle OBC \\ &= \cos \alpha \cos \beta\end{aligned}$$

$$\begin{aligned}|DA| &= |QB| && \square DABQ \\ &= \sin \alpha |CB| && \triangle CQB \\ &= \sin \alpha \sin \beta |OC| && \triangle OBC \\ &= \sin \alpha \sin \beta\end{aligned}$$



# $\sin(\alpha + \beta), \cos(\alpha + \beta)$ via $\sin \alpha, \sin \beta, \cos \alpha, \cos \beta$



$$\begin{aligned}\sin(\alpha + \beta) &= \frac{|CD|}{|OC|} = |CD| \\ &= |QD| + |CQ| \\ &= \sin \alpha \cos \beta + \cos \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}\cos(\alpha + \beta) &= \frac{|OD|}{|OC|} = |OD| \\ &= |OA| - |DA| \\ &= \cos \alpha \cos \beta - \sin \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}|QD| &= |BA| && \square DABQ \\ &= \sin \alpha |OB| && \triangle OAB \\ &= \sin \alpha \cos \beta |OC| && \triangle OBC \\ &= \sin \alpha \cos \beta\end{aligned}$$

$$\begin{aligned}|CQ| &= \cos \alpha |CB| && \triangle CQB \\ &= \cos \alpha \sin \beta |OC| && \triangle OBC \\ &= \cos \alpha \sin \beta\end{aligned}$$

$$\begin{aligned}|OA| &= \cos \alpha |OB| && \triangle OAB \\ &= \cos \alpha \cos \beta |OC| && \triangle OBC \\ &= \cos \alpha \cos \beta\end{aligned}$$

$$\begin{aligned}|DA| &= |QB| && \square DABQ \\ &= \sin \alpha |CB| && \triangle CQB \\ &= \sin \alpha \sin \beta |OC| && \triangle OBC \\ &= \sin \alpha \sin \beta\end{aligned}$$

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

- We gave a geometric proof of the sum formulas when the two angles are acute and their sum is less than  $\pi = 90^\circ$ .

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

- We gave a geometric proof of the sum formulas when the two angles are acute and their sum is less than  $\pi = 90^\circ$ .
- The theorem holds for all angles  $\alpha, \beta$  without any restrictions.

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

- We gave a geometric proof of the sum formulas when the two angles are acute and their sum is less than  $\pi = 90^\circ$ .
- The theorem holds for all angles  $\alpha, \beta$  without any restrictions.
- This can be shown by combining the preceding proof with identities such as  $\cos\left(\frac{\pi}{2} - \alpha\right) = \sin \alpha$ ,  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$ .

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

- We gave a geometric proof of the sum formulas when the two angles are acute and their sum is less than  $\pi = 90^\circ$ .
- The theorem holds for all angles  $\alpha, \beta$  without any restrictions.
- This can be shown by combining the preceding proof with identities such as  $\cos\left(\frac{\pi}{2} - \alpha\right) = \sin \alpha$ ,  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$ .
- There is a theoretically more advanced (but algebraically simpler) proof using Euler's formula (to be studied later/in another course).

# Trig Functions of Sums and Differences of Angles

## Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

- We gave a geometric proof of the sum formulas when the two angles are acute and their sum is less than  $\pi = 90^\circ$ .
- The theorem holds for all angles  $\alpha, \beta$  without any restrictions.
- This can be shown by combining the preceding proof with identities such as  $\cos\left(\frac{\pi}{2} - \alpha\right) = \sin \alpha$ ,  $\cos\left(\frac{\pi}{2} + \alpha\right) = -\sin \alpha$ .
- There is a theoretically more advanced (but algebraically simpler) proof using Euler's formula (to be studied later/in another course).
- The difference formulas are a consequence of the sum formulas and the fact that  $\sin$  is an odd function and  $\cos$  is even.

# Trig Functions of Differences of Angles

## Example

Prove the identities

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

from the (already demonstrated) identities

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin(\alpha + (-\beta))$$

$$= \sin \alpha \cos(-\beta) + \cos \alpha \sin(-\beta)$$

cos is even ,  
sin is odd

$$= \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos(\alpha + (-\beta))$$

$$= \cos \alpha \cos(-\beta) - \sin \alpha \sin(-\beta)$$

cos is even ,  
sin is odd

$$= \cos \alpha \cos \beta + \sin \alpha \sin \beta$$



## Example

Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ)$$

## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

## Example

Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$





## Example

Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ) = \cos(45^\circ + 60^\circ) \\ = ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$   
**Angle sum f-la**



## Example



Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= ? \cdot ? - ? \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot ? - ? \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot ? - ? \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la





## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - ? \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - ? \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot ?$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.



$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$

$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2}$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$



$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2}$$

$$= \frac{\sqrt{2} - \sqrt{6}}{4}$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example

Find the exact value of the trigonometric function using radicals.

$$\cos(105^\circ) = \cos(45^\circ + 60^\circ)$$



$$= \cos(45^\circ) \cos(60^\circ) - \sin(45^\circ) \sin(60^\circ)$$

$$= \frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2}$$

$$= \frac{\sqrt{2} - \sqrt{6}}{4}$$

we know the trig  
f-ns of  $45^\circ$  and  $60^\circ$

Angle sum f-la



## Example


Use the angle sum/difference formulas to simplify.

$$\cos\left(\frac{\pi}{2} - x\right)$$



## Example

Use the angle sum/difference formulas to simplify.



$$\begin{aligned}\cos\left(\frac{\pi}{2} - x\right) &= \cos\left(\frac{\pi}{2}\right)\cos x + \sin\left(\frac{\pi}{2}\right)\sin x \\ &= 0 \cdot \cos(x) + 1 \cdot \sin x \\ &= \sin x\end{aligned}$$

## Example


Use the angle sum/difference formulas to simplify.

$$\cot \left( \frac{3\pi}{2} + x \right)$$

## Example

Use the angle sum/difference formulas to simplify.

$$\begin{aligned}
 \cot\left(\frac{3\pi}{2} + x\right) &= \frac{\cos\left(\frac{3\pi}{2} + x\right)}{\sin\left(\frac{3\pi}{2} + x\right)} \\
 &= \frac{\cos\left(\frac{3\pi}{2}\right)\cos x - \sin\left(\frac{3\pi}{2}\right)\sin x}{\sin\left(\frac{3\pi}{2}\right)\cos x + \cos\left(\frac{3\pi}{2}\right)\sin x} \\
 &= \frac{0 \cdot \cos x - (-1)\sin x}{(-1)\cos x + 0 \cdot \sin x} \\
 &= \frac{-\cos x}{-\sin x} = -\frac{\sin x}{\cos x} \\
 &= -\tan x
 \end{aligned}$$



## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\tan(\pi + x) = \frac{\sin(\pi + x)}{\cos(\pi + x)}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x}\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{? \cdot \cos x + ? \cdot \sin x}{? \cdot \cos x - ? \cdot \sin x}
 \end{aligned}$$



## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{0 \cdot \cos x + ? \cdot \sin x}{? \cdot \cos x - ? \cdot \sin x}
 \end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \text{red } \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{0 \cdot \cos x + \text{red } ? \cdot \sin x}{? \cdot \cos x - ? \cdot \sin x}
 \end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \text{red } \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{0 \cdot \cos x + \text{red } (-1) \cdot \sin x}{? \cdot \cos x - ? \cdot \sin x}
 \end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x} \\
 &= \frac{-\sin x}{-\cos x} = \frac{\sin x}{\cos x} = \tan x
 \end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\ &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\ &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - ? \cdot \sin x}\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\ &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x}\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\ &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\ &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x} \\ &= \frac{-\sin x}{-\cos x}\end{aligned}$$



## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}
 \tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\
 &= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\
 &= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x} \\
 &= \frac{-\sin x}{-\cos x} \\
 &= \frac{\sin x}{\cos x}
 \end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\&= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\&= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x} \\&= \frac{-\sin x}{-\cos x} \\&= \frac{\sin x}{\cos x} \\&= \tan x,\end{aligned}$$

## Example

Show that  $\tan(\pi + x) = \tan x$  using the angle sum formulas.

$$\begin{aligned}\tan(\pi + x) &= \frac{\sin(\pi + x)}{\cos(\pi + x)} \\&= \frac{\sin \pi \cos x + \cos \pi \sin x}{\cos \pi \cos x - \sin \pi \sin x} \\&= \frac{0 \cdot \cos x + (-1) \cdot \sin x}{(-1) \cdot \cos x - 0 \cdot \sin x} \\&= \frac{-\sin x}{-\cos x} \\&= \frac{\sin x}{\cos x} \\&= \tan x,\end{aligned}$$

as desired.

### Proposition ( $\tan, \cot$ are $\pi$ -periodic)

*The tangent and cotangent functions are  $\pi$ -periodic, in other words,*

$$\tan(\theta + \pi) = \tan \theta$$

$$\cot(\theta + \pi) = \cot \theta$$

Recall the angle sum formula  $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ .

### Example

Show that the Pythagorean identity  $\sin^2 \theta + \cos^2 \theta = 1$  follows from the angle difference formula.

Recall the angle sum formula  $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ .

### Example

Show that the Pythagorean identity  $\sin^2 \theta + \cos^2 \theta = 1$  follows from the angle difference formula.

$$\begin{aligned} 1 &= \cos 0 \\ &= \cos(\theta - \theta) \\ &= \cos \theta \cos \theta + \sin \theta \sin \theta \\ &= \cos^2 \theta + \sin^2 \theta, \end{aligned}$$

as desired.

## Example

Prove the angle sum formula  $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$ .

$$\tan(\alpha + \beta) =$$

## Example

Prove the angle sum formula  $\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$ .

$$\begin{aligned}
 \tan(\alpha + \beta) &= \frac{\sin(\alpha + \beta)}{\cos(\alpha + \beta)} \\
 &= \frac{(\sin \alpha \cos \beta + \cos \alpha \sin \beta) \frac{1}{\cos \alpha \cos \beta}}{(\cos \alpha \cos \beta - \sin \alpha \sin \beta) \frac{1}{\cos \alpha \cos \beta}} \\
 &= \frac{\frac{\sin \alpha \cancel{\cos \beta}}{\cos \alpha \cancel{\cos \beta}} + \frac{\cancel{\cos \alpha} \sin \beta}{\cancel{\cos \alpha} \cos \beta}}{\frac{\cancel{\cos \alpha} \cos \beta}{\cancel{\cos \alpha} \cos \beta} - \frac{\sin \alpha \sin \beta}{\cos \alpha \cos \beta}} \\
 &= \frac{\frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta}}{1 - \frac{\sin \alpha}{\cos \alpha} \cdot \frac{\sin \beta}{\cos \beta}} \\
 &= \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}
 \end{aligned}$$



# Double angle formulas

## Proposition (Double angle formulas)

$$\begin{aligned}\sin(2\alpha) &= 2 \sin \alpha \cos \alpha \\ \cos(2\alpha) &= \cos^2 \alpha - \sin^2 \alpha \\ &= 2 \cos^2 \alpha - 1 \\ &= 1 - 2 \sin^2 \alpha\end{aligned}$$

# Double angle formulas

## Proposition (Double angle formulas)

$$\begin{aligned}\sin(2\alpha) &= 2 \sin \alpha \cos \alpha \\ \cos(2\alpha) &= \cos^2 \alpha - \sin^2 \alpha \\ &= 2 \cos^2 \alpha - 1 \\ &= 1 - 2 \sin^2 \alpha\end{aligned}$$

- The double angle formulas play a special role in integration.

## Example

Derive the double-angle formulas.

$$\sin(2\alpha) =$$

$$\cos(2\alpha) =$$

## Example

Derive the double-angle formulas.


$$\begin{aligned}\sin(2\alpha) &= \sin(\alpha + \alpha) \\ &= \sin \alpha \cos \alpha + \cos \alpha \sin \alpha \\ &= 2 \sin \alpha \cos \alpha\end{aligned}$$

$$\begin{aligned}\cos(2\alpha) &= \cos(\alpha + \alpha) \\ &= \cos \alpha \cos \alpha - \sin \alpha \sin \alpha \\ &= \cos^2 \alpha - \sin^2 \alpha \\ &= \cos^2 \alpha - (1 - \cos^2 \alpha) \\ &= 2 \cos^2 \alpha - 1 \\ &= 1 - \sin^2 \alpha - \sin^2 \alpha \\ &= 1 - 2 \sin^2 \alpha\end{aligned}$$

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\cos 105^\circ$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\cos 105^\circ$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\cos 105^\circ = \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\cos 105^\circ = \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} \quad \left| \quad \cos 105^\circ ? 0 \right.$$






Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\cos 105^\circ = \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} \quad \left| \quad \cos 105^\circ < 0 \right.$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned} \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} & \left| \cos 105^\circ < 0 \right. \\ &= - \sqrt{\frac{1 + \cos(210^\circ)}{2}} \end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned}\cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} & \left| \cos 105^\circ < 0 \right. \\ &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}}\end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned} \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} & \left| \cos 105^\circ < 0 \right. \\ &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned} \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} && \left| \cos 105^\circ < 0 \right. \\ &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \\ &= -\sqrt{\frac{1 - \cos(30^\circ)}{2}} \end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned}
 \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} & \left| \cos 105^\circ < 0 \right. \\
 &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \cos(30^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - ?}{2}}
 \end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned}
 \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} & \left| \cos 105^\circ < 0 \right. \\
 &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \cos(30^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}}
 \end{aligned}$$




Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned}
 \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} && \left| \cos 105^\circ < 0 \right. \\
 &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \cos(30^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}} = -\sqrt{\frac{2 - \sqrt{3}}{2 \cdot 2}}
 \end{aligned}$$






Recall the half angle formula  $\cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$ .

## Example

Using radicals, find the exact value of the trigonometric expression.

$$\begin{aligned}
 \cos 105^\circ &= \pm \sqrt{\frac{1 + \cos(2 \cdot 105^\circ)}{2}} && \left| \cos 105^\circ < 0 \right. \\
 &= -\sqrt{\frac{1 + \cos(210^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \cos(30^\circ)}{2}} \\
 &= -\sqrt{\frac{1 - \frac{\sqrt{3}}{2}}{2}} = -\sqrt{\frac{2 - \sqrt{3}}{2 \cdot 2}} \\
 &= -\frac{\sqrt{2 - \sqrt{3}}}{2}
 \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned}\cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha)\end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned}\cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha)\end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\begin{aligned}\cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ \color{red}{2}\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{\color{red}{2}}\end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned}$$





## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha - 1 = \cos(2\alpha)$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned} \qquad \begin{aligned} \color{red}{\cos(2\alpha)} &= 2\cos^2 \alpha - 1 \\ 2\cos^2 \alpha - 1 &= \color{red}{\cos(2\alpha)} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned} \qquad \begin{aligned} \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\cos^2 \alpha - 1 &= \cos(2\alpha) \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} \end{aligned}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha = 1 + \cos(2\alpha)$$

$$\cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \qquad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\cos(2\alpha) = 1 - 2\sin^2 \alpha$$

$$2\sin^2 \alpha = 1 - \cos(2\alpha)$$

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2}$$

$$\cos(2\alpha) = 2\cos^2 \alpha - 1$$

$$2\cos^2 \alpha = 1 + \cos(2\alpha)$$

$$\cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

### Corollary (Half-Angle Formulas)

$$\sin \left( \frac{\beta}{2} \right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos \left( \frac{\beta}{2} \right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$



## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

### Proof.

$$\begin{aligned} \cos(2\alpha) &= 1 - 2\sin^2 \alpha & \cos(2\alpha) &= 2\cos^2 \alpha - 1 \\ 2\sin^2 \alpha &= 1 - \cos(2\alpha) & 2\cos^2 \alpha &= 1 + \cos(2\alpha) \\ \sin^2 \alpha &= \frac{1 - \cos(2\alpha)}{2} & \cos^2 \alpha &= \frac{1 + \cos(2\alpha)}{2} \end{aligned}$$



### Corollary

$$\sin \alpha = \pm \sqrt{\frac{1 - \cos(2\alpha)}{2}} \quad \cos \alpha = \pm \sqrt{\frac{1 + \cos(2\alpha)}{2}}$$

### Corollary (Half-Angle Formulas)

$$\sin\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 - \cos \beta}{2}} \quad \cos\left(\frac{\beta}{2}\right) = \pm \sqrt{\frac{1 + \cos \beta}{2}}$$

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express  $\sin^k \alpha$  and  $\cos^k \alpha$  via lower powers of the sin and cos functions (applied to angles other than  $\alpha$ ).

## Proposition (Power-Reducing Formulas)

$$\sin^2 \alpha = \frac{1 - \cos(2\alpha)}{2} \quad \cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2}$$

- The power reducing formulas are used to express  $\sin^k \alpha$  and  $\cos^k \alpha$  via lower powers of the sin and cos functions (applied to angles other than  $\alpha$ ).
- This technique will play a key role in integration (studied later/in another course).

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\sin^4 \alpha$$

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\sin^4 \alpha = \left( \sin^2 \alpha \right)^2$$

Recall the formulas:  $\sin^2 \beta = ?$  ,  $\cos^2 \beta = ?$  .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= \left( \sin^2 \alpha \right)^2 \\ &= \left( ? \right)^2\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= \left( \sin^2 \alpha \right)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} \left( ? \right) \end{aligned}$$



Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned} \sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} ( ? ) \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\ &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\ &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha))\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = ?$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\&= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\&= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\&= \frac{1}{4} (1 - 2\cos(2\alpha) + ?)\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\&= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\&= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \cos^2(2\alpha) \right) \\&= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\&= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\&= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\&= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right)\end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \cos^2(2\alpha) \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( \textcolor{red}{1} - 2\cos(2\alpha) + \frac{\cos(\textcolor{red}{2} \cdot \textcolor{red}{2}\alpha)}{2} + \textcolor{red}{\frac{1}{2}} \right) \\
 &= \frac{1}{4} \left( \textcolor{red}{\frac{3}{2}} - 2\cos(2\alpha) + \frac{\cos(\textcolor{red}{4}\alpha)}{2} \right)
 \end{aligned}$$

Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}
 \sin^4 \alpha &= (\sin^2 \alpha)^2 \\
 &= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\
 &= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\
 &= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\
 &= \frac{1}{4} \left( \frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\
 &= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))
 \end{aligned}$$



Recall the formulas:  $\sin^2 \beta = \frac{1 - \cos(2\beta)}{2}$ ,  $\cos^2 \beta = \frac{\cos(2\beta) + 1}{2}$ .

## Example

Rewrite  $\sin^4 \alpha$  in terms of first powers of the cosines and sines of multiples of the angle  $\alpha$ .

$$\begin{aligned}\sin^4 \alpha &= (\sin^2 \alpha)^2 \\&= \left( \frac{1 - \cos(2\alpha)}{2} \right)^2 \\&= \frac{1}{4} (1 - 2\cos(2\alpha) + \cos^2(2\alpha)) \\&= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha) + 1}{2} \right) \\&= \frac{1}{4} \left( 1 - 2\cos(2\alpha) + \frac{\cos(2 \cdot 2\alpha)}{2} + \frac{1}{2} \right) \\&= \frac{1}{4} \left( \frac{3}{2} - 2\cos(2\alpha) + \frac{\cos(4\alpha)}{2} \right) \\&= \frac{1}{8} (3 - 4\cos(2\alpha) + \cos(4\alpha))\end{aligned}$$