

# Calculus I

## Volumes of solids of revolution

Todor Milev

2019

# Outline

1

## Volumes

# Outline

1 Volumes

2 Volumes by Cylindrical Shells

# License to use and redistribute


These lecture slides and their L<sup>A</sup>T<sub>E</sub>X source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,


as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/> and the links therein.


# Volumes




Volumes of solids are found/defined via integration.




- How do we find the volume of a solid  $S$ ?




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .



- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .
- Approximate this part of  $S$ :
- Pick a sample point  $x^*$  between  $x_1$  and  $x_2$ . Use a solid that has the same constant cross-sectional area  $A(x^*)$  between  $x_1$  and  $x_2$ .




- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .
- Approximate this part of  $S$ :
- Pick a sample point  $x^*$  between  $x_1$  and  $x_2$ . Use a solid that has the same constant cross-sectional area  $A(x^*)$  between  $x_1$  and  $x_2$ .
- Let  $\Delta x$  be the distance from  $x_1$  to  $x_2$ .



Approx. volume of slab:

$$A(x^*)\Delta x$$

- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .
- Approximate this part of  $S$ :
- Pick a sample point  $x^*$  between  $x_1$  and  $x_2$ . Use a solid that has the same constant cross-sectional area  $A(x^*)$  between  $x_1$  and  $x_2$ .
- Let  $\Delta x$  be the distance from  $x_1$  to  $x_2$ .




Approx. volume of slab:

$$A(x^*)\Delta x$$

Approx. volume of  $S$ :

$$V \approx \sum_{i=1}^n A(x_i^*)\Delta x$$

- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .
- Approximate this part of  $S$ :
- Pick a sample point  $x^*$  between  $x_1$  and  $x_2$ . Use a solid that has the same constant cross-sectional area  $A(x^*)$  between  $x_1$  and  $x_2$ .
- Let  $\Delta x$  be the distance from  $x_1$  to  $x_2$ .



Approx. volume of slab:

$$A(x^*)\Delta x$$

Approx. volume of  $S$ :

$$V \approx \sum_{i=1}^n A(x_i^*)\Delta x$$

Exact volume of  $S$ :

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n A(x_i^*)\Delta x$$

- How do we find the volume of a solid  $S$ ?
- Let  $P_x$  be the plane perpendicular to the  $x$ -axis and passing through the point  $x$ .
- The intersection of  $P_x$  with  $S$  is called a cross-section.
- Let  $A(x)$  be the area of this cross-section.
- Consider the part of  $S$  between two planes  $P_{x_1}$  and  $P_{x_2}$ .
- Approximate this part of  $S$ :
- Pick a sample point  $x^*$  between  $x_1$  and  $x_2$ . Use a solid that has the same constant cross-sectional area  $A(x^*)$  between  $x_1$  and  $x_2$ .
- Let  $\Delta x$  be the distance from  $x_1$  to  $x_2$ .


## Definition (Volume)

Let  $S$  be a solid that lies between  $x = a$  and  $x = b$ . If the cross-sectional area of  $S$  in the plane  $P_x$  is a continuous function  $A(x)$ , then the volume of  $S$  is

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n A(x_i^*) \Delta x = \int_a^b A(x) dx$$


## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.




## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.



## Example


Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = ?$



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .



## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .




$$V = \int_{?}^{?} A(x)dx$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .




$$V = \int_{?}^{?} A(x)dx$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .




$$V = \int_{?}^{?} A(x)dx$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .
- The  $x$  coords. of the solid are between 0 and 1, so its volume is




$$V = \int_0^1 A(x)dx$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .
- The  $x$  coords. of the solid are between 0 and 1, so its volume is




$$V = \int_0^1 A(x)dx = \int_0^1 \pi x dx$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .
- The  $x$  coords. of the solid are between 0 and 1, so its volume is




$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 \pi x \, dx \\ &= \left[ \frac{\pi x^2}{2} \right]_0^1 \end{aligned}$$

## Example

Find the volume of the solid obtained by rotating about the  $x$ -axis the region under the curve  $y = \sqrt{x}$  from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point  $(x, 0)$  has radius  $\sqrt{x}$ .
- The area of the cross-section is  $A(x) = \pi (\sqrt{x})^2 = \pi x$ .
- The volume of a single approximating section is  $A(x)\Delta x$ .
- The  $x$  coords. of the solid are between 0 and 1, so its volume is



$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 \pi x \, dx \\ &= \left[ \pi \frac{x^2}{2} \right]_0^1 = \frac{\pi}{2} \end{aligned}$$


## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis **the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .**




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .  
Cross-section:




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .  
Cross-section: washer, center:  $(x, 0)$ .



## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .  
Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius: ? area: .

Outer disk radius: , area: .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius:  $x$ , area: .

Outer disk radius: , area: .




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius:  $x$ , area: ? .

Outer disk radius: , area:




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius: , area:




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius: ? , area:




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area: ?




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .



## Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int A(x)dx$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int_{?}^{?} A(x)dx$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int_0^1 A(x)dx$$



## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .


Cross-section: washer, center:  $(x, 0)$ . Area:

$$A(x) = \text{Area outer disk} - \text{Area inner disk}$$

Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$



## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .


Cross-section: washer, center:  $(x, 0)$ . Area:

$$A(x) = \text{Area outer disk} - \text{Area inner disk}$$

Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$



## Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$V = \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ ? + ? + ? \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + ? + ? \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + ? + ? \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + \frac{x^3}{3} + ? \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + \frac{x^3}{3} + ? \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \\ &= \pi \left( \frac{1}{5} + \frac{1}{3} + 1 \right) \end{aligned}$$




## Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the  $x$ -axis the region bounded by  $y = x^2 + 1$ ,  $y = x$ ,  $x = 0$ , and  $x = 1$ .

Cross-section: washer, center:  $(x, 0)$ . Area:  
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$   
Inner disk radius:  $x$ , area:  $\pi x^2$ .

Outer disk radius:  $x^2 + 1$ , area:  $\pi(x^2 + 1)^2$ .

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[ \frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \\ &= \pi \left( \frac{1}{5} + \frac{1}{3} + 1 \right) = \frac{23}{15}\pi \end{aligned}$$




## Example (Rotation About a Line Parallel to the $x$ -axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the $x$ -axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .



## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

**Cross-section:** a circle centered at ?

radius: ?

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

**Cross-section:** a circle centered at ?

radius: ?

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at ?

radius: ?

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius: ?

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius: ?

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) =$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = ?$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$V = \int_{?}^{?} A(x)dx$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$V = \int_0^2 A(x)dx$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$V = \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ ? - ? + ? \right]_0^2 \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ \frac{x^5}{5} - ? + ? \right]_0^2 \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ \frac{x^5}{5} - ? + ? \right]_0^2 \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ \frac{x^5}{5} - x^4 + ? \right]_0^2 \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ \frac{x^5}{5} - x^4 + ? \right]_0^2 \end{aligned}$$



## Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[ \frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \end{aligned}$$




## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .



$$\begin{aligned}
 V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\
 &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\
 &= \pi \left[ \frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\
 &= \pi \left( \frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right)
 \end{aligned}$$


## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,

radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .



$$\begin{aligned}
 V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\
 &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\
 &= \pi \left[ \frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\
 &= \pi \left( \frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right)
 \end{aligned}$$


## Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line  $y = 1$  the region bounded by  $y = -x^2 + 2x + 1$  and  $y = 1$ .

Cross-section: a circle centered at  $(x, 1)$ ,


radius:  $(-x^2 + 2x + 1) - 1$ ,

area:  $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$ .




$$\begin{aligned}
 V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\
 &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\
 &= \pi \left[ \frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\
 &= \pi \left( \frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right) \\
 &= \pi \left( \frac{32}{5} - 16 + \frac{32}{3} \right) = \frac{16}{15}\pi.
 \end{aligned}$$

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the **x-axis** around ...




- ... the x-axis.




- ... the y-axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...




- ... the  $x$ -axis.




- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.





- ... the  $y$ -axis.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- ... the  $y$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
  - Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
  - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
  - We understand the problem.
- ... the  $y$ -axis.
  - Approx. with washers:

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
  - Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
  - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
  - We understand the problem.
- ... the  $y$ -axis.
  - Approx. with washers: need **inner rad.  $x_i$** .

Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...



- ... the  $x$ -axis.
  - Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
  - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
  - We understand the problem.
- ... the  $y$ -axis.
  - Approx. with washers: need inner rad.  $x_i$  & outer rad.  $x_o$ .



Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...



- ... the  $x$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

- ... the  $y$ -axis.
- Approx. with washers: need inner rad.  $x_i$  & outer rad.  $x_o$ .
- $x_i$  and  $x_o$ : solutions to cubic:  
 $-x^3 + 2x^2 - y = 0$ .



Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.

- ... the  $y$ -axis.
- Approx. with washers: need inner rad.  $x_i$  & outer rad.  $x_o$ .
- $x_i$  and  $x_o$ : solutions to cubic:  $-x^3 + 2x^2 - y = 0$ . Solving for  $x$  requires lots of algebra.



Find the volume obtained by rotating the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis around ...





- ... the  $x$ -axis.
- Approximate the volume using circular cylinders with radius  $2x^2 - x^3$  and height  $\Delta x$ .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$ .
- We understand the problem.
- ... the  $y$ -axis.
- Approx. with washers: need inner rad.  $x_i$  & outer rad.  $x_o$ .
- $x_i$  and  $x_o$ : solutions to cubic:  $-x^3 + 2x^2 - y = 0$ . Solving for  $x$  requires lots of algebra.
- We show a simpler technique.



- Consider the solid obtained by rotating around the  $y$ -axis the region bounded above by  $y = 2x^2 - x^3$  and below by the  $x$ -axis.




- Consider the solid obtained by rotating around the  $y$ -axis the region bounded above by  $y = 2x^2 - x^3$  and below by the  $x$ -axis.
- Approximate this solid by nested cylindrical shells.




- Consider the solid obtained by rotating around the  $y$ -axis the region bounded above by  $y = 2x^2 - x^3$  and below by the  $x$ -axis.
- Approximate this solid by nested cylindrical shells.
- Cylindrical shells are solids obtained by taking a cylinder and removing from its center another cylinder of equal height but smaller radius.


- Consider a cylindrical shell with:




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .




- Consider a cylindrical shell with:
  - outer radius  $r_2$ ,
  - inner radius  $r_1$ ,
  - height  $h$ .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}}$$




- Consider a cylindrical shell with:
  - outer radius  $r_2$ ,
  - inner radius  $r_1$ ,
  - height  $h$ .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h$$




- Consider a cylindrical shell with:
  - outer radius  $r_2$ ,
  - inner radius  $r_1$ ,
  - height  $h$ .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h$$




- Consider a cylindrical shell with:
  - outer radius  $r_2$ ,
  - inner radius  $r_1$ ,
  - height  $h$ .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$$




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let  $\Delta r = r_2 - r_1$ .




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let  $\Delta r = r_2 - r_1$ .
- Let  $r = \frac{r_2 + r_1}{2}$ .




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let  $\Delta r = r_2 - r_1$ .
- Let  $r = \frac{r_2 + r_1}{2}$ .
- Then  $V_{\text{shell}} = 2\pi rh\Delta r$ .







- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h$ .
- Let  $\Delta r = r_2 - r_1$ .
- Let  $r = \frac{r_2 + r_1}{2}$ .
- Then  $V_{\text{shell}} = 2\pi r h \Delta r$ .




- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h$ .
- Let  $\Delta r = r_2 - r_1$ .
- Let  $r = \frac{r_2 + r_1}{2}$ .
- Then  $V_{\text{shell}} = 2\pi r h \Delta r$ .

- Consider a cylindrical shell with:
- outer radius  $r_2$ ,
- inner radius  $r_1$ ,
- height  $h$ .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let  $\Delta r = r_2 - r_1$ .
- Let  $r = \frac{r_2 + r_1}{2}$ .
- Then  $V_{\text{shell}} = 2\pi rh\Delta r$ .







Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.




Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.




Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.

$$V_{\text{shell}} = 2\pi r h \Delta r$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.  
Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).

$$V_{\text{shell}} = 2\pi r h \Delta r = 2\pi r f(r) \Delta r.$$




Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.  
Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$




Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.  
Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi r f(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$




Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
Approximate the volume by cylindrical shells.  
Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  
 $h = f(r)$  ( $r$ =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x = \int_a^b 2\pi x f(x) dx.$$



Consider a solid obtained by rotating the region under  $f(x)$  around the  $y$  axis.  
 Approximate the volume by cylindrical shells.  
 Select the height of an individual shell to be  $h = f(r)$  ( $r$ =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are  $n$  cylindrical shells and let  $x_1, \dots, x_n$  be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to  $\infty$  to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x = \int_a^b 2\pi x f(x) dx.$$

The endpoints of integration are the endpoints of the rotated region.

## Definition (Volume by Cylindrical Shells)

The volume of the solid obtained by rotating around the  $y$ -axis the region under the curve  $y = f(x)$  from  $a$  to  $b$  is


$$V = \int_a^b 2\pi x f(x) dx.$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.


## Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height: ? ; circumference: ? ; infinitesimal volume: ? .


## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height: ? ; circumference: ? ; infinitesimal volume: ? .


## Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference: ? ; infinitesimal volume: ? .


## Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference: ? ; infinitesimal volume: ? .


## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume: ?


## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume: ?


## Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .


## Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .


## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .


## Example



Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; **infinitesimal** volume:  $2\pi x(2x^2 - x^3)dx$ .

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$V = \int (2\pi x)(2x^2 - x^3)dx$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$V = \int_{?}^{?} (2\pi x)(2x^2 - x^3)dx$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$V = \int_0^2 (2\pi x)(2x^2 - x^3)dx$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$V = \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ ? - ? \right]_0^2 \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - ? \right]_0^2 \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - ? \right]_0^2 \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the  $y$ -axis the region bounded by  $y = 2x^2 - x^3$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left( \frac{2^4}{2} - \frac{2^5}{5} \right) \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left( \frac{2^4}{2} - \frac{2^5}{5} \right) \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left( \frac{2^4}{2} - \frac{2^5}{5} \right) = 2\pi \left( 8 - \frac{32}{5} \right) \end{aligned}$$

## Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by  $y = 2x^2 - x^3$  and the x-axis.

Cylindrical shell: outer radius  $x$ ; height:  $2x^2 - x^3$ ; circumference:  $2\pi x$ ; infinitesimal volume:  $2\pi x(2x^2 - x^3)dx$ .

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[ \frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left( \frac{2^4}{2} - \frac{2^5}{5} \right) = 2\pi \left( 8 - \frac{32}{5} \right) = \frac{16}{5}\pi. \end{aligned}$$


## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and **the  $x$ -axis**.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about **the line  $x = 1$**  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.



## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius ? ; height:

? ; circumference: ? ;

infinitesimal volume: ? .



## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.


Cylindrical shell: outer radius  $x - 1$ ; height:

?

; circumference: ?

;

infinitesimal volume: ?



## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.


Cylindrical shell: outer radius  $x - 1$ ; height:

?

; circumference: ?

;

infinitesimal volume: ?



## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.


Cylindrical shell: outer radius  $x - 1$ ; height:

$$|x^3 - 3x^2 + 2x|$$

; circumference: ?

;

infinitesimal volume: ?




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference: ? ;

infinitesimal volume: ? .




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference: ?

infinitesimal volume: ?




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume: ?




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;

infinitesimal volume: ?




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;

infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;

**infinitesimal** volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .



## Example (Rotated About a Line Other Than the $y$ -axis)


Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;

infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$V = \int_{?}^{?} 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$V = \int_{?}^{?} 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$V = \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (? ) dx \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (? ) dx \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + \frac{4x^4}{4} - \frac{5x^3}{3} + 2x^2 \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + ? - ? + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + ? - ? + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + x^4 - ? + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + x^4 - ? + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + ? \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \end{aligned}$$




## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .




$$\begin{aligned}
 V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left( \left( -\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left( -\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right)
 \end{aligned}$$

## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .




$$\begin{aligned}
 V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left( \left( -\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left( -\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right)
 \end{aligned}$$

## Example (Rotated About a Line Other Than the $y$ -axis)

Find the volume obtained by rotating about the line  $x = 1$  the region to the right of  $x = 1$  bounded by  $y = x^3 - 3x^2 + 2x$  and the  $x$ -axis.

Cylindrical shell: outer radius  $x - 1$ ; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$ ; circumference:  $2\pi(x - 1)$ ;  
infinitesimal volume:  $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$ .



$$\begin{aligned}
 V &= \int_1^2 2\pi(x-1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[ -\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left( \left( -\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left( -\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right) = \frac{4}{15}\pi.
 \end{aligned}$$

	Rotate about ...	
	... a horizontal line	... a vertical line
$y$ is a function of $x$	Cross-sections $\int \cdot dx$	Cylindrical shells $\int \cdot dx$
$x$ is a function of $y$	Cylindrical shells $\int \cdot dy$	Cross-sections $\int \cdot dy$

- $\int \cdot dx$  means integrate with respect to  $x$ .
- $\int \cdot dy$  means integrate with respect to  $y$ .
- Some equations express  $y$  as a function of  $x$  and  $x$  as a function of  $y$ . In such cases, you may use either method.