

Precalculus

Euler's formula memorization

Todor Milev

2019

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + \frac{(-1)^n x^{2n+1}}{(2n+1)!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + \frac{(-1)^n x^{2n}}{(2n)!} + \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} + \frac{z^5}{5!} + \dots$$

Rearrange.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix + \frac{(ix)^2}{2!} + \frac{(ix)^3}{3!} + \frac{(ix)^4}{4!} + \frac{(ix)^5}{5!} + \dots$$

Rearrange. **Plug-in** $z = ix$.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$. Multiply $\sin x$ by i .



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$. Multiply $\sin x$ by i . **Add to get**
 $e^{ix} = \cos x + i \sin x.$



Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

All trigonometric formulas can be easily derived using the above formulas.

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \quad .\end{aligned}$$

Proof.

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \quad .\end{aligned}$$

Proof.

$$e^{i(x+y)} =$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \quad .\end{aligned}$$

Proof.

$$e^{i(x+y)} = \cos(x+y) + i \sin(x+y)$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \quad .\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Compare coefficient in front of i and

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \sin y \cos x \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(x+y)} &= \cos(x+y) + i \sin(x+y) \\ e^{ix} e^{iy} &= \cos(x+y) + i \sin(x+y) \\ (\cos x + i \sin x)(\cos y + i \sin y) &= \cos(x+y) + i \sin(x+y) \\ \cos x \cos y - \sin x \sin y + i(\sin x \cos y + \sin y \cos x) &= \cos(x+y) + i \sin(x+y)\end{aligned}$$

Compare coefficient in front of i and **remaining terms** to get the desired equalities.



Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$1 = e^0$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = \end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = \end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned} 1 &= e^0 \\ &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned}
 1 &= e^0 \\
 &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\
 &= (\cos x + i \sin x)(\cos x - i \sin x) =
 \end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned}
 1 &= e^0 \\
 &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\
 &= (\cos x + i \sin x)(\cos x - i \sin x) = \cos^2 x - i^2 \sin^2 x
 \end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\sin^2 x + \cos^2 x = 1$$

Proof.

$$\begin{aligned}
 1 &= e^0 \\
 &= e^{ix-ix} = e^{ix} e^{-ix} = (\cos x + i \sin x)(\cos(-x) + i \sin(-x)) \\
 &= (\cos x + i \sin x)(\cos x - i \sin x) = \cos^2 x - i^2 \sin^2 x \\
 &= \cos^2 x + \sin^2 x .
 \end{aligned}$$



Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

$$e^{i(2x)} =$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

$$e^{i(2x)} = \cos(2x) + i \sin(2x)$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)(\cos x + i \sin x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \quad .\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x$, $\cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Compare coefficient in front of i and

Trigonometric Identities Revisited

- $e^{ix} = \cos x + i \sin x$ (Euler's Formula).
- $e^{ix} e^{iy} = e^{ix+iy} = e^{i(x+y)}$ (exponentiation rule: valid for \mathbb{C}).
- $e^0 = 1$ (exponentiation rule).
- $\sin(-x) = -\sin x, \cos(-x) = \cos x$ (easy to remember).

Example

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x.\end{aligned}$$

Proof.

$$\begin{aligned}e^{i(2x)} &= \cos(2x) + i \sin(2x) \\ e^{ix} e^{ix} &= \cos(2x) + i \sin(2x) \\ (\cos x + i \sin x)^2 &= (\cos x + i \sin x)(\cos x + i \sin x) = \cos(2x) + i \sin(2x) \\ \cos^2 x - \sin^2 x + i(2 \sin x \cos x) &= \cos(2x) + i \sin(2x)\end{aligned}$$

Compare coefficient in front of i and **remaining terms** to get the desired equalities.

