Calculus I

Reference: the Chain Rule statement and notation

Todor Milev

2019

• What is the derivative of $f(x) = \sqrt{x^2 + 1}$?

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- y = g(u) = ? .
- u = h(x) = ?
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- y = g(u) = ? .
- u = h(x) = ?
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- u = h(x) =?
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- u = h(x) = ?
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- u = h(x) = ?
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then y = f(x) = g(h(x)) =

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of *g* and *h*:
- g'(u) = ?
- h'(x) = ?.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of *g* and *h*:
- g'(u) = ?
- h'(x) = ?.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of *g* and *h*:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = ?.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of g and h:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = ?.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of *g* and *h*:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = 2x.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of g and h:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = 2x.
- It would be nice if we could find the derivative of f in terms of the derivatives of y and u.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of g and h:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = 2x.
- It would be nice if we could find the derivative of f in terms of the derivatives of y and u.
- It turns out that the derivative of the composition $g \circ h$ is the product of the derivative of g and the derivative of h.

- What is the derivative of $f(x) = \sqrt{x^2 + 1}$?
- The Power Rule doesn't tell us how to find the derivative.
- f is a composite function $g \circ h$:
- $y = g(u) = \sqrt{u}$.
- $u = h(x) = x^2 + 1$.
- Then $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$.
- We know the derivatives of *g* and *h*:
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$.
- h'(x) = 2x.
- It would be nice if we could find the derivative of f in terms of the derivatives of y and u.
- It turns out that the derivative of the composition $g \circ h$ is the product of the derivative of g and the derivative of h.
- This important fact is called the Chain Rule.

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

$$f'(x) = g'(h(x)) \cdot h'(x)$$
 (notation 1)

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

Let h be differentiable at x and let g be a differentiable at h(x). Then the composite function $f = g \circ h$ is differentiable at x and f' is given by the product

$$f'(x) = g'(h(x)) \cdot h'(x)$$

(notation 1) equivalently:

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

$$f'(x) = g'(h(x)) \cdot h'(x)$$
 (notation 1)
equivalently:
 $f'(x) = (g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

$$f'(x) = g'(h(x)) \cdot h'(x) \qquad (notation 1)$$

$$equivalently:$$

$$f'(x) = (g(u))' = g'(u)u' \qquad where u = h(x) \quad (notation 2)$$

$$\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx} \qquad where y = g(u) \quad (notation 3)$$

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

$$f'(x) = g'(h(x)) \cdot h'(x)$$
 (notation 1) equivalently:
 $f'(x) = (g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2) $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u}\frac{\mathrm{d}u}{\mathrm{d}x}$ where $y = g(u)$ (notation 3).

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

$$f'(x) = g'(h(x)) \cdot h'(x)$$
 (notation 1) equivalently:
$$f'(x) = (g(u))' = g'(u)u'$$
 where $u = h(x)$ (notation 2)
$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$
 where $y = g(u)$ (notation 3) .

Let g and h be functions. Recall that the composite function $f = g \circ h$ is defined via f(x) = g(h(x)).

Theorem

Let h be differentiable at x and let g be a differentiable at h(x). Then the composite function $f = g \circ h$ is differentiable at x and f' is given by the product

$$f'(x) = g'(h(x)) \cdot h'(x) \qquad \qquad \text{(notation 1)}$$

$$equivalently:$$

$$f'(x) = (g(u))' = g'(u)u' \qquad \text{where } u = h(x) \quad \text{(notation 2)}$$

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}u}\frac{\mathrm{d}u}{\mathrm{d}x} \qquad \text{where } y = g(u) \quad \text{(notation 3)} \quad .$$

The last equality uses the Leibniz notation (due to G. Leibniz (1646-1716)).

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{dx} \frac{du}{dx}$ where $y = g(u)$ (notation 3).

 As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}$ where $y = g(u)$ (notation 3).

 The three notations are all accepted and can be used interchangeably.

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}$ where $y = g(u)$ (notation 3).

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}$ where $y = g(u)$ (notation 3).

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}$ where $y = g(u)$ (notation 3).

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.
- There are additional notations (not covered here) used in practice.

$$(g(h(x)))' = g'(h(x)) \cdot h'(x)$$
 (notation 1)
 $(g(u))' = g'(u)u'$ where $u = h(x)$ (notation 2)
 $\frac{dy}{dx} = \frac{dy}{du}\frac{du}{dx}$ where $y = g(u)$ (notation 3).

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.
- There are additional notations (not covered here) used in practice.
- Whenever in doubt about derivative notation, if possible, request clarification.