

Calculus I

Implicit derivatives, related rates

Todor Milev

2019

Outline

1 Implicit Differentiation

Outline

1 Implicit Differentiation

2 Related Rates

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.


Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.
- Implicitly it gives two functions:




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.
- Implicitly it gives two functions: $y = \sqrt{1 - x^2}$ and




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.
- Implicitly it gives two functions: $y = \sqrt{1 - x^2}$ and $y = -\sqrt{1 - x^2}$.




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.
- Implicitly it gives two functions: $y = \sqrt{1 - x^2}$ and $y = -\sqrt{1 - x^2}$.
- How do we differentiate these functions?




Implicit Differentiation

- So far, we have seen functions with formulas that express one variable explicitly in terms of the other.
- $y = \sqrt{x^3 + 1}$, $y = x \sin x$, etc.
- Some functions are given implicitly by a relation between x and y .
- $x^2 + y^2 = 1$ isn't the equation of any one function.
- Implicitly it gives two functions: $y = \sqrt{1 - x^2}$ and $y = -\sqrt{1 - x^2}$.
- How do we differentiate these functions?
- Differentiate both sides with respect to x , and then solve for y' .




Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Example


Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

+ ?
= ?

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx}(25)$$

$$2(x - 1) \frac{d}{dx}(x - 1) + ? = ?$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$
$$2(x - 1) \frac{d}{dx} (x - 1) + \text{?} = \text{?}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx}(25)$$

$$2(x - 1) \frac{d}{dx}(x - 1) + 2(y + 2) \frac{d}{dx}(y + 2) = ?$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx}(25)$$
$$2(x - 1) \frac{d}{dx}(x - 1) + 2(y + 2) \frac{d}{dx}(y + 2) = ?$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx}(25)$$

$$2(x - 1) \frac{d}{dx}(x - 1) + 2(y + 2) \frac{d}{dx}(y + 2) = 0$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:


$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(x - 1)(?) + 2(y + 2) \left(? \right) = 0$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:


$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(x - 1)(1) + 2(y + 2) \left(? \right) = 0$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\begin{aligned} \frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) &= \frac{d}{dx}(25) \\ 2(x - 1) \frac{d}{dx}(x - 1) + 2(y + 2) \frac{d}{dx}(y + 2) &= 0 \\ 2(x - 1)(1) + 2(y + 2) \left(? \right) &= 0 \end{aligned}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.




Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\begin{aligned}\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) &= \frac{d}{dx}(25) \\ 2(x - 1) \frac{d}{dx}(x - 1) + 2(y + 2) \frac{d}{dx}(y + 2) &= 0 \\ 2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) &= 0\end{aligned}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$


$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(1 - x)$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$


$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(\mathbf{x} - \mathbf{1})(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(\mathbf{1} - \mathbf{x})$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$


$$2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(1 - x)$$

$$\frac{dy}{dx} = \frac{1 - x}{y + 2}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Plug in $(-2, 2)$:


$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2}$$

Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\begin{aligned} \frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) &= \frac{d}{dx} (25) \\ 2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) &= 0 \\ 2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) &= 0 \\ 2(y + 2) \left(\frac{dy}{dx} \right) &= 2(1 - x) \\ \frac{dy}{dx} &= \frac{1 - x}{y + 2} \end{aligned}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Plug in $(-2, 2)$:


$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2}$$

Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\begin{aligned} \frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) &= \frac{d}{dx} (25) \\ 2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) &= 0 \\ 2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) &= 0 \\ 2(y + 2) \left(\frac{dy}{dx} \right) &= 2(1 - x) \\ \frac{dy}{dx} &= \frac{1 - x}{y + 2} \end{aligned}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Plug in $(-2, 2)$:


$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2} = \frac{3}{4}$$

Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\begin{aligned} \frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) &= \frac{d}{dx} (25) \\ 2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) &= 0 \\ 2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) &= 0 \\ 2(y + 2) \left(\frac{dy}{dx} \right) &= 2(1 - x) \\ \frac{dy}{dx} &= \frac{1 - x}{y + 2} \end{aligned}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Plug in $(-2, 2)$:

$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2} = \frac{3}{4}$$

Point-slope form:

$$y - 2 = \frac{3}{4}(x + 2)$$

Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$


$$2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(1 - x)$$

$$\frac{dy}{dx} = \frac{1 - x}{y + 2}$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(1 - x)$$

$$\frac{dy}{dx} = \frac{1 - x}{y + 2}$$

Plug in $(-2, 2)$:


$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2} = \frac{3}{4}$$

Point-slope form:

$$y - 2 = \frac{3}{4}(x + 2)$$

Example

Find an equation of the tangent line to $(x - 1)^2 + (y + 2)^2 = 25$ at $(-2, 2)$.



Find $\frac{dy}{dx}$, given $(x - 1)^2 + (y + 2)^2 = 25$:

$$\frac{d}{dx} \left((x - 1)^2 \right) + \frac{d}{dx} \left((y + 2)^2 \right) = \frac{d}{dx} (25)$$

$$2(x - 1) \frac{d}{dx} (x - 1) + 2(y + 2) \frac{d}{dx} (y + 2) = 0$$

$$2(x - 1)(1) + 2(y + 2) \left(\frac{dy}{dx} \right) = 0$$

$$2(y + 2) \left(\frac{dy}{dx} \right) = 2(1 - x)$$

$$\frac{dy}{dx} = \frac{1 - x}{y + 2}$$

Plug in $(-2, 2)$:


$$\frac{dy}{dx} = \frac{1 - (-2)}{2 + 2} = \frac{3}{4}$$

Point-slope form:

$$y - 2 = \frac{3}{4}(x + 2)$$


(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$?




(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? **Yes.**



(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.




We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.




We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.

- Elementary algorithm: fix large rectangle.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.



We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ **have different sign** then H must become zero somewhere on the segment between P and Q .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign **then H must become zero somewhere on the segment between P and Q .**

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

Set $H(x, y) = x^2 + 2y^2 - 1$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- **For each triangle:**
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.




Illustrate the algorithm for:

$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.

(Elementary Computer algorithm for sketching graphs)

Let H -continuous; is there simple algorithm to sketch $H(x, y) = 0$? Yes.




Illustrate the algorithm for:

$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
 - Fix two corners $P(x_P, y_P)$ and $Q(x_Q, y_Q)$.
 - If $H(x_P, y_P)$ and $H(x_Q, y_Q)$ have different sign then H must become zero somewhere on the segment between P and Q .
 - Select a point between P and Q and “guess” that H is zero there.
 - In our implementation, we select the midpoint (i.e., $\frac{1}{2}P + \frac{1}{2}Q$).
 - Connect the selected pts. for each triangle.
 - Repeat for ever finer grid.


Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$

Example




Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$

Example



?


Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$=?$$

Example




Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = ?$$

Example




Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = ?$$

Example




Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (\text{?}) = \text{?} \cos(2x) + y^2 \text{?}$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = ? \quad \cos(2x) + y^2?$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = ? \cos(2x) + y^2 ?$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2?$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 \text{?}$$

Example



Find y' as an expression of x and y .


$$\sin(2(x + y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$


$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x + y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x)?$$

Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$


$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x + y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x)?$$

Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$


$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x + y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) \text{ ?}$$

Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$


$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x + y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) \quad 2$$

Example



Find y' as an expression of x and y .

$$\sin(2(x + y)) = y^2 \cos(2x).$$


$$\frac{d}{dx}(\sin(2(x + y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x + y)) \frac{d}{dx}(2(x + y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x + y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x + y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$


$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$


$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$


$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

$$y' \cos(2(x+y)) - yy' \cos(2x) = -\cos(2(x+y)) - y^2 \sin(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$


$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

$$y' \cos(2(x+y)) - yy' \cos(2x) = -\cos(2(x+y)) - y^2 \sin(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$


$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

$$y' \cos(2(x+y)) - yy' \cos(2x) = -\cos(2(x+y)) - y^2 \sin(2x)$$

$$y'(\cos(2(x+y)) - y \cos(2x)) = -\cos(2(x+y)) - y^2 \sin(2x)$$

Example



Find y' as an expression of x and y .

$$\sin(2(x+y)) = y^2 \cos(2x).$$

$$\frac{d}{dx}(\sin(2(x+y))) = \frac{d}{dx}(y^2 \cos(2x))$$

$$\cos(2(x+y)) \frac{d}{dx}(2(x+y)) = \frac{d}{dx}(y^2) \cos(2x) + (y^2) \frac{d}{dx}(\cos(2x))$$

$$\cos(2(x+y)) (2 + 2y') = 2yy' \cos(2x) + y^2 (-\sin(2x)) \frac{d}{dx}(2x)$$

$$2 \cos(2(x+y)) (1 + y') = 2yy' \cos(2x) - y^2 \sin(2x) 2$$

$$\cos(2(x+y)) + y' \cos(2(x+y)) = yy' \cos(2x) - y^2 \sin(2x)$$

$$y' \cos(2(x+y)) - yy' \cos(2x) = -\cos(2(x+y)) - y^2 \sin(2x)$$

$$y'(\cos(2(x+y)) - y \cos(2x)) = -\cos(2(x+y)) - y^2 \sin(2x)$$

$$y' = \frac{-\cos(2(x+y)) - y^2 \sin(2x)}{\cos(2(x+y)) - y \cos(2x)}.$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3 y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right)$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3 y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx} (x^3) y^3 - x^3 \frac{d}{dx} (y^3)}{(y^3)^2}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3 y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx} (x^3) y^3 - x^3 \frac{d}{dx} (y^3)}{(y^3)^2}$$

$$= -\frac{(\textcolor{red}{?}) y^3 - x^3 (\textcolor{red}{?})}{y^6}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned} y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2} \\ &= -\frac{(3x^2)y^3 - x^3(?) }{y^6} \end{aligned}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3 y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx} (x^3) y^3 - x^3 \frac{d}{dx} (y^3)}{(y^3)^2}$$

$$= -\frac{(3x^2)y^3 - x^3(\text{?})}{y^6}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned} y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2} \\ &= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} \end{aligned}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3 y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned} y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx} (x^3) y^3 - x^3 \frac{d}{dx} (y^3)}{(y^3)^2} \\ &= -\frac{(3x^2)y^3 - x^3(3y^2 y')}{y^6} = -\frac{3x^2 y^3 - 3x^3 y^2 \left(-\frac{x^3}{y^3} \right)}{y^6} \end{aligned}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned} y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2} \\ &= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} = -\frac{3x^2y^3 - 3x^3y^2\left(-\frac{x^3}{y^3}\right)}{y^6} \\ &= -\frac{3x^2(y^3 + \frac{x^4}{y})}{y^6} \end{aligned}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned} y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2} \\ &= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} = -\frac{3x^2y^3 - 3x^3y^2\left(-\frac{x^3}{y^3}\right)}{y^6} \\ &= -\frac{3x^2\left(y^3 + \frac{x^4}{y}\right)}{y^6} = -\frac{3x^2\left(\frac{y^4 + x^4}{y}\right)}{y^6} \end{aligned}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2}$$

$$= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} = -\frac{3x^2y^3 - 3x^3y^2\left(-\frac{x^3}{y^3}\right)}{y^6}$$

$$= -\frac{3x^2(y^3 + \frac{x^4}{y})}{y^6} = -\frac{3x^2\left(\frac{y^4 + x^4}{y}\right)}{y^6}$$

$$= -\frac{3x^2(y^4 + x^4)}{y^7}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$y'' = \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2}$$

$$= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} = -\frac{3x^2y^3 - 3x^3y^2\left(-\frac{x^3}{y^3}\right)}{y^6}$$

$$= -\frac{3x^2\left(y^3 + \frac{x^4}{y}\right)}{y^6} = -\frac{3x^2\left(\frac{y^4 + x^4}{y}\right)}{y^6}$$

$$= -\frac{3x^2(y^4 + x^4)}{y^7} = -\frac{3x^2(16)}{y^7}$$

Example

Let $x^4 + y^4 = 16$. Find y'' .

$$4x^3 + 4y^3y' = 0$$

$$y' = -\frac{x^3}{y^3}.$$

$$\begin{aligned}y'' &= \frac{d}{dx} \left(-\frac{x^3}{y^3} \right) = -\frac{\frac{d}{dx}(x^3) y^3 - x^3 \frac{d}{dx}(y^3)}{(y^3)^2} \\&= -\frac{(3x^2)y^3 - x^3(3y^2y')}{y^6} = -\frac{3x^2y^3 - 3x^3y^2\left(-\frac{x^3}{y^3}\right)}{y^6} \\&= -\frac{3x^2\left(y^3 + \frac{x^4}{y}\right)}{y^6} = -\frac{3x^2\left(\frac{y^4 + x^4}{y}\right)}{y^6} \\&= -\frac{3x^2(y^4 + x^4)}{y^7} = -\frac{3x^2(16)}{y^7} = -48\frac{x^2}{y^7}.\end{aligned}$$

Related Rates

- Suppose we are pumping a balloon with air.
- The balloon's volume is increasing.
- The balloon's radius is increasing.
- The rates of increase of these quantities are related to one another.

Related Rates

- Suppose we are pumping a balloon with air.
- The balloon's volume is increasing.
- The balloon's radius is increasing.
- The rates of increase of these quantities are related to one another.
- It is easier to measure the rate of increase of volume.

Related Rates

- Suppose we are pumping a balloon with air.
- The balloon's volume is increasing.
- The balloon's radius is increasing.
- The rates of increase of these quantities are related to one another.
- It is easier to measure the rate of increase of volume.
- In a related rates problem, we compute the rate of change of one quantity in terms of the rate of change of another (which may be more easily measured).

Related Rates

- Suppose we are pumping a balloon with air.
- The balloon's volume is increasing.
- The balloon's radius is increasing.
- The rates of increase of these quantities are related to one another.
- It is easier to measure the rate of increase of volume.
- In a related rates problem, we compute the rate of change of one quantity in terms of the rate of change of another (which may be more easily measured).
- Procedure:
 - 1 Find an equation relating the two quantities.
 - 2 Use the Chain Rule to differentiate both sides with respect to time.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given:
- Unknown:

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.
- **Given:**
- **Unknown:**

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
- Unknown:

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
- **Unknown:**

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. **How fast is the radius of the balloon increasing when the diameter is 50 cm?**

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
- **Unknown: $\frac{dr}{dt}$ when $r = 25 \text{ cm}$.**

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
 - Let r denote its radius.
 - Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
 - Unknown: $\frac{dr}{dt}$ when $r = 25 \text{ cm}$.
- 1 Find an equation relating the two quantities.
 - 2 Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
- Unknown: $\frac{dr}{dt}$ when $r = 25 \text{ cm}$.

1 Find an equation relating the two quantities.

2 Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

$$V = \frac{4}{3}\pi r^3$$

- Let V denote the balloon's volume.
- Let r denote its radius.
- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.
- Unknown: $\frac{dr}{dt}$ when $r = 25 \text{ cm}$.

- Find an equation relating the two quantities.
- Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

$$V = \frac{4}{3}\pi r^3$$

- Let r denote its radius.

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

- Find an equation relating the two quantities.

- Use the Chain Rule to **differentiate both sides**.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

$$V = \frac{4}{3}\pi r^3$$

- Let r denote its radius.

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

- Find an equation relating the two quantities.

- Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

$$V = \frac{4}{3}\pi r^3$$

- Let r denote its radius.

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

$$\frac{dV}{dt} = ? \quad \frac{dr}{dt}$$

- Find an equation relating the two quantities.
- Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

$$V = \frac{4}{3}\pi r^3$$

- Let r denote its radius.

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

- Find an equation relating the two quantities.
- Use the Chain Rule to differentiate both sides.

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

- Let r denote its radius.

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

- Unknown:** $\frac{dr}{dt}$ when $r = 25$ cm.

- Find an equation relating the two quantities.

- Use the Chain Rule to differentiate both sides.

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$$

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

- Let r denote its radius.

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

- Find an equation relating the two quantities.

- Use the Chain Rule to differentiate both sides.

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$$

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

- Let r denote its radius.

- Given:** $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

- Find an equation relating the two quantities.

- Use the Chain Rule to differentiate both sides.

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi (25\text{cm})^2} 100 \frac{\text{cm}^3}{\text{s}}$$

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

- Let r denote its radius.

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

- Unknown: $\frac{dr}{dt}$ when $r = 25 \text{ cm}$.

- Find an equation relating the two quantities.

- Use the Chain Rule to differentiate both sides.

$$V = \frac{4}{3}\pi r^3$$

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$$

$$\frac{dr}{dt} = \frac{1}{4\pi (25\text{cm})^2} 100 \frac{\text{cm}^3}{\text{s}}$$

Example

Air is being pumped into a balloon such that its volume changes at a rate of $100 \text{ cm}^3/\text{s}$. How fast is the radius of the balloon increasing when the diameter is 50 cm?

- Let V denote the balloon's volume.

$$V = \frac{4}{3}\pi r^3$$

- Let r denote its radius.

$$\frac{dV}{dt} = \frac{d}{dt} \left(\frac{4}{3}\pi r^3 \right)$$

- Given: $\frac{dV}{dt} = 100 \text{ cm}^3/\text{s}$.

$$\frac{dV}{dt} = \frac{d}{dr} \left(\frac{4}{3}\pi r^3 \right) \frac{dr}{dt}$$

- Unknown: $\frac{dr}{dt}$ when $r = 25$ cm.

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$


- Find an equation relating the two quantities.

$$\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$$

- Use the Chain Rule to differentiate both sides.


$$\frac{dr}{dt} = \frac{1}{4\pi(25\text{cm})^2} 100 \frac{\text{cm}^3}{\text{s}} = \frac{1}{25\pi} \text{cm/s}$$

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: ?
- Unknown: ?


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: ?


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: ?


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. **How fast is the ladder top sliding down when the bottom is 6 ft from the wall?**

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

Example




10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

- Relationship b/n quantities.
- Differentiate (use Chain Rule).


Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.
- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example




10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

$$x^2 + y^2 = 10^2 = 100$$

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?


- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.


$$x^2 + y^2 = 10^2 = 100$$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = - \frac{x}{y} \frac{dx}{dt}$$

$$\frac{dy}{dt} = - \frac{?}{?} \cdot ?$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = - \frac{x}{y} \frac{dx}{dt}$$

$$\frac{dy}{dt} = - \frac{?}{?} \cdot ?$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given:** $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = - \frac{x}{y} \frac{dx}{dt}$$

$$\frac{dy}{dt} = - \frac{?}{?} \cdot 1 \text{ ft/s}$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = - \frac{x}{y} \frac{dx}{dt}$$

$$\frac{dy}{dt} = - \frac{?}{?} \cdot 1 \text{ ft/s}$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$x^2 + y^2 = 10^2 = 100$$


$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} = -\frac{x}{y} \frac{dx}{dt}$$

$$\frac{dy}{dt} = -\frac{6 \text{ ft}}{?} \cdot 1 \text{ ft/s}$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example




10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.

$$\begin{aligned}
 x^2 + y^2 &= 10^2 = 100 \\
 2x \frac{dx}{dt} + 2y \frac{dy}{dt} &= 0 \\
 \frac{dy}{dt} &= -\frac{x}{y} \frac{dx}{dt} \\
 \frac{dy}{dt} &= -\frac{6 \text{ ft}}{\text{?}} \cdot 1 \text{ ft/s}
 \end{aligned}$$

- Relationship b/n quantities.
- Differentiate (use Chain Rule).

Example




10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.
- Pythagorean Theorem:**
 $y = \sqrt{10^2 - 6^2} = 8.$
- Relationship b/n quantities.
- Differentiate (use Chain Rule).

$$\begin{aligned}
 x^2 + y^2 &= 10^2 = 100 \\
 2x \frac{dx}{dt} + 2y \frac{dy}{dt} &= 0 \\
 \frac{dy}{dt} &= -\frac{x}{y} \frac{dx}{dt} \\
 \frac{dy}{dt} &= -\frac{6 \text{ ft}}{8 \text{ ft}} \cdot 1 \text{ ft/s}
 \end{aligned}$$

Example




10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.
- Pythagorean Theorem:
 $y = \sqrt{10^2 - 6^2} = 8$.
- Relationship b/n quantities.
- Differentiate (use Chain Rule).

$$\begin{aligned}
 x^2 + y^2 &= 10^2 = 100 \\
 2x \frac{dx}{dt} + 2y \frac{dy}{dt} &= 0 \\
 \frac{dy}{dt} &= -\frac{x}{y} \frac{dx}{dt} \\
 \frac{dy}{dt} &= -\frac{6 \text{ ft}}{8 \text{ ft}} \cdot 1 \text{ ft/s} \\
 &= -3/4 \text{ ft/s.}
 \end{aligned}$$

Example



10 ft ladder rests against a vertical wall. The bottom of the ladder slides away from the wall at a rate of 1 ft/s. How fast is the ladder top sliding down when the bottom is 6 ft from the wall?

- Let y = dist. from top to ground.
- Let x = dist. from bottom to wall.
- Given: $\frac{dx}{dt} = 1$ ft/s.
- Unknown: $\frac{dy}{dt}$ when $x = 6$ ft.
- Pythagorean Theorem:
 $y = \sqrt{10^2 - 6^2} = 8$.
- Relationship b/n quantities.
- Differentiate (use Chain Rule).

$$\begin{aligned}
 x^2 + y^2 &= 10^2 = 100 \\
 2x \frac{dx}{dt} + 2y \frac{dy}{dt} &= 0 \\
 \frac{dy}{dt} &= -\frac{x}{y} \frac{dx}{dt} \\
 \frac{dy}{dt} &= -\frac{6 \text{ ft}}{8 \text{ ft}} \cdot 1 \text{ ft/s} \\
 &= -3/4 \text{ ft/s.}
 \end{aligned}$$

Therefore the top of the ladder is **falling** at a rate of 3/4 ft/s.