

## Precalculus

# Equations formed by setting trigonometric sum equal to 0

Todor Milev

2019

## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

Proof.



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

Proof.

?

$$= \cos(\alpha + \beta)$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\cos \alpha \cos \beta - \sin \alpha \sin \beta = \cos(\alpha + \beta)$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}\textcolor{red}{?} &= \textcolor{red}{\cos(\alpha - \beta)} \\ \cos \alpha \cos \beta - \sin \alpha \sin \beta &= \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}\cos \alpha \cos \beta + \sin \alpha \sin \beta &= \cos(\alpha - \beta) \\ \cos \alpha \cos \beta - \sin \alpha \sin \beta &= \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad \cos \alpha \cos \beta + \sin \alpha \sin \beta &= \cos(\alpha - \beta) \\ \cos \alpha \cos \beta - \sin \alpha \sin \beta &= \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad \cos \alpha \cos \beta + \sin \alpha \sin \beta &= \cos(\alpha - \beta) \\ \cos \alpha \cos \beta - \sin \alpha \sin \beta &= \cos(\alpha + \beta) \\ \quad ? &= \cos(\alpha - \beta) + \cos(\alpha + \beta)\end{aligned}$$





## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad & \cos \alpha \cos \beta + \sin \alpha \sin \beta = \cos(\alpha - \beta) \\ & \cos \alpha \cos \beta - \sin \alpha \sin \beta = \cos(\alpha + \beta) \\ & \quad ? = \cos(\alpha - \beta) + \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} &= \cos(\alpha - \beta) \\ \cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} &= \cos(\alpha + \beta) \\ 2 \cos \alpha \cos \beta &= \cos(\alpha - \beta) + \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad & \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} = \cos(\alpha - \beta) \\ & \cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} = \cos(\alpha + \beta) \\ & 2 \cos \alpha \cos \beta = \cos(\alpha - \beta) + \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\sin \alpha \sin \beta = \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta))$$

$$\cos \alpha \cos \beta = \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta))$$

$$\sin \alpha \cos \beta = \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))$$

## Proof.

$$\begin{aligned}
 &+ \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} = \cos(\alpha - \beta) \\
 &\cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} = \cos(\alpha + \beta) \\
 &\quad \quad \quad 2 \cos \alpha \cos \beta = \cos(\alpha - \beta) + \cos(\alpha + \beta)
 \end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{aligned}+ \quad \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} &= \cos(\alpha - \beta) \\ \cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} &= \cos(\alpha + \beta) \\ \textcolor{red}{2} \cos \alpha \cos \beta &= \cos(\alpha - \beta) + \cos(\alpha + \beta)\end{aligned}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}
 \sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\
 \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\
 \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))
 \end{aligned}$$

## Proof.

$$\begin{array}{rcl}
 + & \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} & = \cos(\alpha - \beta) \\
 & \cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} & = \cos(\alpha + \beta) \\
 & \hline
 & 2 \cos \alpha \cos \beta & = \cos(\alpha - \beta) + \cos(\alpha + \beta) \\
 - & \cos \alpha \cos \beta + \sin \alpha \sin \beta & = \cos(\alpha - \beta) \\
 & \cos \alpha \cos \beta - \sin \alpha \sin \beta & = \cos(\alpha + \beta) \\
 & \hline
 & 2 \sin \alpha \sin \beta & = \cos(\alpha - \beta) - \cos(\alpha + \beta)
 \end{array}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

## Proof.

$$\begin{array}{rcll} + & \cos \alpha \cos \beta + \cancel{\sin \alpha \sin \beta} & = & \cos(\alpha - \beta) \\ & \cos \alpha \cos \beta - \cancel{\sin \alpha \sin \beta} & = & \cos(\alpha + \beta) \\ & 2 \cos \alpha \cos \beta & = & \cos(\alpha - \beta) + \cos(\alpha + \beta) \\ \hline - & \cos \alpha \cos \beta + \sin \alpha \sin \beta & = & \cos(\alpha - \beta) \\ & \cos \alpha \cos \beta - \sin \alpha \sin \beta & = & \cos(\alpha + \beta) \\ & 2 \sin \alpha \sin \beta & = & \cos(\alpha - \beta) - \cos(\alpha + \beta) \\ \hline + & \sin \alpha \cos \beta + \cos \alpha \sin \beta & = & \sin(\alpha + \beta) \\ & \sin \alpha \cos \beta - \cos \alpha \sin \beta & = & \sin(\alpha - \beta) \\ & 2 \sin \alpha \cos \beta & = & \sin(\alpha + \beta) + \sin(\alpha - \beta) \end{array}$$



## Proposition (Product to sum formulas)

$$\begin{aligned}\sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \\ \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha - \beta) + \cos(\alpha + \beta)) \\ \sin \alpha \cos \beta &= \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))\end{aligned}$$

- Product to sum formulas are used when integrating (a topic to be studied later/in another course).



## Proposition (Sum to product formulas)

$$\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$$

$$\sin \alpha - \sin \beta = 2 \sin \left( \frac{\alpha - \beta}{2} \right) \cos \left( \frac{\alpha + \beta}{2} \right)$$


$$\cos \alpha + \cos \beta = 2 \cos \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$$

$$\cos \alpha - \cos \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \sin \left( \frac{\alpha - \beta}{2} \right)$$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .

$$\sin(2x) + \sin(5x) = 0$$




Recall the formula  $\sin \alpha + \sin \beta = ?$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .

$$\sin(2x) + \sin(5x) = 0 \quad | \quad \text{use f-l-a}$$




Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .


$$\sin(2x) + \sin(5x) = 0 \quad | \quad \text{use f-l-a}$$



Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{2x + 5x}{2} \right) \cos \left( \frac{2x - 5x}{2} \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{2x + 5x}{2} \right) \cos \left( \frac{2x - 5x}{2} \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\begin{aligned} \sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{2x + 5x}{2} \right) \cos \left( \frac{2x - 5x}{2} \right) &= 0 \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( -\frac{3}{2}x \right) &= 0 \end{aligned}$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\begin{aligned} \sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{2x + 5x}{2} \right) \cos \left( \frac{2x - 5x}{2} \right) &= 0 \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( -\frac{3}{2}x \right) &= 0 \quad | \begin{array}{l} \cos \\ \text{is even} \end{array} \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) &= 0 \end{aligned}$$



Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\begin{aligned} \sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{2x + 5x}{2} \right) \cos \left( \frac{2x - 5x}{2} \right) &= 0 \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( -\frac{3}{2}x \right) &= 0 \quad | \text{ cos is even} \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) &= 0 \end{aligned}$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .





$$\sin(2x) + \sin(5x) = 0 \quad | \quad \text{use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$



$$\cos \left( \frac{3}{2}x \right) = 0$$

or

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$



$$\cos \left( \frac{3}{2}x \right) = 0$$

or

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = ?$$

$$\sin(2x) + \sin(5x) = 0 \quad | \quad \text{use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .




$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = ?$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$




$$\cos \left( \frac{3}{2}x \right) = 0$$

or

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = ?$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = ?$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

or


$$\cos \left( \frac{3}{2}x \right) = 0$$



Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$k$  – integer


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\begin{aligned}\sin \left( \frac{7}{2}x \right) &= 0 \\ \frac{7}{2}x &= k\pi \\ x &= \frac{2k\pi}{7}\end{aligned}$$

$$\begin{aligned}\sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) &= 0\end{aligned}$$

$k$  – integer


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\begin{aligned} \sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) &= 0 \end{aligned}$$

$$\begin{aligned} \sin \left( \frac{7}{2}x \right) &= 0 \\ \frac{7}{2}x &= k\pi \\ x &= \frac{2k\pi}{7} \end{aligned}$$

$k$  – integer

$$x = \dots, -\frac{2\pi}{7}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \dots$$


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions **in the interval  $[0, 2\pi)$** .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \cancel{\frac{14\pi}{7}}, \cancel{.}$$


or

$$\cos \left( \frac{3}{2}x \right) = 0$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{.}$$

or


$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\frac{3}{2}x = ?$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{.}$$

or


$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\frac{3}{2}x = ?$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{-\frac{2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{.}$$

or


$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\frac{3}{2}x = ?$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{-\frac{2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{.}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$


$$\frac{3}{2}x = ?$$



Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{\cdot}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{\cdot}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$


$$\frac{3}{2}x = \frac{\pi}{2} + k\pi$$

$k$  – integer

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\begin{aligned} \sin(2x) + \sin(5x) &= 0 \quad | \text{ use f-l-a} \\ 2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) &= 0 \end{aligned}$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{.}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{.}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$


$$\frac{3}{2}x = \frac{\pi}{2} + k\pi = \frac{(2k+1)\pi}{2}$$

$k$  – integer

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{\cdot}, \cancel{\frac{-2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{\cdot}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\textcolor{red}{2}x = \frac{\pi}{2} + k\pi = \frac{(2k+1)\pi}{2}$$


$k$  – integer

$$x = \frac{(2k+1)\pi}{\textcolor{red}{3}}$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{\dots}, \cancel{\frac{2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \cancel{\frac{14\pi}{7}}, \cancel{\dots}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\frac{3}{2}x = \frac{\pi}{2} + k\pi = \frac{(2k+1)\pi}{2}$$

$k$  – integer


$$x = \frac{(2k+1)\pi}{3}$$

$$x = \dots, -\frac{\pi}{3}, \frac{\pi}{3}, \frac{3\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \dots$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions **in the interval  $[0, 2\pi)$** .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use f-l-a}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$\sin \left( \frac{7}{2}x \right) = 0$$

$$\frac{7}{2}x = k\pi$$

$$x = \frac{2k\pi}{7}$$

$k$  – integer

$$x = \cancel{\cdot}, \cancel{\frac{2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \cancel{\frac{14\pi}{7}}, \cancel{\cdot}$$

or

$$\cos \left( \frac{3}{2}x \right) = 0$$

$$\frac{3}{2}x = \frac{\pi}{2} + k\pi = \frac{(2k+1)\pi}{2}$$

$k$  – integer


$$x = \frac{(2k+1)\pi}{3}$$

$$x = \cancel{\cdot}, \cancel{\frac{\pi}{3}}, \frac{\pi}{3}, \frac{3\pi}{3}, \frac{5\pi}{3}, \cancel{\frac{7\pi}{3}}, \cancel{\cdot}$$

Recall the formula  $\sin \alpha + \sin \beta = 2 \sin \left( \frac{\alpha + \beta}{2} \right) \cos \left( \frac{\alpha - \beta}{2} \right)$

## Example

Find all solutions in the interval  $[0, 2\pi)$ .



$$\sin(2x) + \sin(5x) = 0 \quad | \text{ use formula}$$

$$2 \sin \left( \frac{7}{2}x \right) \cos \left( \frac{3}{2}x \right) = 0$$

$$x = \cancel{\cdot}, \cancel{-\frac{2\pi}{7}}, 0, \frac{2\pi}{7}, \frac{4\pi}{7}, \frac{6\pi}{7}, \frac{8\pi}{7}, \frac{10\pi}{7}, \frac{12\pi}{7}, \frac{14\pi}{7}, \cancel{\cdot}$$

or

$$x = \cancel{\cdot}, \cancel{-\frac{\pi}{3}}, \frac{\pi}{3}, \frac{3\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}, \cancel{\cdot}$$

