

# Precalculus

## Graphs of trig functions; inverse trig

Todor Milev


2019

# Outline


## 1 Inverse Trigonometric Functions

- The arcsine function
- The arccosine function
- The arctangent and the remaining inverse trig functions
- Trigonometric Functions with Inverse Trig Arguments

# Inverse Trigonometric Functions




# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.

# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .

# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .


# Inverse Trigonometric Functions




- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .



# Inverse Trigonometric Functions



- $\sin x$  isn't one-to-one.
- It is if we restrict the domain to  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- Then it has an inverse function.
- We call it arcsin or  $\sin^{-1}$ .
- $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .





## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

## Observation

- $\arcsin y =$  *the appropriate angle* whose sine equals  $y$ .

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\textcolor{red}{?}\right) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin(?) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\text{?}\right) = \frac{1}{2}$ .

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\textcolor{red}{?}\right) = \frac{1}{2}.$

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}.$

## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*
- *Important: the output angle must lie in the interval  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ .*

## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$ .
- $-\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}$ .



## Observation

- $\arcsin y =$  *the appropriate angle whose sine equals  $y$ .*
- *Important: the output angle must lie in the interval  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .*


## Example

Find  $\arcsin\left(\frac{1}{2}\right)$ .

- $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$ .
- $-\frac{\pi}{2} \leq \frac{\pi}{6} \leq \frac{\pi}{2}$ .
- Therefore  $\arcsin\left(\frac{1}{2}\right) = \frac{\pi}{6}$ .

## Example


Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with **opposite side 1** and hypotenuse 3.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and **hypotenuse 3**.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled.



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .

- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .






## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side = ?



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side =  $\sqrt{3^2 - 1^2}$



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .


- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .
- Then  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right) = ?$



## Example

Find  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right)$ .

- Let  $\theta = \arcsin \left( \frac{1}{3} \right)$ , so  $\sin \theta = \frac{1}{3}$ .
- Draw a right triangle with opposite side 1 and hypotenuse 3.
- Let the angle  $\theta$  be as labeled. Then  $\sin \theta = \frac{1}{3}$  and so  $\theta = \arcsin \left( \frac{1}{3} \right)$ .
- Length of adjacent side  $= \sqrt{3^2 - 1^2} = \sqrt{8} = 2\sqrt{2}$ .
- Then  $\tan \left( \arcsin \left( \frac{1}{3} \right) \right) = \frac{1}{2\sqrt{2}}$ .



## Example

Find  $\arcsin(\sin(1.5))$ .

## Example

Find  $\arcsin(\sin(1.5))$ .

●  $\frac{\pi}{2} \approx ?$

## Example

Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .



## Example

Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .
- Therefore  $-\frac{\pi}{2} \leq 1.5 \leq \frac{\pi}{2}$ .



## Example

Find  $\arcsin(\sin(1.5))$ .

- $\frac{\pi}{2} \approx 1.57$ .
- Therefore  $-\frac{\pi}{2} \leq 1.5 \leq \frac{\pi}{2}$ .
- Therefore  $\arcsin(\sin 1.5) = 1.5$ .

## Example



Find  $\arcsin(\sin 2)$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .







## Example

Find  $\arcsin(\sin 2)$ .



- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .



## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  **$a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$**  for which  $\sin 2 = \sin a$ .





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = ?$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\text{Therefore } \arcsin(\sin 2) = \arcsin(\sin a)$$





## Example

Find  $\arcsin(\sin 2)$ .

- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  **$a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$**  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\begin{aligned} \text{Therefore } \arcsin(\sin 2) &= \arcsin(\sin a) \\ &= a \end{aligned}$$





## Example

Find  $\arcsin(\sin 2)$ .


- 2 is not between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .
- We need the angle  $a$  between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  for which  $\sin 2 = \sin a$ .

$$a = \pi - 2.$$

$$\begin{aligned} \text{Therefore } \arcsin(\sin 2) &= \arcsin(\sin a) \\ &= a = \pi - 2. \end{aligned}$$




## Important facts about arcsin:



- 1 Domain: ?
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .




## Important facts about arcsin:




- 1 Domain: ?
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 Range: ?
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 **Range: ?**
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .


## Important facts about arcsin:




- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .
- 3  $\arcsin x = y \Leftrightarrow \sin y = x$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ .
- 4  $\arcsin(\sin x) = x$  for  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .
- 5  $\sin(\arcsin x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$ .




- Same for  $\cos x$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called arccos or  $\cos^{-1}$ .




- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called  $\arccos$  or  $\cos^{-1}$ .








- Same for  $\cos x$ .
- Restrict the domain to  $[0, \pi]$ .
- The inverse is called arccos or  $\cos^{-1}$ .
- $\arccos(x) = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .




## Important facts about arccos:



- 1 Domain:
- 2 Range:
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:



1 Domain: ?

2 Range:


3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .

4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .

5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .


6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:




- 1 Domain:  $[-1, 1]$ .
- 2 Range:
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:




- 1 Domain:  $[-1, 1]$ .
- 2 **Range: ?**
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .

## Important facts about arccos:



- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[0, \pi]$ .
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .



## Important facts about arccos:



- 1 Domain:  $[-1, 1]$ .
- 2 Range:  $[0, \pi]$ .
- 3  $\arccos x = y \Leftrightarrow \cos y = x$  and  $0 \leq y \leq \pi$ .
- 4  $\arccos(\cos x) = x$  for  $0 \leq x \leq \pi$ .
- 5  $\cos(\arccos x) = x$  for  $-1 \leq x \leq 1$ .
- 6  $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$ .  
(The proof is similar to the proof of the formula for the derivative of  $\arcsin x$ .)

## Example

Find  $\arccos(\cos 4)$ .







## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .



## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .



## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .



## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .



## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .



## Example

Find  $\arccos(\cos 4)$ .



- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .



## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  **$a$  between 0 and  $\pi$**  for which  $\cos 4 = \cos a$ .





## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = ?$$







## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = \pi - (4 - \pi)$$





## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = \pi - (4 - \pi) = 2\pi - 4$$





## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = \pi - (4 - \pi) = 2\pi - 4$$

$$\text{Therefore } \arccos(\cos 4) = \arccos(\cos a)$$





## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = \pi - (4 - \pi) = 2\pi - 4$$

$$\begin{aligned} \text{Therefore } \arccos(\cos 4) &= \arccos(\cos a) \\ &= a \end{aligned}$$





## Example

Find  $\arccos(\cos 4)$ .

- 4 is not between 0 and  $\pi$ .
- We need the angle  $a$  between 0 and  $\pi$  for which  $\cos 4 = \cos a$ .

$$a = \pi - (4 - \pi) = 2\pi - 4$$


$$\begin{aligned} \text{Therefore } \arccos(\cos 4) &= \arccos(\cos a) \\ &= a = 2\pi - 4. \end{aligned}$$



## Example


The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed?

## Example



The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? **Assume earth is round** with radius 6371 km and that the ship sails along the shortest curved path.


## Example



The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with **radius** 6371 km and that the ship sails along the shortest curved path.



## Example




not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.

## Example




not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth.


## Example



The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.


## Example



The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that **the ship sails along the shortest curved path**.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

## Example




not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

## Example




The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with **radius 6371 km** and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r=6371\text{km}$$

## Example




The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r=6371\text{km}$$

$$d=?$$

## Example



The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.


- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km}$$



## Example



not to scale


The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km}$$

## Example




The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km}$$

## Example



not to scale


The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.


- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = ?$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.


- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .


$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos \left( \frac{r}{d} \right)$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$


$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos\left(\frac{r}{d}\right)$$

$$x = ?$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$

$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$


$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos\left(\frac{r}{d}\right)$$

$$x = r\alpha$$



## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$


$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos\left(\frac{r}{d}\right)$$

$$x = r\alpha = r \arccos\left(\frac{r}{d}\right)$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$


$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos \left( \frac{r}{d} \right)$$

$$x = r\alpha = r \arccos \left( \frac{r}{d} \right) = 6371 \text{ km} \arccos \left( \frac{6371 \text{ km}}{6371.01 \text{ km}} \right)$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$


$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$

$$\alpha = \arccos \left( \frac{r}{d} \right)$$

$$x = r\alpha = r \arccos \left( \frac{r}{d} \right) = 6371 \text{ km} \arccos \left( \frac{6371 \text{ km}}{6371.01 \text{ km}} \right)$$

## Example



not to scale

The horizon at sea is viewed at from a ship height of 10m. How far does the ship need to travel to reach the horizon just observed? Assume earth is round with radius 6371 km and that the ship sails along the shortest curved path.

- Let  $d$  be the distance from eyes of seaman to the center of earth.
- Let  $r$  be the radius of earth. Let  $\alpha$  be the indicated angle.
- Let the distance to the horizon be  $x$ .

$$r = 6371 \text{ km}$$


$$d = 6371 \text{ km} + 0.01 \text{ km} = 6371.01 \text{ km}$$

$$\cos \alpha = \frac{r}{d}$$


$$\alpha = \arccos \left( \frac{r}{d} \right)$$


$$x = r\alpha = r \arccos \left( \frac{r}{d} \right) = 6371 \text{ km} \arccos \left( \frac{6371 \text{ km}}{6371.01 \text{ km}} \right) \approx 11.29 \text{ km}$$

- $\tan x$  isn't one-to-one.




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .






- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .







- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- **Domain of  $\arctan$ : ?**
- Range of  $\arctan$ :




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- **Domain of  $\arctan$ :  $(-\infty, \infty)$ .**
- Range of  $\arctan$ :




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ : ?




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = ?$
- $\lim_{x \rightarrow -\infty} \arctan x =$



- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x =$



- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x = ?$




- $\tan x$  isn't one-to-one.
- Restrict the domain to  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- The inverse is called  $\tan^{-1}$  or  $\arctan$ .
- $\arctan x = y \Leftrightarrow \tan y = x$  and  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ .
- Domain of  $\arctan$ :  $(-\infty, \infty)$ .
- Range of  $\arctan$ :  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .
- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$ .
- $\lim_{x \rightarrow -\infty} \arctan x = -\frac{\pi}{2}$ .



## Example


Simplify the expression  $\cos(\arctan x)$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .
- Length of hypotenuse = ?



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .



## Example

Simplify the expression  $\cos(\arctan x)$ .


- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .
- Then  $\cos(\arctan x) = ?$



## Example

Simplify the expression  $\cos(\arctan x)$ .

- Let  $y = \arctan x$ , so  $\tan y = x$ .
- Draw a right triangle with opposite  $x$  and adjacent  $1$ .
- Length of hypotenuse =  $\sqrt{1^2 + x^2}$ .
- Then  $\cos(\arctan x) = \frac{1}{\sqrt{1+x^2}}$ .



The remaining inverse trigonometric functions aren't used as often:

$$\begin{aligned}y = \operatorname{arccsc} x \quad (|x| \geq 1) &\Leftrightarrow \csc y = x \quad \text{and} \quad y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right] \\y = \operatorname{arcsec} x \quad (|x| \geq 1) &\Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\y = \operatorname{arccot} x \quad (|x| \in \mathbb{R}) &\Leftrightarrow \cot y = x \quad \text{and} \quad y \in (0, \pi)\end{aligned}$$




The remaining inverse trigonometric functions aren't used as often:

$$\begin{aligned} y = \operatorname{arccsc} x \quad (|x| \geq 1) &\Leftrightarrow \csc y = x \quad \text{and} \quad y \in \left(0, \frac{\pi}{2}\right] \cup \left(\pi, \frac{3\pi}{2}\right] \\ y = \operatorname{arcsec} x \quad (|x| \geq 1) &\Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right) \\ y = \operatorname{arccot} x \quad (|x| \in \mathbb{R}) &\Leftrightarrow \cot y = x \quad \text{and} \quad y \in (0, \pi) \end{aligned}$$

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$


• Plot  $\sec x$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$


- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$


- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$

- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in ?$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and  
 $x \in [0, \frac{\pi}{2}) \cup [\pi, \frac{3\pi}{2})$ .
- $x \in [0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$




- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**



We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.


$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

We will however make use of  $\operatorname{arcsec} x$ : we discuss in detail its domain.

$$y = \operatorname{arcsec} x \quad (|x| \geq 1) \Leftrightarrow \sec y = x \quad \text{and} \quad y \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$$



- Plot  $\sec x$ .
- Restrict domain to make one-to-one: Two common choices:  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and  
 $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$ .
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  is good because the domain is easiest to remember: an interval without a point. **NOT our choice.**
- $x \in \left[0, \frac{\pi}{2}\right) \cup \left[\pi, \frac{3\pi}{2}\right)$  is good because  $\tan x$  is positive on both intervals, resulting in easier differentiation and integration formulas. **Our choice.**

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .

$$\sin(2 \arccos(x))$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ .

$$\sin(2 \arccos(x))$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\sin(2 \arccos(x))$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\sin(2 \arccos(x)) = \sin(2y)$$

$$| \text{ Set } y = \arccos x$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= ?\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$



## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to **rewrite the expression only using the cos function**.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right)\end{aligned} \quad \left| \begin{array}{l} \text{Set } y = \arccos x \\ \text{Express via } \sin y, \cos y \\ \text{Express } \sin y \text{ via } \cos y \end{array} \right.$$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y}\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$   
Express  $\sin y$  via  $\cos y$   
 $\sin y > 0$  because  
 $0 \leq y \leq \pi$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y}\end{aligned}$$

Set  $y = \arccos x$

Express via  $\sin y, \cos y$

Express  $\sin y$  via  $\cos y$

$\sin y > 0$  because

$0 \leq y \leq \pi$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y} \\ &= 2x \sqrt{1 - x^2}\end{aligned}$$

Set  $y = \arccos x$

Express via  $\sin y, \cos y$

Express  $\sin y$  via  $\cos y$

$\sin y > 0$  because

$0 \leq y \leq \pi$

use  $x = \cos y$

## Example

Rewrite  $\sin(2 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\sin(2 \arccos(x)) &= \sin(2y) \\ &= 2 \cos y \sin y \\ &= 2 \cos y \left( \pm \sqrt{1 - \cos^2 y} \right) \\ &= 2 \cos y \sqrt{1 - \cos^2 y} \\ &= 2x \sqrt{1 - x^2}\end{aligned}$$

Set  $y = \arccos x$   
Express via  $\sin y, \cos y$   
Express  $\sin y$  via  $\cos y$   
 $\sin y > 0$  because  
 $0 \leq y \leq \pi$   
use  $x = \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .

$$\cos(3 \arccos(x))$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ .  
To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ .

$$\cos(3 \arccos(x))$$



## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x))$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) \quad \Big| \quad y = \arccos x$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) = \cos(2y + y) \quad \bigg| \quad y = \arccos x$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\cos(3 \arccos(x)) = \cos(3y) = \cos(2y + y) \\ = ?$$

$y = \arccos x$   
Angle sum f-la

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\ &= \cos(2y) \cos y - \sin(2y) \sin y\end{aligned} \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\text{?}) \cos y \\
 &\quad - \text{?} \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - ? \qquad \qquad \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - \sin y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$



## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\ &= \cos(2y) \cos y - \sin(2y) \sin y \\ &= (\cos^2 y - \sin^2 y) \cos y \\ &\quad - 2 \sin y \cos y \sin y\end{aligned}$$

$y = \arccos x$   
Angle sum f-la  
Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y
 \end{aligned}
 \quad \left| \begin{array}{l} y = \arccos x \\ \text{Angle sum f-la} \\ \text{Express via} \\ \sin y, \cos y \end{array} \right.$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}\cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\ &= \cos(2y) \cos y - \sin(2y) \sin y \\ &= (\cos^2 y - \sin^2 y) \cos y \\ &\quad - 2 \sin y \cos y \sin y \\ &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\ &= \cos^3 y - 3 \sin^2 y \cos y\end{aligned}$$

$y = \arccos x$   
Angle sum f-la  
Express via  
 $\sin y, \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(\text{?}) \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) & y = \arccos x \\
 &= \cos(2y) \cos y - \sin(2y) \sin y & \text{Angle sum f-la} \\
 &= (\cos^2 y - \sin^2 y) \cos y & \text{Express via} \\
 &\quad - 2 \sin y \cos y \sin y & \sin y, \cos y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y & \text{Express } \sin y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y & \text{via } \cos y \\
 &= 4\cos^3 y - 3 \cos y
 \end{aligned}$$



## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) & y = \arccos x \\
 &= \cos(2y) \cos y - \sin(2y) \sin y & \text{Angle sum f-la} \\
 &= (\cos^2 y - \sin^2 y) \cos y & \text{Express via} \\
 &\quad - 2 \sin y \cos y \sin y & \sin y, \cos y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y & \text{Express } \sin y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y & \text{via } \cos y \\
 &= 4\cos^3 y - 3\cos y
 \end{aligned}$$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1 - x^2}$ . To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) \\
 &= \cos(2y) \cos y - \sin(2y) \sin y \\
 &= (\cos^2 y - \sin^2 y) \cos y \\
 &\quad - 2 \sin y \cos y \sin y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y \\
 &= 4\cos^3 y - 3\cos y \\
 &= 4x^3 - 3x
 \end{aligned}$$

$y = \arccos x$   
 Angle sum f-la  
 Express via  
 $\sin y, \cos y$

Express  $\sin y$   
 via  $\cos y$

$x = \cos y$

## Example

Rewrite  $\cos(3 \arccos(x))$  as an algebraic expression of  $x$  and  $\sqrt{1-x^2}$ .

To simplify  $\arccos x$  we try to use  $\cos(\arccos x) = x$ . Therefore our aim is to rewrite the expression only using the  $\cos$  function.

$$\begin{aligned}
 \cos(3 \arccos(x)) &= \cos(3y) = \cos(2y + y) & y = \arccos x \\
 &= \cos(2y) \cos y - \sin(2y) \sin y & \text{Angle sum f-la} \\
 &= (\cos^2 y - \sin^2 y) \cos y & \text{Express via} \\
 &\quad - 2 \sin y \cos y \sin y & \sin y, \cos y \\
 &= \cos^3 y - \sin^2 y \cos y - 2 \sin^2 y \cos y \\
 &= \cos^3 y - 3 \sin^2 y \cos y & \text{Express } \sin y \\
 &= \cos^3 y - 3(1 - \cos^2 y) \cos y & \text{via } \cos y \\
 &= 4\cos^3 y - 3 \cos y \\
 &= 4x^3 - 3x & x = \cos y
 \end{aligned}$$