

Calculus I

Exponents and logarithms review

Todor Milev

2019

Outline

- 1 Exponential Functions
 - Two ways to define exponents
 - Basic properties
 - The Natural Exponential Function

Outline

1 Exponential Functions

- Two ways to define exponents
- Basic properties
- The Natural Exponential Function

2 Logarithmic Functions

- Logarithm basics
- Natural Logarithms

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$7^3 \cdot 7^2 = (?) (\quad)$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$7^3 \cdot 7^2 = (7 \cdot 7 \cdot 7)(\quad)$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$7^3 \cdot 7^2 = (7 \cdot 7 \cdot 7)(?)$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$7^3 \cdot 7^2 = (7 \cdot 7 \cdot 7)(7 \cdot 7)$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\begin{aligned} 7^3 \cdot 7^2 &= (7 \cdot 7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\begin{aligned} 7^3 \cdot 7^2 &= (7 \cdot 7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^5 \\ &= 7^{3+2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\begin{aligned} 7^3 \cdot 7^2 &= (7 \cdot 7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^5 \\ &= 7^{3+2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\begin{aligned} 7^3 \cdot 7^2 &= (7 \cdot 7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^5 \\ &= 7^{3+2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\begin{aligned} 7^3 \cdot 7^2 &= (7 \cdot 7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^5 \\ &= 7^{3+2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\frac{7^3}{7^2} = \frac{?}{?}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\frac{7^3}{7^2} = \frac{7 \cdot 7 \cdot 7}{?}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\frac{7^3}{7^2} = \frac{7 \cdot 7 \cdot 7}{?}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\frac{7^3}{7^2} = \frac{7 \cdot 7 \cdot 7}{7 \cdot 7}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\begin{aligned}\frac{7^3}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\begin{aligned}\frac{7^3}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7 \\ &= 7^1\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

① $a^x a^y = a^{x+y}$

$$\begin{aligned}\frac{7^3}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7 \\ &= 7^1 \\ &= 7^{3-2}.\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} \frac{7^3}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7 \\ &= 7^1 \\ &= 7^{3-2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^{\textcolor{red}{x}}}{a^y} = a^{\textcolor{red}{x}-y}$$

$$\begin{aligned} \frac{7^{\textcolor{red}{3}}}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7 \\ &= 7^1 \\ &= 7^{\textcolor{red}{3}-2}. \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} \frac{7^3}{7^2} &= \frac{\cancel{7} \cdot \cancel{7} \cdot 7}{\cancel{7} \cdot \cancel{7}} \\ &= 7 \\ &= 7^1 \\ &= 7^{3-2} \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\left(7^2\right)^4 = 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^8 \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^8 \\ &= 7^{2 \cdot 4} \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^8 \\ &= 7^{2 \cdot 4} \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^8 \\ &= 7^{2 \cdot 4} \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned} (7^2)^4 &= 7^2 \cdot 7^2 \cdot 7^2 \cdot 7^2 \\ &= (7 \cdot 7)(7 \cdot 7)(7 \cdot 7)(7 \cdot 7) \\ &= 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \\ &= 7^8 \\ &= 7^{2 \cdot 4} \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$(5 \cdot 7)^3 = (5 \cdot 7)(5 \cdot 7)(5 \cdot 7)$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$① \quad a^x a^y = a^{x+y}$$

$$② \quad \frac{a^x}{a^y} = a^{x-y}$$

$$③ \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$① \quad a^x a^y = a^{x+y}$$

$$② \quad \frac{a^x}{a^y} = a^{x-y}$$

$$③ \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= ? \cdot ?\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$① \quad a^x a^y = a^{x+y}$$

$$② \quad \frac{a^x}{a^y} = a^{x-y}$$

$$③ \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= 5^3 \cdot ?\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot \textcolor{red}{7} \cdot \textcolor{red}{7} \cdot \textcolor{red}{7} \\ &= 5^3 \cdot \textcolor{red}{?}\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= 5^3 \cdot 7^3\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\textcircled{4} \quad (ab)^x = a^x b^x$$

$$\begin{aligned} (5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= 5^3 \cdot 7^3 \end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\textcircled{4} \quad (ab)^x = a^x b^x$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= 5^3 \cdot 7^3\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\textcircled{4} \quad (ab)^x = a^x b^x$$

$$\begin{aligned}(5 \cdot 7)^3 &= (5 \cdot 7)(5 \cdot 7)(5 \cdot 7) \\ &= 5 \cdot 7 \cdot 5 \cdot 7 \cdot 5 \cdot 7 \\ &= 5 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 7 \\ &= 5^3 \cdot 7^3\end{aligned}$$

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\textcircled{4} \quad (ab)^x = a^x b^x$$

These rules do continue to hold for all $a > 0$, $b > 0$ and arbitrary x and y .

Properties of exponential expressions.

For integer x, y and bases a, b , we demonstrate the exponent rules by example.

$$\textcircled{1} \quad a^x a^y = a^{x+y}$$

$$\textcircled{2} \quad \frac{a^x}{a^y} = a^{x-y}$$

$$\textcircled{3} \quad (a^x)^y = a^{xy}$$

$$\textcircled{4} \quad (ab)^x = a^x b^x$$

These rules do continue to hold for all $a > 0$, $b > 0$ and arbitrary x and y . **The rules do fail when $a < 0$, $b < 0$ and x, y are not integers.**

Exponents overview

- For integer x , we know how to compute a^x as a function of a .

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.
- Choosing one definition makes some statements easier to prove and others more difficult.

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.
- Choosing one definition makes some statements easier to prove and others more difficult.
- We shall discuss pros and cons of the two. In a nutshell:

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.
- Choosing one definition makes some statements easier to prove and others more difficult.
- We shall discuss pros and cons of the two. In a nutshell:
 - the first elementary definition is easier to motivate;

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.
- Choosing one definition makes some statements easier to prove and others more difficult.
- We shall discuss pros and cons of the two. In a nutshell:
 - the first elementary definition is easier to motivate;
 - the second alternative definition is easier to compute with.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid x^q \leq a\}$.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid \text{for which } x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid \text{for which } x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.
- We can then define

$$a^x = \lim_{\substack{y \rightarrow x \\ y\text{-rational}}} a^y$$

For example, a^π would be defined as the limit of the sequence $a^{3.14}, a^{3.141}, a^{3.1415}, \dots$

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid \text{for which } x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.
- We can then define

$$a^x = \lim_{\substack{y \rightarrow x \\ y\text{-rational}}} a^y$$

For example, a^π would be defined as the limit of the sequence $a^{3.14}, a^{3.141}, a^{3.1415}, \dots$

- Cons: not computationally effective; not how computers compute.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.
- We can then define

$$a^x = \lim_{\substack{y \rightarrow x \\ y\text{-rational}}} a^y$$

For example, a^π would be defined as the limit of the sequence $a^{3.14}, a^{3.141}, a^{3.1415}, \dots$

- Cons: not computationally effective; not how computers compute.
- Pros: for non-integer x and y , it is very easy to prove that $a^{x+y} = a^x a^y$ - this follows from the definition of limit above.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid \text{for which } x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.
- We can then define

$$a^x = \lim_{\substack{y \rightarrow x \\ y\text{-rational}}} a^y$$

For example, a^π would be defined as the limit of the sequence $a^{3.14}, a^{3.141}, a^{3.1415}, \dots$

- Cons: not computationally effective; not how computers compute.
- Pros: for non-integer x and y , it is very easy to prove that $a^{x+y} = a^x a^y$ - this follows from the definition of limit above.
- This is the definition assumed in many elementary courses.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ and is read “ n factorial”.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Infinite sum studied much later.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Infinite sum studied much later.

- For arbitrary $a > 0$ define a^x as $a^x = e^{x \ln a}$.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Infinite sum studied much later.

- For arbitrary $a > 0$ define a^x as $a^x = e^{x \ln a}$.
- Cons: more difficult to prove $e^{x+y} = e^x e^y$ and $e^{\ln(1+x)} = 1+x$, proof done later.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdots (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define


$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Infinite sum studied much later.

- For arbitrary $a > 0$ define a^x as $a^x = e^{x \ln a}$.
- Cons: more difficult to prove $e^{x+y} = e^x e^y$ and $e^{\ln(1+x)} = 1+x$, proof done later.
- Pros: this is how e^x and a^x are actually computed (by modern computers and by humans in the past).

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	
1	
0	
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	?
1	
0	
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	
0	
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	?
0	
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	1/2
-2	1/4

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	?
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	?
-2	

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	$\frac{1}{4}$

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	?

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	$\frac{1}{4}$

Exponential Functions


The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.



x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	$\frac{1}{4}$

Exponential Functions

The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.




x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	$\frac{1}{4}$


(Exponential Function Terminology)

An exponential function is a function of the form $f(x) = a^x$, where a is a positive constant.


Graphs of various exponential functions.




Graphs of various exponential functions.




Graphs of various exponential functions.




Graphs of various exponential functions.




Graphs of various exponential functions.



Graphs of various exponential functions.




Graphs of various exponential functions.



Observations


Graphs of various exponential functions.



Observations

- a^x is always ?


Graphs of various exponential functions.



Observations

- a^x is always positive.


Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = ?$ for all a .


Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = ?$
- $\lim_{x \rightarrow -\infty} a^x =$

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x =$
- $\lim_{x \rightarrow -\infty} a^x =$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x =$

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x =$
- $\lim_{x \rightarrow -\infty} a^x =$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = ?$

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x =$
- $\lim_{x \rightarrow -\infty} a^x =$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = 0$.

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x =$
- $\lim_{x \rightarrow -\infty} a^x =$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = 0$.

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x = ?$
- $\lim_{x \rightarrow -\infty} a^x =$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = 0$.

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x = 0$.
- $\lim_{x \rightarrow -\infty} a^x = \infty$.

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = 0$.

$a < 1$:

- $\lim_{x \rightarrow \infty} a^x = 0$.
- $\lim_{x \rightarrow -\infty} a^x = ?$

Graphs of various exponential functions.



Observations

- a^x is always positive.
- $a^0 = 1$ for all a .


For $a > 1$:

- $\lim_{x \rightarrow \infty} a^x = \infty$.
- $\lim_{x \rightarrow -\infty} a^x = 0$.


$a < 1$:

- $\lim_{x \rightarrow \infty} a^x = 0$.
- $\lim_{x \rightarrow -\infty} a^x = \infty$.

Graphical comparison of $y = 2^x$ with $y = x^2$. Axes have different scales.




Graphical comparison of $y = 2^x$ with $y = x^2$. Axes have different scales.




Example

Draw the graph of the function $y = 2^{-x} - 1 = 0.5^x - 1 = \left(\frac{1}{2}\right)^x - 1$.
Assume the graph of $y = 2^x$ given.



Example


Draw the graph of the function $y = 2^{-x} - 1 = 0.5^x - 1 = \left(\frac{1}{2}\right)^x - 1$.
Assume the graph of $y = 2^x$ given.



- Plot of 2^x assumed given.

Example


Draw the graph of the function $y = 2^{-x} - 1 = 0.5^x - 1 = \left(\frac{1}{2}\right)^x - 1$. Assume the graph of $y = 2^x$ given.



- Plot of 2^x assumed given.
- Plot $f(-x) =$ reflect $f(x)$ across y axis.

Example

Draw the graph of the function $y = 2^{-x} - 1 = 0.5^x - 1 = \left(\frac{1}{2}\right)^x - 1$. Assume the graph of $y = 2^x$ given.



- Plot of 2^x assumed given.
- Plot $f(-x) =$ reflect $f(x)$ across y axis.
- Plot $g(x) - 1 =$ shift graph $g(x)$ 1 unit down.

Example (Solve exponential equation without logarithms)

Solve for t .

$$16^{4t} = 8^{t-2}$$

Example (Solve exponential equation without logarithms)

Solve for t .

$$\begin{array}{lcl} 16^{4t} & = & 8^{t-2} \\ \text{Find a common base: } (?)^{4t} & = & (?)^{t-2} \end{array}$$

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned} 16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \end{aligned}$$

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned} 16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \\ 2^{16t} &= 2^{3t-6} \end{aligned}$$

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned} 16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \\ 2^{16t} &= 2^{3t-6} \\ 16t &= 3t - 6 \end{aligned}$$

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned}16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \\ 2^{16t} &= 2^{3t-6} \\ 16t &= 3t - 6 \\ 13t &= -6\end{aligned}$$

Example (Solve exponential equation without logarithms)

Solve for t .

Find a common base:

$$\begin{aligned}16^{4t} &= 8^{t-2} \\ (2^4)^{4t} &= (2^3)^{t-2} \\ 2^{16t} &= 2^{3t-6} \\ 16t &= 3t - 6 \\ 13t &= -6 \\ t &= -\frac{6}{13}.\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$9^x = 2 \cdot 3^x + 63$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{array}{rcl} 9^x & = & 2 \cdot 3^x + 63 \\ 9^x - 2 \cdot 3^x - 63 & = & 0 \\ ? - 2u - 63 & = & 0 \end{array} \quad \left| \text{Substitute } u = 3^x \right.$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{array}{rclcl} 9^x & = & 2 \cdot 3^x + 63 & & \\ \textcolor{red}{9}^x - 2 \cdot 3^x - 63 & = & 0 & \left| \text{Substitute } u = 3^x \right. & \\ \textcolor{red}{?} - 2u - 63 & = & 0 & & \end{array}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$9^x = 2 \cdot 3^x + 63$$

$$9^x - 2 \cdot 3^x - 63 = 0$$

$$u^2 - 2u - 63 = 0$$

Substitute $u = 3^x$

Example (Solving a quadratic exponential equation)

Solve for x .

$$9^x = 2 \cdot 3^x + 63$$

$$9^x - 2 \cdot 3^x - 63 = 0$$

Substitute $u = 3^x$

$$u^2 - 2u - 63 = 0$$

$$(\textcolor{red}{?})(\textcolor{red}{?}) = 0$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$9^x = 2 \cdot 3^x + 63$$

$$9^x - 2 \cdot 3^x - 63 = 0$$

$$u^2 - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

Substitute $u = 3^x$

Example (Solving a quadratic exponential equation)

Solve for x .

$$9^x = 2 \cdot 3^x + 63$$

$$9^x - 2 \cdot 3^x - 63 = 0$$

Substitute $u = 3^x$

$$u^2 - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u = 9 \text{ or } u = -7$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\u^2 - 2u - 63 &= 0 \\(u - 9)(u + 7) &= 0 \\u &= 9 \text{ or } u = -7 \\3^x &= 9 \text{ or } 3^x = -7\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned} 9^x &= 2 \cdot 3^x + 63 \\ 9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\ u^2 - 2u - 63 &= 0 \\ (u - 9)(u + 7) &= 0 \end{aligned}$$

$$\begin{aligned} u &= 9 \quad \text{or} \quad u = -7 \\ 3^x &= 9 \quad \text{or} \quad 3^x = -7 \\ x &= ? \end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\u^2 - 2u - 63 &= 0 \\(u - 9)(u + 7) &= 0\end{aligned}$$

$$\begin{aligned}u &= 9 \quad \text{or} \quad u = -7 \\3^x &= 9 \quad \text{or} \quad 3^x = -7 \\x &= 2\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\u^2 - 2u - 63 &= 0 \\(u - 9)(u + 7) &= 0\end{aligned}$$

$$\begin{aligned}u &= 9 \quad \text{or} \quad u = -7 \\3^x &= 9 \quad \text{or} \quad 3^x = -7 \\x &= 2 & \quad \text{?}\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned}9^x &= 2 \cdot 3^x + 63 \\9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\u^2 - 2u - 63 &= 0 \\(u - 9)(u + 7) &= 0\end{aligned}$$

$$\begin{aligned}u &= 9 \quad \text{or} \quad u = -7 \\3^x &= 9 \quad \text{or} \quad 3^x = -7 \\x &= 2 & \text{no real solution}\end{aligned}$$

Example (Solving a quadratic exponential equation)

Solve for x .

$$\begin{aligned} 9^x &= 2 \cdot 3^x + 63 \\ 9^x - 2 \cdot 3^x - 63 &= 0 & \left| \text{Substitute } u = 3^x \right. \\ u^2 - 2u - 63 &= 0 \\ (u - 9)(u + 7) &= 0 \end{aligned}$$

$$\begin{aligned} u &= 9 \quad \text{or} \quad u = -7 \\ 3^x &= 9 \quad \text{or} \quad 3^x = -7 \\ x &= 2 & \quad \text{no real solution} \end{aligned}$$

Therefore $x = 2$ is the solution.

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months.

When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t :

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same **number of chickens** as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : **$c(t)$**

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have **the same** number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) =$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same **number of** chickens as **rabbits**?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

=

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

$$48 \cdot 2^t =$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

$$48 \cdot 2^t =$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

$$48 \cdot 2^t = 6 \cdot 4^t$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

Solve for t : $c(t) = r(t)$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^? \cdot 2^t = 2^?$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^?$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^?$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^{2t}$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^{2t}$$

$$2^{t+3} = 2^{2t}$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^{2t}$$

$$2^{t+3} = 2^{2t}$$

$$t + 3 = 2t$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^{2t}$$

$$2^{t+3} = 2^{2t}$$

$$t + 3 = 2t$$

$$t = 3.$$

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let $c(t)$ denote the number of chickens after t years, and let $r(t)$ denote the number of rabbits after t years.

$$\text{Solve for } t: c(t) = r(t)$$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

$$\text{Find a common base: } 2^3 \cdot 2^t = 2^{2t}$$

$$2^{t+3} = 2^{2t}$$



$$t + 3 = 2t$$

$$t = 3.$$

Therefore the chicken and rabbit populations are equal after 3 years.



The Natural Exponential Function

- One base for an exponential function is especially useful.






The Natural Exponential Function

- One base for an exponential function is especially useful.
- It has a special property: its tangent line at $x = 0$ has slope $m = 1$.






The Natural Exponential Function

- One base for an exponential function is especially useful.
- It has a special property: its tangent line at $x = 0$ has slope $m = 1$.
- We call this number e , known as Euler's number or Napier's constant.






The Natural Exponential Function

- One base for an exponential function is especially useful.
- It has a special property: its tangent line at $x = 0$ has slope $m = 1$.
- We call this number e , known as Euler's number or Napier's constant.
- e is a number between 2 and 3.




The Natural Exponential Function

- One base for an exponential function is especially useful.
- It has a special property: its tangent line at $x = 0$ has slope $m = 1$.
- We call this number e , known as Euler's number or Napier's constant.
- e is a number between 2 and 3.
- In fact, $e = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots \approx 2.71828$.




Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.

Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.

Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.

Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.

Logarithmic Functions



- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.
- Therefore f has an inverse function, f^{-1} .

Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.
- Therefore f has an inverse function, f^{-1} .

Definition ($\log_a x$)

The inverse function of $f(x) = a^x$ is called the logarithmic function with base a , and is written $\log_a x$. It is defined by the formula

$$\log_a x = y \quad \Leftrightarrow \quad a^y = x.$$

Logarithmic Functions




- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.
- Therefore f has an inverse function, f^{-1} .
- The graph shows $y = a^x$ for $a > 1$.

Definition ($\log_a x$)

The inverse function of $f(x) = a^x$ is called the logarithmic function with base a , and is written $\log_a x$. It is defined by the formula

$$\log_a x = y \quad \Leftrightarrow \quad a^y = x.$$

Logarithmic Functions



- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.
- Therefore f has an inverse function, f^{-1} .
- The graph shows $y = a^x$ for $a > 1$.
- The graph of $y = \log_a x$ is the reflection of this in the line $y = x$.

Definition ($\log_a x$)

The inverse function of $f(x) = a^x$ is called the logarithmic function with base a , and is written $\log_a x$. It is defined by the formula

$$\log_a x = y \quad \Leftrightarrow \quad a^y = x.$$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:

① $\log_3 81 =$

② $\log_{25} 5 =$

③ $\log_{10} 0.001 =$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:

① $\log_3 81 = ?$

② $\log_{25} 5 = ?$

③ $\log_{10} 0.001 = ?$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:

① $\log_3 81 = 4$ because $3^4 = 81$.

② $\log_{25} 5 = ?$

③ $\log_{10} 0.001 = ?$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:

① $\log_3 81 = 4$ because $3^4 = 81$.

② $\log_{25} 5 = ?$

③ $\log_{10} 0.001 = ?$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:

- 1 $\log_3 81 = 4$ because $3^4 = 81$.
- 2 $\log_{25} 5 = \frac{1}{2}$ because $25^{\frac{1}{2}} = \sqrt{25} = 5$.
- 3 $\log_{10} 0.001 = ?$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .

Example

Evaluate:


- 1 $\log_3 81 = 4$ because $3^4 = 81$.
- 2 $\log_{25} 5 = \frac{1}{2}$ because $25^{\frac{1}{2}} = \sqrt{25} = 5$.
- 3 $\log_{10} 0.001 = ?$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .


Example

Evaluate:


- 1 $\log_3 81 = 4$ because $3^4 = 81$.
- 2 $\log_{25} 5 = \frac{1}{2}$ because $25^{\frac{1}{2}} = \sqrt{25} = 5$.
- 3 $\log_{10} 0.001 = -3$ because $10^{-3} = 0.001$.




- Suppose $a > 1$.




- Suppose $a > 1$.
- Domain of a^x : ?
- Range of a^x : ?
- Domain of $\log_a x$: ?
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : ?
- Range of a^x : ?
- Domain of $\log_a x$: ?
- Range of $\log_a x$: ?



- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : ?
- Domain of $\log_a x$: ?
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : ?
- Domain of $\log_a x$: ?
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$:
?
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$:
?
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$: $(0, \infty)$.
- Range of $\log_a x$: ?




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$: $(0, \infty)$.
- Range of $\log_a x$: ?





- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$: $(0, \infty)$.
- Range of $\log_a x$: \mathbb{R} .




- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$: $(0, \infty)$.
- Range of $\log_a x$: \mathbb{R} .
- $\log_a(a^x) = x$ for $x \in \mathbb{R}$.
- $a^{\log_a x} = x$ for $x > 0$.

Graphs of various logarithmic functions with $a > 1$ 

Graphs of various logarithmic functions with $a > 1$ 

Graphs of various logarithmic functions with $a > 1$ 

Graphs of various logarithmic functions with $a > 1$ 

Theorem (Properties of Logarithmic Functions)

If $a > 1$, the function $f(x) = \log_a x$ is a one-to-one, continuous, increasing function with domain $(0, \infty)$ and range \mathbb{R} . If $x, y, a, b > 0$ and r is any real number, then

$$\textcircled{1} \log_a(xy) = \log_a x + \log_a y.$$

$$\textcircled{2} \log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y.$$

$$\textcircled{3} \log_a(x^r) = r \log_a x.$$

$$\textcircled{4} \log_a(x) = \log_b x \log_a b = \frac{\log_b x}{\log_b a} = \frac{\ln x}{\ln a}.$$

Use the properties of logarithms to evaluate the following.

Example

$$\log_4 2 + \log_4 32$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\log_4 2 + \log_4 32 = \log_4 (2 \cdot 32)$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\log_4 2 + \log_4 32 = \log_4 (2 \cdot 32)$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= ?\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\log_2 80 - \log_2 5 = \log_2 \left(\frac{80}{5} \right)$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16)\end{aligned}$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16) \\ &= ?\end{aligned}$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16) \\ &= 4 \\ &\quad (\text{because } 2^4 = 16.)\end{aligned}$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16) \\ &= 4 \\ &\quad (\text{because } 2^4 = 16.)\end{aligned}$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example

$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16) \\ &= 4 \\ &\quad (\text{because } 2^4 = 16.)\end{aligned}$$

Use the properties of logarithms to evaluate the following.

Example

$$\begin{aligned}\log_4 2 + \log_4 32 &= \log_4(2 \cdot 32) \\ &= \log_4(64) \\ &= 3 \\ &\quad (\text{because } 4^3 = 64.)\end{aligned}$$

Example


$$\begin{aligned}\log_2 80 - \log_2 5 &= \log_2 \left(\frac{80}{5} \right) \\ &= \log_2(16) \\ &= 4 \\ &\quad (\text{because } 2^4 = 16.)\end{aligned}$$

Natural Logarithms

Definition ($\ln x$)

The logarithm with base e is called the natural logarithm, and has a special notation:

$$\log_e x = \ln x.$$




Natural Logarithms

Definition ($\ln x$)

The logarithm with base e is called the natural logarithm, and has a special notation:

$$\log_e x = \ln x.$$




$$\bullet \ln x = y \quad \Leftrightarrow \quad e^y = x.$$

Natural Logarithms

Definition ($\ln x$)

The logarithm with base e is called the natural logarithm, and has a special notation:

$$\log_e x = \ln x.$$




- $\ln x = y \quad \Leftrightarrow \quad e^y = x.$
- $\ln(e^x) = x$ for $x \in \mathbb{R}.$

Natural Logarithms

Definition ($\ln x$)

The logarithm with base e is called the natural logarithm, and has a special notation:

$$\log_e x = \ln x.$$



- $\ln x = y \quad \Leftrightarrow \quad e^y = x.$
- $\ln(e^x) = x$ for $x \in \mathbb{R}.$
- $e^{\ln x} = x$ for $x > 0.$

Summary of logarithm notation conventions

	Name	ISO notation	Other notation	Used in
$\log_2(x)$	binary logarithm	$\text{lb}(x)$	$\text{ld}(x)$, $\log(x)$, $\text{lg}(x)$	computer science, information theory, music theory, photography
$\log_e(x)$	natural logarithm	$\ln(x)$	$\log(x)$	mathematics, physics, chemistry, statistics, economics, information theory, and engineering
$\log_{10}(x)$	common logarithm	$\text{lg}(x)$	$\log(x)$	various engineering, logarithm tables, handheld calculators, spectroscopy

Table source: Wikipedia

- Standardized in ISO_31-11 (International Standards Organization).

Example

Solve the equation.

$$e^{5-3x} = 10$$

Example

Solve the equation.

$$\begin{array}{rclcl} e^{5-3x} & = & 10 & & \text{apply } \ln \\ \ln(e^{5-3x}) & = & \ln 10 & & \end{array}$$

Example

Solve the equation.

$$\begin{aligned} e^{5-3x} &= 10 && \text{apply } \ln \\ \ln(e^{5-3x}) &= \ln 10 \\ 5 - 3x &= \ln 10 \end{aligned}$$

Example

Solve the equation.

$$\begin{aligned} e^{5-3x} &= 10 && \text{apply } \ln \\ \ln(e^{5-3x}) &= \ln 10 \\ \textcolor{red}{5} - 3x &= \ln 10 \\ 3x &= \textcolor{red}{5} - \ln 10 \end{aligned}$$

Example

Solve the equation.

$$\begin{aligned} e^{5-3x} &= 10 && \text{apply ln} \\ \ln(e^{5-3x}) &= \ln 10 \\ 5 - 3x &= \ln 10 \\ \textcolor{red}{3}x &= 5 - \ln 10 \\ x &= \frac{5 - \ln 10}{\textcolor{red}{3}} \end{aligned}$$

Example

Solve the equation.

$$\begin{aligned}e^{5-3x} &= 10 && \text{apply } \ln \\ \ln(e^{5-3x}) &= \ln 10 \\ 5 - 3x &= \ln 10 \\ 3x &= 5 - \ln 10 \\ x &= \frac{5 - \ln 10}{3} \\ \text{Calculator: } x &\approx 0.8991.\end{aligned}$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$.

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = ?$.

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(\textcolor{red}{?} \quad \textcolor{red}{?}) (\textcolor{red}{?} \quad \textcolor{red}{?}) = 0$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

$$e^x = 4$$

or

$$e^x = -1$$

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

$$e^x = 4$$

or

$$e^x = -1$$

$$x = \ln 4$$

or

no real solution

Example

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4)(u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

$$e^x = 4$$

or

$$e^x = -1$$

$$x = \ln 4$$

or

no real solution

$$x \approx 1.3863$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = ?$.

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$.

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$.

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = ?$,

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$,

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = ?$.

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(\text{?}) (\text{?}) = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = -\frac{3}{2} \quad \text{or} \quad u =$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u =$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right)$$

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right)$$

or no real solution

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right) = \frac{\ln(?)}{\ln ?}$$

or no real solution

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$

$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right) = \frac{\ln \left(\frac{3}{2} \right)}{\ln 2}$$

or no real solution

Example

Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \quad \text{or} \quad 2u + 1 = 0$$


$$u = \frac{3}{2} \quad \text{or} \quad u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \quad \text{or} \quad 2^x = -\frac{1}{2}$$

$$x = \log_2 \left(\frac{3}{2} \right) = \frac{\ln \left(\frac{3}{2} \right)}{\ln 2} \approx 0.58496 \quad \text{or} \quad \text{no real solution}$$


Example

Draw the graph of $y = \ln(x - 2) - 1$.



Example


Draw the graph of $y = \ln(x - 2) - 1$.



- Graph $y = \ln(x)$ assumed known.

Example


Draw the graph of $y = \ln(x - 2) - 1$.



- Graph $y = \ln(x)$ assumed known.
- $f(x - 2)$ shifts graph 2 units to the right.

Example

Draw the graph of $y = \ln(x - 2) - 1$.




- Graph $y = \ln(x)$ assumed known.
- $f(x - 2)$ shifts graph 2 units to the right.
- $g(x) - 1$ shifts graph 1 unit down.

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$




Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$



Example


Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - ?)}{(u + ?)} = y$$

Set $u = e^x$



Example

Find $f^{-1}(x)$ for


$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - ?)}{(u + ?)} = y$$

Set $u = e^x$


$$e^{-x} = ?$$



Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$
$$\frac{(u - \frac{1}{u})}{(u + \frac{1}{u})} = y$$

Set $u = e^x$

$$e^{-x} = \frac{1}{e^x} = \frac{1}{u}$$

Example


Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) \textcolor{red}{u}}{(u + \frac{1}{u}) \textcolor{red}{u}} = y$$


$$\begin{array}{|l} \text{Set } u = e^x \\ e^{-x} = \frac{1}{e^x} = \frac{1}{u} \end{array}$$



Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$




$$\begin{aligned}\frac{e^x - e^{-x}}{e^x + e^{-x}} &= y \\ \frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} &= y \\ \frac{u^2 - 1}{u^2 + 1} &= y\end{aligned}$$

$$\begin{aligned}\text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u}\end{aligned}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$


$$u^2 - 1 = y(u^2 + 1)$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$


$$u^2(1 - y) = 1 + y$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$


$$u^2(1 - y) = 1 + y$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$


$$u^2 = \frac{1 + y}{1 - y}$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$


Set $u = e^x$

$$e^{-x} = \frac{1}{e^x} = \frac{1}{u}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$


Set $u = e^x$

$$e^{-x} = \frac{1}{e^x} = \frac{1}{u}$$

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$


$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Take \ln

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$


$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Take \ln

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



answer

$$f^{-1}(y) = \frac{1}{2} \ln \left(\frac{1+y}{1-y} \right)$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1+y}{1-y}$$

$$(e^x)^2 = \frac{1+y}{1-y}$$

$$e^{2x} = \frac{1+y}{1-y}$$

$$x = \frac{1}{2} \ln \left(\frac{1+y}{1-y} \right)$$


$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Take \ln

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$



Final answer, **relabelled**:

$$f^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln \left(\frac{1 + y}{1 - y} \right)$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$


Take \ln

Example

Find $f^{-1}(x)$ for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

$f = \tanh =$ **hyperbolic tangent function.**



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$$

$$\frac{e^x - e^{-x}}{e^x + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^2 - 1}{u^2 + 1} = y$$

$$u^2 - 1 = y(u^2 + 1)$$

$$u^2(1 - y) = 1 + y$$

$$u^2 = \frac{1 + y}{1 - y}$$

$$(e^x)^2 = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln \left(\frac{1 + y}{1 - y} \right)$$

$$\begin{aligned} \text{Set } u &= e^x \\ e^{-x} &= \frac{1}{e^x} = \frac{1}{u} \end{aligned}$$

Take \ln