

Calculus II

Comparison and limit-comparison tests, part 3

Todor Milev

2019

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7} + n^5}$ for convergence or divergence.

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \text{---}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7} + n^5}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7} + n^5}, \quad b_n = \frac{2n^2}{n^5}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{\sqrt{7 + n^5}}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}}
 \end{aligned}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^2}$ is a constant multiple of a p -series with $p =$

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^{\frac{1}{2}}}$ is a constant multiple of a p -series with $p = \frac{1}{2}$.

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^2}$ is a constant multiple of a p -series with $p = \frac{1}{2}$.
- Therefore $\sum \frac{2}{n^2}$ is

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^{1/2}}$ is a constant multiple of a p -series with $p = \frac{1}{2}$.
- Therefore $\sum \frac{2}{n^{1/2}}$ is divergent

Example

Test the series $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ for convergence or divergence.

- The dominant part of the numerator is $2n^2$ and the dominant part of the denominator is $\sqrt{n^5} = n^{5/2}$.

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^{1/2}}$ is a constant multiple of a p -series with $p = \frac{1}{2}$.
- Therefore $\sum \frac{2}{n^{1/2}}$ is divergent, and so is $\sum \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$.