

Calculus II

Power series expansion of arctangent, part 1

Todor Milev

2019

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$\arctan(x)$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\arctan(x) = \int d(\arctan x) \quad \Bigg| \quad \text{up to const.}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\arctan(x) = \int d(\arctan x) = \int (\arctan x)' dx \quad \Bigg| \text{ up to const.}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned} \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\ &= \int \left(? \right) dx \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}\arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\ &= \int \left(\frac{1}{1+x^2} \right) dx\end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}\arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\ &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx\end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1 - (-x^2)} \right) dx \\
 &= \int \left(? \right) dx && \left| \text{for } |x| < 1 \right.
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1 - (-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right.
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right)
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} ?
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int \left(1 - x^2 + x^4 - x^6 + \dots \right) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} ?
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}
 \end{aligned}$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = ?$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.
- Therefore the theorem on integrating power series implies that

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}, \text{ for } |x| < 1.$$

Example

Find a power series for $\arctan x$ and state its radius of convergence.

$$\begin{aligned}
 \arctan(x) &= \int d(\arctan x) = \int (\arctan x)' dx && \left| \text{up to const.} \right. \\
 &= \int \left(\frac{1}{1+x^2} \right) dx = \int \left(\frac{1}{1-(-x^2)} \right) dx \\
 &= \int (1 - x^2 + x^4 - x^6 + \dots) dx && \left| \text{for } |x| < 1 \right. \\
 &= \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots \right) + C \\
 &= C + \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}
 \end{aligned}$$

- To find C , plug in $x = 0$: $C = 0$.
- Therefore the theorem on integrating power series implies that

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}, \text{ for } |x| < 1.$$

- By the same theorem, the radius of convergence remains $R = 1$.