Precalculus Factor quadratic with irrational real roots

Todor Miley

2019

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$

$$x_1 x_2 = \frac{c}{a}$$
$$x_1 + x_2 = -\frac{b}{a}$$

$$x^2 + 3x + 1$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$

$$x_1 x_2 = \frac{c}{a}$$

$$x_1 + x_2 = -\frac{b}{a}$$

Factor the quadratic.

$$x^2 + 3x + 1 = (x + ?)$$

• The product of the two roots: $x_1x_2 = 1$.

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1$, $x_2 = 1$ and $x_1 = -1$, $x_2 = -1$.

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = (x + ?)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1$, $x_2 = 1$ and $x_1 = -1$, $x_2 = -1$.

•
$$(x-1)(x-1)$$

 $(x+1)(x+1)$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = (x + ?)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.

•
$$(x-1)(x-1)$$

 $(x+1)(x+1)$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$
 $\begin{vmatrix} x_1x_2 &= \frac{c}{a} \\ x_1 + x_2 &= -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- → No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$

$$\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- → No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$

$$\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$$

$$x^2 + \frac{3}{3}x + 1 = \left(x + ? \right) \left(x + ?\right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- → No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$

$$\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- → No easy factorization; must use quadratic formula.

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = \left(x + ? \right) \left(x + ? \right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1$, $x_2 = 1$ and $x_1 = -1$, $x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- No easy factorization; must use quadratic formula.

$$\Rightarrow \text{ No easy factorization; must use quadratic formula.}$$

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{-3 \pm \sqrt{5}}{2}$$

$$ax^2 + bx + c = a(x - x_1)(x - x_2),$$
 $\begin{vmatrix} x_1x_2 &= \frac{c}{a} \\ x_1 + x_2 &= -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = (x + ?)$$
 $(x + ?)$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- No easy factorization; must use quadratic formula.

$$x_{1}, x_{2} = \frac{-b \pm \sqrt{b^{2} - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^{2} - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$
$$= \frac{-3 \pm \sqrt{5}}{2}$$

$$ax^{2} + bx + c = a(x - x_{1})(x - x_{2}),$$
 $\begin{vmatrix} x_{1}x_{2} &=& \frac{c}{a} \\ x_{1} + x_{2} &=& -\frac{b}{a} \end{vmatrix}$

$$x^2 + 3x + 1 = \left(x - \left(\frac{-3 + \sqrt{5}}{2}\right)\right) \left(x - \left(\frac{-3 - \sqrt{5}}{2}\right)\right)$$

- The product of the two roots: $x_1x_2 = 1$.
- Integer options: $x_1 = 1, x_2 = 1$ and $x_1 = -1, x_2 = -1$.
- $(x-1)(x-1) = (x-1)^2 = x^2 2x + 1$ $(x+1)(x+1) = (x+1)^2 = x^2 + 2x + 1$ both don't work.
- No easy factorization; must use quadratic formula.

$$x_{1}, x_{2} = \frac{-b \pm \sqrt{b^{2} - 4ac}}{2a} = \frac{-3 \pm \sqrt{3^{2} - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$
$$= \frac{-3 \pm \sqrt{5}}{2}$$