

Precalculus

Polynomial inequalities

Todor Milev

2019

Outline

1 Polynomial inequalities

Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (?)(?) &\geq 0 \end{aligned}$$

Example

Solve the inequality.


$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Example

Solve the inequality.

$$\begin{aligned}2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0\end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.




Example

Solve the inequality.

$$\begin{aligned}2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0\end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.




Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$			


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(?) (?)$		


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(?)$		

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(?)$		

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$		

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$?	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$	$+$	


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(?) (?)$		
$(1, \infty)$	$(+)(+)$	+	


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(?)$		
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(?)$		
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$		
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$?	
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	
$(1, \infty)$	$(+)(+)$	$+$	


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(\textcolor{red}{?})(\textcolor{red}{?})$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	


Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(?)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(?)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$?	
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$	$+$	
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	
$(1, \infty)$	$(+)(+)$	$+$	

Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$

Example


Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

$x \in ?$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$

Example

Solve the inequality.

$$\begin{aligned}
 2x^2 + 3x - 5 &\geq 0 \\
 (2x + 5)(x - 1) &\geq 0 \\
 x &\in (-\infty, -\frac{5}{2}] \cup [1, \infty)
 \end{aligned}$$


Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
 The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.



Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$

Example

Solve the inequality.




$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

$$x \in (-\infty, -\frac{5}{2}] \cup [1, \infty)$$

Left hand side vanishes when $x = -\frac{5}{2}$ and when $x = 1$.
The two roots split the real line into three intervals:
 $(-\infty, -\frac{5}{2})$, $(-\frac{5}{2}, 1)$, $(1, \infty)$.




Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	$+$	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	$+$	100	$f(100) > 0$

Example

Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


Example



Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - \quad) (x - \quad) (x - \quad) > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - ?) (x - ?)(x - ?) > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - (-\frac{1}{2})) (x - 1)(x - 2) > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - (-\frac{1}{2})) (x - 1)(x - 2) > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$.



Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$.



Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$.



Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		

Example



Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		

Example



Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		
$(1, 2)$		

Example



Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		
$(1, 2)$		
$(2, \infty)$		

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$?	?
$(-\frac{1}{2}, 1)$?	?
$(1, 2)$?	?
$(2, \infty)$?	?

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	$-$
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	$+$
$(1, 2)$	$(+)(+)(-)$	$-$
$(2, \infty)$	$(+)(+)(+)$	$+$

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$


$$x \in ?$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$


$$x \in (-\frac{1}{2}, 1) \cup (2, \infty)$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+

Example




Plot the function $2x^3 - 5x^2 + x + 2$. Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

$$x \in (-\frac{1}{2}, 1) \cup (2, \infty)$$

Left hand side vanishes when $x = -\frac{1}{2}$, when $x = 1$ and when $x = 2$. The two roots split the real line into four intervals: $(-\infty, -\frac{1}{2})$, $(-\frac{1}{2}, 1)$, $(1, 2)$, $(2, \infty)$.



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+