

Calculus II

Polar coordinates

Todor Milev

2019

Outline

1 Polar Coordinates

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Polar Coordinates

- The polar coordinate system is an alternative to the Cartesian coordinate system.

Polar Coordinates

- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).




A diagram showing the origin of a coordinate system. It consists of a small blue dot with the letter O placed to its left.

O

Polar Coordinates

- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.




Polar Coordinates

- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.


• P

- Let P be a point in the plane.



Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- Let P be a point in the plane.
- Let θ denote the angle between the polar axis and the line OP .

Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- Let P be a point in the plane.
- Let θ denote the angle between the polar axis and the line OP .
- Let r denote the length of the segment OP .

Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- Let P be a point in the plane.
- Let θ denote the angle between the polar axis and the line OP .
- Let r denote the length of the segment OP .
- Then P is represented by the ordered pair (r, θ) .

Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- The letters (x, y) imply Cartesian coordinates and the letters (r, θ) - polar.

Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- The letters (x, y) imply Cartesian coordinates and the letters (r, θ) - polar. When we use other letters, it should be clear from context whether we mean Cartesian or polar coordinates.

Polar Coordinates


- The polar coordinate system is an alternative to the Cartesian coordinate system.
- Choose a point in the plane called O (the origin).
- Draw a ray starting at O . The ray is called the polar axis. This ray is usually drawn horizontally to the right.



- The letters (x, y) imply Cartesian coordinates and the letters (r, θ) - polar. When we use other letters, it should be clear from context whether we mean Cartesian or polar coordinates. If not, one must request clarification.



- 1 What if θ is negative?
- 2 What if r is negative?
- 3 What if r is 0?

- 1 What if θ is negative?
- 2 What if r is negative?
- 3 What if r is 0?





- 1 Positive angles θ are measured in the counterclockwise direction from O . Negative angles are measured in the clockwise direction.

- 1 What if θ is negative?
- 2 What if r is negative?
- 3 What if r is 0?




- 1 Positive angles θ are measured in the counterclockwise direction from O . Negative angles are measured in the clockwise direction.
- 2 Points with polar coordinates $(-r, \theta)$ and (r, θ) lie on the same line through O and at the same distance from O , but on opposite sides.



- 1 What if θ is negative?
- 2 What if r is negative?
- 3 What if r is 0?






- 1 Positive angles θ are measured in the counterclockwise direction from O . Negative angles are measured in the clockwise direction.
- 2 Points with polar coordinates $(-r, \theta)$ and (r, θ) lie on the same line through O and at the same distance from O , but on opposite sides.
- 3 If $r = 0$, then $(0, \theta)$ represents O for all values of θ .







- There are many ways to represent the same point.



- There are many ways to represent the same point.
- We could use a negative θ .



- There are many ways to represent the same point.
- We could use a negative θ .
- We could go around more than once.



- There are many ways to represent the same point.
- We could use a negative θ .
- We could go around more than once.
- We could use a negative r .

- Let P_1 be point with polar coordinates (r_1, θ_1) .
- Let P_2 be point with polar coordinates (r_2, θ_2) .

- Let P_1 be point with polar coordinates (r_1, θ_1) .
- Let P_2 be point with polar coordinates (r_2, θ_2) .

Observation

P_1 coincides with P_2 if one of the three mutually exclusive possibilities holds:



- *$r_1 = r_2 \neq 0$ and $\theta_2 = \theta_1 + 2k\pi, k \in \mathbb{Z}$,*
- *$r_1 = -r_2 \neq 0$ and $\theta_2 = \theta_1 + (2k + 1)\pi, k \in \mathbb{Z}$,*
- *$r_1 = r_2 = 0$ and θ is arbitrary.*

- Let P_1 be point with polar coordinates (r_1, θ_1) .
- Let P_2 be point with polar coordinates (r_2, θ_2) .

Observation

P_1 coincides with P_2 if one of the three mutually exclusive possibilities holds:

- $r_1 = r_2 \neq 0$ and $\theta_2 = \theta_1 + 2k\pi, k \in \mathbb{Z}$,
- $r_1 = -r_2 \neq 0$ and $\theta_2 = \theta_1 + (2k + 1)\pi, k \in \mathbb{Z}$,
- $r_1 = r_2 = 0$ and θ is arbitrary.




- Let P_1 be point with polar coordinates (r_1, θ_1) .
- Let P_2 be point with polar coordinates (r_2, θ_2) .


Observation

P_1 coincides with P_2 if one of the three mutually exclusive possibilities holds:

- $r_1 = r_2 \neq 0$ and $\theta_2 = \theta_1 + 2k\pi, k \in \mathbb{Z}$,
- $r_1 = -r_2 \neq 0$ and $\theta_2 = \theta_1 + (2k + 1)\pi, k \in \mathbb{Z}$,
- $r_1 = r_2 = 0$ and θ is arbitrary.



- How do we go from polar coordinates to Cartesian coordinates?




$$x =$$

$$y =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .




$$x =$$

$$y =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .



$$x =$$

$$y =$$


$$\cos \theta =$$

$$\sin \theta =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .



$$x =$$

$$y =$$


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .



$$x =$$

$$y =$$


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .



$$x =$$

$$y =$$


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .



$$x = r \cos \theta$$

$$y = r \sin \theta$$


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$


$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\cos \theta = \frac{x}{r}$$


$$\sin \theta = \frac{y}{r}$$

$$r^2 =$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\cos \theta = \frac{x}{r}$$


$$\sin \theta = \frac{y}{r}$$

$$r^2 = x^2 + y^2$$

$$r =$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\cos \theta = \frac{x}{r}$$


$$\sin \theta = \frac{y}{r}$$

$$r^2 = x^2 + y^2$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\cos \theta = \frac{x}{r}$$


$$\sin \theta = \frac{y}{r}$$

$$r^2 = x^2 + y^2$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta =$$

- How do we go from polar coordinates to Cartesian coordinates?
- Suppose a point has polar coordinates (r, θ) and Cartesian coordinates (x, y) .
- How do we go from Cartesian coordinates to polar coordinates?



$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

$$r^2 = x^2 + y^2$$

$$r = \sqrt{x^2 + y^2}$$

$$\theta = \arcsin\left(\frac{y}{r}\right) \quad \text{if } x > 0$$

$$= \arccos\left(\frac{x}{r}\right) \quad \text{if } y > 0$$

$$= \arctan\left(\frac{y}{x}\right) \quad \text{if } x > 0$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta =$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = \quad \cos$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3}$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\right)$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right)$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta =$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3}$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3} = 2 \left(\quad \right)$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3} = 2 \left(\frac{\sqrt{3}}{2} \right)$$

Example

Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3} = 2 \left(\frac{\sqrt{3}}{2} \right) = \sqrt{3}$$

Example


Convert the point $(2, \frac{\pi}{3})$ from polar to Cartesian coordinates.

$$x = r \cos \theta = 2 \cos \frac{\pi}{3} = 2 \left(\frac{1}{2} \right) = 1$$

$$y = r \sin \theta = 2 \sin \frac{\pi}{3} = 2 \left(\frac{\sqrt{3}}{2} \right) = \sqrt{3}$$


Therefore the point with polar coordinates $(2, \frac{\pi}{3})$ has Cartesian coordinates $(1, \sqrt{3})$.

Example



Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

$$r = \pm \sqrt{x^2 + y^2}$$

$$\tan \theta = \frac{y}{x}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.

$$r = \pm \sqrt{x^2 + y^2}$$

$$\tan \theta = \frac{y}{x}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.

$$r = \sqrt{x^2 + y^2}$$

$$\tan \theta = \frac{y}{x}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\tan \theta = \frac{y}{x}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.

$$\begin{aligned}r &= \sqrt{x^2 + y^2} \\&= \sqrt{1^2 + (-1)^2} \\&= \sqrt{2}\end{aligned}$$

$$\begin{aligned}\tan \theta &= \frac{y}{x} \\&= -1\end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\begin{aligned} \tan \theta &= \frac{y}{x} \\ &= -1 \end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.

$$\begin{aligned}
 r &= \sqrt{x^2 + y^2} \\
 &= \sqrt{1^2 + (-1)^2} \\
 &= \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 \tan \theta &= \frac{y}{x} \\
 &= -1
 \end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.

$$\begin{aligned}
 r &= \sqrt{x^2 + y^2} \\
 &= \sqrt{1^2 + (-1)^2} \\
 &= \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 \tan \theta &= \frac{y}{x} \\
 &= -1
 \end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.
- Of the two values above, only $\theta =$ gives a point in the fourth quadrant.

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\begin{aligned} \tan \theta &= \frac{y}{x} \\ &= -1 \end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.
- Of the two values above, only $\theta = \frac{7\pi}{4}$ gives a point in the fourth quadrant.

$$\begin{aligned}
 r &= \sqrt{x^2 + y^2} \\
 &= \sqrt{1^2 + (-1)^2} \\
 &= \sqrt{2}
 \end{aligned}$$

$$\begin{aligned}
 \tan \theta &= \frac{y}{x} \\
 &= -1
 \end{aligned}$$

Example




Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.
- Of the two values above, only $\theta = \frac{7\pi}{4}$ gives a point in the fourth quadrant.
- \Rightarrow one representation of $(1, -1)$ in polar coordinates is $(\sqrt{2}, \frac{7\pi}{4})$.

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\begin{aligned} \tan \theta &= \frac{y}{x} \\ &= -1 \end{aligned}$$

Example



Represent the point with Cartesian coordinates $(1, -1)$ in terms of polar coordinates.

- Suppose r is positive.
- $\tan \theta = -1$ for $\theta = \frac{3\pi}{4}, \frac{7\pi}{4}$, and many other angles.
- $(1, -1)$ is in the fourth quadrant.
- Of the two values above, only $\theta = \frac{7\pi}{4}$ gives a point in the fourth quadrant.
- \Rightarrow one representation of $(1, -1)$ in polar coordinates is $(\sqrt{2}, \frac{7\pi}{4})$.
- $(\sqrt{2}, -\frac{\pi}{4})$ is another.

$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ &= \sqrt{1^2 + (-1)^2} \\ &= \sqrt{2} \end{aligned}$$

$$\begin{aligned} \tan \theta &= \frac{y}{x} \\ &= -1 \end{aligned}$$