

**Arithmetics**  
**Division and fractions**  
**[calculator-algebra.org](http://calculator-algebra.org)**

Todor Milev

2019

## Definition (Division, exact)

To **divide** a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

answer = ?

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

$$\text{answer} = \frac{5}{3}$$



## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

$$\text{answer} = \frac{5}{3}$$

## Observation

*The quotient of two integers equals the fraction formed by putting the dividend as the numerator and the divisor as the denominator.*

$$q = \frac{p}{d}$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

$$\text{answer} = \frac{5}{3}$$

## Observation

*The quotient of two integers equals **the fraction** formed by putting the dividend as the numerator and the divisor as the denominator.*

$$q = \frac{p}{d}$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

$$\text{answer} = \frac{5}{3}$$

## Observation

*The quotient of two integers equals the fraction formed by putting the **dividend as the numerator** and the divisor as the denominator.*

$$q = \frac{p}{d}$$

## Definition (Division, exact)

To divide a number  $p$  (**dividend**) by a number  $d$  (**divisor**) means to find a number  $q$  (**quotient**) so that

$$q \cdot d = p$$

## Example

Divide 5 by 3.

$$\text{answer} = \frac{5}{3}$$

## Observation

*The quotient of two integers equals the fraction formed by putting the dividend as the numerator and the **divisor as the denominator**.*

$$q = \frac{p}{d}$$

- Recall exact division: 
$$\begin{aligned} p' &= q' \cdot d' \\ q' &= \frac{p'}{d'} \end{aligned}$$
- Quotient may fail to reduce to an integer.
- What if we want an integer quotient?

## Definition (Integer division with remainder)

To **divide** an integer  $p > 0$  by an integer  $d > 0$  **with remainder**  $r \geq 0$  means to find the largest integer  $q \geq 0$  and the smallest  $0 \leq r$  so that:

$$p = q \cdot d + r$$

$p$  is called the **dividend**,  $d$  is called the **divisor**,  $q$  is called the **quotient** and  $r$  is called the **remainder**.

## Example

Divide 7 by 3 with remainder.  $7 = 2 \cdot 3 + 1$ .

- Differences between exact division integer division.
  - Integer division quotient is integer, exact division quotient is fraction.
  - Exact division: no notion of remainder.

- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:

## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:

## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = ?$



Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:

## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*


- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6            ? times.

Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6                  ? times.

Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6                  ? times.

Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6

3 times.

Divide 7 by 3.





- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 **exactly** 3 times.

Divide 7 by 3.



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = ?$



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.

## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$ .



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.


## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.


## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$



- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.


## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$



3 divides 7 ? times

- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.


## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$



3 divides 7 2 times

- Recall integer division of  $p$  by  $d$  with remainder:  $p = q \cdot d + r$ .
- Integer division of  $p$  by  $d$  answers the following question:


## Question

*How many times does the length  $d$  fit inside the length  $p$ ?*

- $p$  may be divisible by  $d$  an exact number of times.
- $p$  may fail to be integer-divisible by  $d$ , then we have remainder.


## Example

Divide 6 by 2 with remainder. Solution:  $6 = 3 \cdot 2 + 0$ .



2 divides 6 exactly 3 times.

Divide 7 by 3. Solution:  $7 = 2 \cdot 3 + 1$



3 divides 7 **non-exactly** 2 times **with remainder 1**.



### Example (Divisor 1-digit, quotient 1-digit)

Divide 7 by 2 with remainder. To solve, we need to answer: what is the largest integer which, when multiplied by 2, stays smaller than 7? Try:

$$0 \cdot 2 = 0$$

$$1 \cdot 2 = 2$$

$$2 \cdot 2 = 4$$

$$3 \cdot 2 = 6$$

$$4 \cdot 2 = 8 > 7$$

$\Rightarrow 7 = 3 \cdot 2 + x$ . Solve:  $x = 7 - 3 \cdot 2 = 1$ . Therefore  $7 = 3 \cdot 2 + 1$ .

### Observation (Question to answer when dividing with remainder)

*What is the largest integer which, when multiplied by  $d$ , remains smaller than  $p$ ?*

- To answer this question, we guess quickly as shown above.
- Later on we learn to divide large numbers without guessing.
- However, we still need the guessing approach as a building block of the complete division algorithm.

## Example (Integer division: 1-digit dividend, 1-digit divisor)

Divide with remainder:

$$5 \text{ by } 2 : \quad 5 = 2 \cdot 2 + 1$$

$$3 \text{ by } 5 : \quad 5 = 0 \cdot 5 + 3$$

$$8 \text{ by } 4 : \quad 5 = 2 \cdot 4$$

$$9 \text{ by } 8 : \quad 5 = 1 \cdot 8 + 1$$

$$3 \text{ by } 1 : \quad 3 = 3 \cdot 1$$

$$9 \text{ by } 2 : \quad 6 = 4 \cdot 2 + 1$$

All quotients are known to be one-digit numbers.

## Example (Integer division: 1-digit divisor, 1-digit quotient)

Divide with remainder:

$$12 \text{ by } 6 : \quad 12 = 2 \cdot 6$$

$$14 \text{ by } 5 : \quad 14 = 2 \cdot 5 + 4$$

$$37 \text{ by } 5 : \quad 37 = 7 \cdot 5 + 2$$

$$40 \text{ by } 9 : \quad 40 = 4 \cdot 9 + 4$$

$$49 \text{ by } 7 : \quad 49 = 7 \cdot 7$$

$$57 \text{ by } 8 : \quad 57 = 7 \cdot 8 + 1$$

$$67 \text{ by } 7 : \quad 67 = 9 \cdot 7 + 4$$


$$82 \text{ by } 9 : \quad 82 = 9 \cdot 9 + 1$$

All quotients are known to be one-digit numbers.

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .


In computer science this function is called the *floor* function, also the *round-down* function.



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = ?$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor 1.5 \rfloor =$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor 1.5 \rfloor =$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = ?$$

$$\lfloor 1.5 \rfloor =$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor =$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$

$$\left\lfloor \frac{3}{2} \right\rfloor =$$


$$\lfloor \pi \rfloor =$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = ?$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor =$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = ?$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor =$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = ?$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\lfloor \pi \rfloor =$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$


$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor = ?$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function, also the *round-down* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor 1.5 \rfloor = \lfloor 1 + 0.5 \rfloor = 1$$

$$\left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\left\lfloor \frac{3}{2} \right\rfloor = \left\lfloor \frac{2+1}{2} \right\rfloor = \left\lfloor \frac{2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 1 + \frac{1}{2} \right\rfloor = 1$$

$$\lfloor \pi \rfloor = \lfloor 3.1415\dots \rfloor = 3$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = ?$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$



## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor =$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = ?$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor =$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor = ?$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor = 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor = 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor =$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}\left\lfloor \frac{1}{3} \right\rfloor &= 0 \\ \left\lfloor \frac{6}{7} \right\rfloor &= 0 \\ \left\lfloor 2 + \frac{3}{11} \right\rfloor &= 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2 \\ \left\lfloor 10 + \frac{1}{7} \right\rfloor &= \end{aligned}$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$



## Example

Compute the floor (round-down) function.

$$\begin{aligned}\left\lfloor \frac{1}{3} \right\rfloor &= 0 \\ \left\lfloor \frac{6}{7} \right\rfloor &= 0 \\ \left\lfloor 2 + \frac{3}{11} \right\rfloor &= 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2 \\ \left\lfloor 10 + \frac{1}{7} \right\rfloor &= \end{aligned}$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}\left\lfloor \frac{1}{3} \right\rfloor &= 0 \\ \left\lfloor \frac{6}{7} \right\rfloor &= 0 \\ \left\lfloor 2 + \frac{3}{11} \right\rfloor &= 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2 \\ \left\lfloor 10 + \frac{1}{7} \right\rfloor &= \end{aligned}$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor = 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor = ?$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$\left\lfloor \frac{6}{7} \right\rfloor = 0$$

$$\left\lfloor 2 + \frac{3}{11} \right\rfloor = 2 + \left\lfloor \frac{3}{11} \right\rfloor = 2$$

$$\left\lfloor 10 + \frac{1}{7} \right\rfloor = 10 + \left\lfloor \frac{1}{7} \right\rfloor = 10$$

## Observation

$$\left\lfloor \frac{p}{q} \right\rfloor = 0 \quad \text{whenever } 0 \leq p < q$$

## Observation

$$\lfloor n + x \rfloor = n + \lfloor x \rfloor \quad \text{whenever } n \text{ is integer}$$

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

|

## Observation

*The floor (round-down) of  $\frac{p}{q}$  is computed as*

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

*where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .*

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder.

## Observation

*The floor (round-down) of  $\frac{p}{q}$  is computed as*

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

*where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .*

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder.

## Observation

*The floor (round-down) of  $\frac{p}{q}$  is computed as*

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

*where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .*

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder.

## Observation

*The floor (round-down) of  $\frac{p}{q}$  is computed as*

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

*where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .*



## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

## Observation

*The floor (round-down) of  $\frac{p}{q}$  is computed as*

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

*where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .*

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder. Try:

$$\begin{array}{rcl} 0 \cdot 3 & = & 0 \\ 1 \cdot 3 & = & 3 \end{array}$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder. Try:

|             |     |   |
|-------------|-----|---|
| $0 \cdot 3$ | $=$ | 0 |
| $1 \cdot 3$ | $=$ | 3 |
| $2 \cdot 3$ | $=$ | 6 |

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor =$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + ?}{3} \right\rfloor$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + ?}{3} \right\rfloor$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .



## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\begin{aligned}\left\lfloor \frac{8}{3} \right\rfloor &= \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor \\ &= \left\lfloor \frac{2 \cdot 3}{3} + \frac{2}{3} \right\rfloor\end{aligned}$$

Divide 8 by 3 with remainder. Try:

$$\begin{aligned}0 \cdot 3 &= 0 \\ 1 \cdot 3 &= 3 \\ 2 \cdot 3 &= 6 \\ 3 \cdot 3 &= 9 > 8\end{aligned}$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

$$= \left\lfloor \frac{2 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \right\rfloor$$

$$= \left\lfloor 2 + \frac{2}{3} \right\rfloor$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

$$= \left\lfloor \frac{2 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \right\rfloor$$

$$= \left\lfloor 2 + \frac{2}{3} \right\rfloor = 2$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

$$\text{because } 2 \leq 2 + \frac{2}{3} < 3$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

$$= \left\lfloor \frac{2 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \right\rfloor$$

$$= \left\lfloor 2 + \frac{2}{3} \right\rfloor = 2$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

because  $2 \leq 2 + \frac{2}{3} < 3$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

$$= \left\lfloor \frac{2 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \right\rfloor$$

$$= \left\lfloor 2 + \frac{2}{3} \right\rfloor = 2$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

because  $2 \leq 2 + \frac{2}{3} < 3$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) of  $\frac{8}{3}$ .

$$\left\lfloor \frac{8}{3} \right\rfloor = \left\lfloor \frac{2 \cdot 3 + 2}{3} \right\rfloor$$

$$= \left\lfloor \frac{2 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \right\rfloor$$

$$= \left\lfloor 2 + \frac{2}{3} \right\rfloor = 2$$

Divide 8 by 3 with remainder. Try:

$$0 \cdot 3 = 0$$

$$1 \cdot 3 = 3$$

$$2 \cdot 3 = 6$$

$$3 \cdot 3 = 9 > 8$$

$$\text{because } 2 \leq 2 + \frac{2}{3} < 3$$

## Observation

The floor (round-down) of  $\frac{p}{q}$  is computed as

$$\left\lfloor \frac{p}{d} \right\rfloor = q,$$

where  $q$  is the the quotient obtained by integer division of  $p$  by  $d$ .

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{4}{3} \right\rfloor = ?$$
$$\left\lfloor \frac{15}{2} \right\rfloor =$$
$$\left\lfloor \frac{24}{4} \right\rfloor =$$
$$\left\lfloor \frac{43}{5} \right\rfloor =$$
$$\left\lfloor \frac{56}{7} \right\rfloor =$$
$$\left\lfloor \frac{79}{8} \right\rfloor =$$
$$\left\lfloor \frac{80}{9} \right\rfloor =$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{4}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1$$

$$\left\lfloor \frac{15}{2} \right\rfloor =$$

$$\left\lfloor \frac{24}{4} \right\rfloor =$$

$$\left\lfloor \frac{43}{5} \right\rfloor =$$

$$\left\lfloor \frac{56}{7} \right\rfloor =$$

$$\left\lfloor \frac{79}{8} \right\rfloor =$$

$$\left\lfloor \frac{80}{9} \right\rfloor =$$



## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{4}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1$$

$$\left\lfloor \frac{15}{2} \right\rfloor = ?$$

$$\left\lfloor \frac{24}{4} \right\rfloor =$$

$$\left\lfloor \frac{43}{5} \right\rfloor =$$

$$\left\lfloor \frac{56}{7} \right\rfloor =$$

$$\left\lfloor \frac{79}{8} \right\rfloor =$$

$$\left\lfloor \frac{80}{9} \right\rfloor =$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\left\lfloor \frac{4}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1$$

$$\left\lfloor \frac{15}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7$$

$$\left\lfloor \frac{24}{4} \right\rfloor = ?$$

$$\left\lfloor \frac{43}{5} \right\rfloor =$$

$$\left\lfloor \frac{56}{7} \right\rfloor =$$

$$\left\lfloor \frac{79}{8} \right\rfloor =$$

$$\left\lfloor \frac{80}{9} \right\rfloor =$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned} \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\ \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\ \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\ \left\lfloor \frac{43}{5} \right\rfloor &= \\ \left\lfloor \frac{56}{7} \right\rfloor &= \\ \left\lfloor \frac{79}{8} \right\rfloor &= \\ \left\lfloor \frac{80}{9} \right\rfloor &= \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= ? \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\
 \left\lfloor \frac{56}{7} \right\rfloor &= ? \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}\left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\ \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\ \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\ \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\ \left\lfloor \frac{56}{7} \right\rfloor &= \left\lfloor \frac{8 \cdot 7}{7} \right\rfloor = \lfloor 8 \rfloor = 8 \\ \left\lfloor \frac{79}{8} \right\rfloor &= \\ \left\lfloor \frac{80}{9} \right\rfloor &= \end{aligned}$$



## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \left\lfloor \frac{8 \cdot 7}{7} \right\rfloor = \lfloor 8 \rfloor = 8 \\
 \left\lfloor \frac{79}{8} \right\rfloor &= ? \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \left\lfloor \frac{8 \cdot 7}{7} \right\rfloor = \lfloor 8 \rfloor = 8 \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \left\lfloor \frac{9 \cdot 8 + 7}{8} \right\rfloor = \left\lfloor \frac{9 \cdot 8}{8} + \frac{7}{8} \right\rfloor = \left\lfloor 9 + \frac{7}{8} \right\rfloor = 9 \\
 \left\lfloor \frac{80}{9} \right\rfloor &=
 \end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}\left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\ \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\ \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\ \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\ \left\lfloor \frac{56}{7} \right\rfloor &= \left\lfloor \frac{8 \cdot 7}{7} \right\rfloor = \lfloor 8 \rfloor = 8 \\ \left\lfloor \frac{79}{8} \right\rfloor &= \left\lfloor \frac{9 \cdot 8 + 7}{8} \right\rfloor = \left\lfloor \frac{9 \cdot 8}{8} + \frac{7}{8} \right\rfloor = \left\lfloor 9 + \frac{7}{8} \right\rfloor = 9 \\ \left\lfloor \frac{80}{9} \right\rfloor &= ?\end{aligned}$$

## Example

Compute the floor (round-down) function.

$$\begin{aligned}
 \left\lfloor \frac{4}{3} \right\rfloor &= \left\lfloor \frac{1 \cdot 3 + 1}{3} \right\rfloor = \left\lfloor \frac{1 \cdot 3}{3} + \frac{1}{3} \right\rfloor = \left\lfloor 1 + \frac{1}{3} \right\rfloor = 1 \\
 \left\lfloor \frac{15}{2} \right\rfloor &= \left\lfloor \frac{7 \cdot 2 + 1}{2} \right\rfloor = \left\lfloor \frac{7 \cdot 2}{2} + \frac{1}{2} \right\rfloor = \left\lfloor 7 + \frac{1}{2} \right\rfloor = 7 \\
 \left\lfloor \frac{24}{4} \right\rfloor &= \left\lfloor \frac{6 \cdot 4}{4} \right\rfloor = \lfloor 6 \rfloor = 6 \\
 \left\lfloor \frac{43}{5} \right\rfloor &= \left\lfloor \frac{8 \cdot 5 + 3}{5} \right\rfloor = \left\lfloor \frac{8 \cdot 5}{5} + \frac{3}{5} \right\rfloor = \left\lfloor 8 + \frac{3}{5} \right\rfloor = 8 \\
 \left\lfloor \frac{56}{7} \right\rfloor &= \left\lfloor \frac{8 \cdot 7}{7} \right\rfloor = \lfloor 8 \rfloor = 8 \\
 \left\lfloor \frac{79}{8} \right\rfloor &= \left\lfloor \frac{9 \cdot 8 + 7}{8} \right\rfloor = \left\lfloor \frac{9 \cdot 8}{8} + \frac{7}{8} \right\rfloor = \left\lfloor 9 + \frac{7}{8} \right\rfloor = 9 \\
 \left\lfloor \frac{80}{9} \right\rfloor &= \left\lfloor \frac{8 \cdot 9 + 8}{9} \right\rfloor = \left\lfloor \frac{8 \cdot 9}{9} + \frac{8}{9} \right\rfloor = \left\lfloor 8 + \frac{8}{9} \right\rfloor = 8
 \end{aligned}$$