

Calculus I

Trigonometry review

Todor Milev

2019

Outline

1 Trigonometry

- Angles
- The Trigonometric Functions
- Trigonometric Identities
- Graphs of the Trigonometric Functions

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\begin{aligned}\pi \text{ rad} &= 180^{\circ} \\ 1 \text{ rad} &= \frac{180^{\circ}}{\pi} \approx 57.3^{\circ}\end{aligned}$$

- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

$$1 \text{ rad} = \frac{180^{\circ}}{\pi} \approx 57.3^{\circ}$$

$$1^{\circ} = \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.$$

- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

$$1 \text{ rad} = \frac{180^{\circ}}{\pi} \approx 57.3^{\circ}$$

$$1^{\circ} = \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.$$


- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .
- Degrees are useful because the most frequently encountered fractions of a half turn are measured by a whole number of degrees.

Degrees and radians

- Degrees is a unit for measuring angles, denoted by $^{\circ}$.
- The relationship between degrees and radians is:


$$\begin{aligned}\pi \text{ rad} &= 180^{\circ} \\ 1 \text{ rad} &= \frac{180^{\circ}}{\pi} \approx 57.3^{\circ} \\ 1^{\circ} &= \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.\end{aligned}$$

- In other words, a half-turn is measured by $\pi \text{ rad}$ or 180° .
- Degrees are useful because the most frequently encountered fractions of a half turn are measured by a whole number of degrees.
- If a measurement unit is not specified, it is implied to be radians. For example, in $\sin 5$, the number 5 stands for 5 radians.




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	?										




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0										




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	?									




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$									




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$?								




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$								




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$?							




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$							




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$?						




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$						




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$?					




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$					




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$?				




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$				




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$?			




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$			




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$?		




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π		




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	?	




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$?




The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{3\pi}{2}$	2π




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.



- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.





- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.





- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.
- For example to determine in which quadrant is an angle of k radians located one needs to know the numerical value of $\frac{k}{\pi}$, which requires knowledge of π with great numerical accuracy.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta$ $\sin \theta$ $\tan \theta$	$0 < \theta < \frac{\pi}{2}$ $\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta$ $\tan \theta$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

 <p> $\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta$ </p>	<p>$0 < \theta < \frac{\pi}{2}$</p>  <p> $\cos \theta$ $\sin \theta$ $\tan \theta$ </p>
<p>sec θ csc θ cot θ</p> <p>All angles</p>	<p>sec θ csc θ cot θ</p> <p>Acute angles</p>

- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta$ $\csc \theta$ $\cot \theta$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

<p>Diagram showing an angle θ in standard position. The terminal arm passes through point (x, y). The distance from the origin to (x, y) is r. The horizontal distance is x (green) and the vertical distance is y (blue). A right angle is shown at (x, y).</p>	<p>Diagram showing a right triangle for an acute angle θ. The hypotenuse is the terminal arm, the adjacent side is the horizontal distance, and the opposite side is the vertical distance. A right angle is shown at the vertex of the adjacent and opposite sides.</p>
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$0 < \theta < \frac{\pi}{2}$ $\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta$ $\csc \theta$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

 $\cos \theta = \frac{x}{r} \qquad \sec \theta = \frac{r}{x}$ $\sin \theta = \frac{y}{r} \qquad \csc \theta = \frac{r}{y}$ $\tan \theta = \frac{y}{x} \qquad \cot \theta = \frac{x}{y}$ <p style="text-align: center;">All angles</p>	 $0 < \theta < \frac{\pi}{2}$ $\cos \theta \qquad \sec \theta$ $\sin \theta \qquad \csc \theta$ $\tan \theta \qquad \cot \theta$ <p style="text-align: center;">Acute angles</p>
--	--



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

 <p>Diagram showing an angle θ in standard position on a Cartesian coordinate system. The terminal arm passes through point (x, y) and has length r. A right triangle is formed with legs of length x and y.</p>	 <p>Diagram showing a right triangle with angle θ. The sides are labeled: adjacent (horizontal), opposite (vertical), and hypotenuse (slanted). The angle θ is at the vertex between the adjacent and hypotenuse sides.</p>
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$ $\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\cos \theta$ $\sin \theta$ $\tan \theta$ $\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of **acute θ (between 0 and $\frac{\pi}{2}$)** can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta$ $\sin \theta$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of **sides of right angle triangle** with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta$ $\csc \theta$ $\cot \theta$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles



- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$ $\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$ $\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles


- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Trigonometric Functions and Right Angle Triangles

	
$\cos \theta = \frac{x}{r}$ $\sin \theta = \frac{y}{r}$ $\tan \theta = \frac{y}{x}$	$\cos \theta = \frac{\text{adj}}{\text{hyp}}$ $\sin \theta = \frac{\text{opp}}{\text{hyp}}$ $\tan \theta = \frac{\text{opp}}{\text{adj}}$
$\sec \theta = \frac{r}{x}$ $\csc \theta = \frac{r}{y}$ $\cot \theta = \frac{x}{y}$	$\sec \theta = \frac{\text{hyp}}{\text{adj}}$ $\csc \theta = \frac{\text{hyp}}{\text{opp}}$ $\cot \theta = \frac{\text{adj}}{\text{opp}}$
All angles	Acute angles


- The trigonometric functions can be defined without requesting that the pt. (x, y) on the terminal arm of the angle lie on the unit circle.
- To do so we rescale by the distance r from the origin.
- The trig functions of acute θ (between 0 and $\frac{\pi}{2}$) can be interpreted as ratios of sides of right angle triangle with angle θ .

Geometric interpretation of all trigonometric functions




Fix unit circle, center O , coordinates $(0, 0)$.

Geometric interpretation of all trigonometric functions




Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$.

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A .

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$\sin \theta$

$\cos \theta$


$\tan \theta$

$\cot \theta$

$\sec \theta$

$\csc \theta$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$\sin \theta$

$\cos \theta$


$\tan \theta$

$\cot \theta$

$\sec \theta$

$\csc \theta$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1}$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1}$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1}$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

Geometric interpretation of all trigonometric functions



$$\beta = ?$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$\angle OED = ?$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0,0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

Geometric interpretation of all trigonometric functions



Fix unit circle, center O , coordinates $(0, 0)$.
 Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$


$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

$$\angle OED = ?$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = ?$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\angle OED = 180^\circ - 90^\circ - \beta$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta)\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$\sec \theta$

$\csc \theta$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|OD|}{|DE|} = \frac{1}{|DE|}$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|OD|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1}$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OB|}{|AC|} = \frac{1}{\sin \theta}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1}$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Geometric interpretation of all trigonometric functions



$$\begin{aligned}\beta &= 180^\circ - 90^\circ - \theta \\ &= 90^\circ - \theta\end{aligned}$$

$$\begin{aligned}\angle OED &= 180^\circ - 90^\circ - \beta \\ &= 90^\circ - (90^\circ - \theta) \\ &= \theta\end{aligned}$$

Fix unit circle, center O , coordinates $(0, 0)$.
Let $\angle DOB = \theta$. Let OB intersect the circle at point A . Coordinates of A are $(\cos \theta, \sin \theta)$.

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{|AC|}{|OA|} = \frac{|AC|}{1} = |AC|$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{|OC|}{|OA|} = \frac{|OC|}{1} = |OC|$$


$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{|BD|}{|OD|} = \frac{|BD|}{1} = |BD|$$

$$\cot \theta = \frac{\text{adj}}{\text{opp}} = \frac{|DE|}{|OD|} = \frac{|DE|}{1} = |DE|$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{|OB|}{|OD|} = \frac{|OB|}{1} = |OB|$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{|OE|}{|DO|} = \frac{|OE|}{1} = |OE|$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) =$$

$$\cos\left(\frac{2\pi}{3}\right) =$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = ?$$

$$\csc\left(\frac{2\pi}{3}\right) =$$


$$\cos\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) =$$


$$\cos\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = ?$$

$$\csc\left(\frac{2\pi}{3}\right) = \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2}$$

$$\cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$


$$\tan\left(\frac{2\pi}{3}\right) =$$

$$\csc\left(\frac{2\pi}{3}\right) =$$

$$\sec\left(\frac{2\pi}{3}\right) =$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$
$$\csc\left(\frac{2\pi}{3}\right) = \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = ?$$
$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) = \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\text{csc}\left(\frac{2\pi}{3}\right) = ? \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\text{csc}\left(\frac{2\pi}{3}\right) = \frac{2}{\sqrt{3}} \quad \sec\left(\frac{2\pi}{3}\right) =$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example



Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$


$$\sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2} \quad \cos\left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$\csc\left(\frac{2\pi}{3}\right) = \frac{2}{\sqrt{3}} \quad \sec\left(\frac{2\pi}{3}\right) = ?$$

$$\tan\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{-1} = -\sqrt{3}$$

$$\cot\left(\frac{2\pi}{3}\right) =$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= \end{aligned}$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= ? \end{aligned}$$

Example




Find the exact values of the trigonometric functions of

$$\theta = \frac{2\pi}{3} = 120^\circ.$$

$$\begin{aligned} \sin\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{2} & \cos\left(\frac{2\pi}{3}\right) &= -\frac{1}{2} & \tan\left(\frac{2\pi}{3}\right) &= \frac{\sqrt{3}}{-1} = -\sqrt{3} \\ \csc\left(\frac{2\pi}{3}\right) &= \frac{2}{\sqrt{3}} & \sec\left(\frac{2\pi}{3}\right) &= -\frac{2}{1} = -2 & \cot\left(\frac{2\pi}{3}\right) &= -\frac{1}{\sqrt{3}} \end{aligned}$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.

$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.

$$\sin \theta =$$

$$\tan \theta =$$


$$\csc \theta =$$

$$\sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = ?$, so $x = ?$.


$$\sin \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = ? \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta =$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = ?$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = ? \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta =$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = ?$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta =$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .



- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.


$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$

$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta = ?$$

Example

If $\cos \theta = \frac{2}{5}$ and $0 < \theta < \frac{\pi}{2}$, find the other five trigonometric functions of θ .




- Label the hypotenuse with length 5 and the adjacent side with length 2.
- Pythagorean theorem: $x^2 + 2^2 = 5^2$.
- Therefore $x^2 = 21$, so $x = \sqrt{21}$.

$$\sin \theta = \frac{\sqrt{21}}{5} \quad \tan \theta = \frac{\sqrt{21}}{2}$$


$$\csc \theta = \frac{5}{\sqrt{21}} \quad \sec \theta = \frac{5}{2}$$

$$\cot \theta = \frac{2}{\sqrt{21}}$$




- Positive angles are obtained by rotating counterclockwise.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.
- \sin is an **odd function**.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- Positive angles are obtained by rotating counterclockwise.
- Negative angles are obtained by rotating clockwise.
- If (x, y) is on the terminal arm of the angle θ , then $(x, -y)$ is on the terminal arm of $-\theta$.
- $\sin(-\theta) = \frac{-y}{r} = -\frac{y}{r} = -\sin \theta$.
- $\cos(-\theta) = \frac{x}{r} = \cos \theta$.
- \sin is an odd function.
- \cos is an even function.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- 2π represents a full rotation.

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$




- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .
- $\sin(\theta + 2\pi) = \sin \theta$.
- $\cos(\theta + 2\pi) = \cos \theta$.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$



- 2π represents a full rotation.
- $\theta + 2\pi$ has the same terminal arm as θ .
- $\theta + 2\pi$ uses the same point (x, y) and the same length r .
- $\sin(\theta + 2\pi) = \sin \theta$.
- $\cos(\theta + 2\pi) = \cos \theta$.
- We say \sin and \cos are 2π -periodic.

$$\begin{array}{ll} \sin \theta = \frac{y}{r} & \csc \theta = \frac{r}{y} \\ \cos \theta = \frac{x}{r} & \sec \theta = \frac{r}{x} \\ \tan \theta = \frac{y}{x} & \cot \theta = \frac{x}{y} \end{array}$$

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

Trigonometric Identities

Definition (Trigonometric Identity)

A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.


- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.

Trigonometric Identities

Definition (Trigonometric Identity)


A trigonometric identity is an equality between the trigonometric functions in one or more variables that holds for all values of the involved variables in the domains of all of the expressions.

- By convention, when dealing with trigonometric identities we do not account for the domains of the involved expressions.
- For example, $\frac{\sin \theta}{\sin \theta} = 1$ is considered a valid trigonometric identity, although, when considered as a function, the left hand side is not defined for $\theta \neq 0$.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

- $\csc \theta = \frac{1}{\sin \theta}$
- $\sec \theta = \frac{1}{\cos \theta}$
- $\cot \theta = \frac{1}{\tan \theta}$
- $\tan \theta = \frac{\sin \theta}{\cos \theta}$
- $\cot \theta = \frac{\cos \theta}{\sin \theta}$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\sin^2 \theta + \cos^2 \theta$$

$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2}\end{aligned}$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$


$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

$$\begin{aligned}& \sin^2 \theta + \cos^2 \theta \\ &= \frac{y^2}{r^2} + \frac{x^2}{r^2} \\ &= \frac{y^2 + x^2}{r^2} \\ &= \frac{r^2}{r^2} \\ &= 1\end{aligned}$$

Therefore $\sin^2 \theta + \cos^2 \theta = 1$.




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\sin^2 \theta + \cos^2 \theta = 1$$




$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta}\end{aligned}$$



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$

Example ($\tan^2 \theta + 1 = \sec^2 \theta$)

Prove the identity

$$\tan^2 \theta + 1 = \sec^2 \theta.$$

$$\begin{aligned}\sin^2 \theta + \cos^2 \theta &= 1 \\ \frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} &= \frac{1}{\cos^2 \theta} \\ \tan^2 \theta + 1 &= \sec^2 \theta\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

Substitute $-y$ for y , and use the fact that $\sin(-y) = -\sin y$ and $\cos(-y) = \cos y$:

$$\begin{aligned}\sin(x - y) &= \sin x \cos y - \cos x \sin y \\ \cos(x - y) &= \cos x \cos y + \sin x \sin y\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute x for y :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute x for y :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using $\cos^2 x = 1 - \sin^2 x$ and $\sin^2 x = 1 - \cos^2 x$:

$$\cos(2x) = 2 \cos^2 x - 1$$

$$\cos(2x) = 1 - 2 \sin^2 x$$

The remaining identities are consequences of the addition formulas:

$$\sin(x + y) = \sin x \cos y + \cos x \sin y$$

$$\cos(x + y) = \cos x \cos y - \sin x \sin y$$

To get the double angle formulas, substitute x for y :

$$\sin(2x) = 2 \sin x \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x$$

Rewrite the second double angle formula in two ways, using $\cos^2 x = 1 - \sin^2 x$ and $\sin^2 x = 1 - \cos^2 x$:

$$\cos(2x) = 2 \cos^2 x - 1$$

$$\cos(2x) = 1 - 2 \sin^2 x$$

To get the half-angle formulas, solve these equations for $\cos^2 x$ and $\sin^2 x$ respectively.

$$\cos^2 x = \frac{1 + \cos(2x)}{2}, \quad \sin^2 x = \frac{1 - \cos(2x)}{2}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$

Divide the first equation by the second, and then cancel $\cos x \cos y$ from the top and bottom:

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

The remaining identities are consequences of the addition formulas:

$$\begin{aligned}\sin(x + y) &= \sin x \cos y + \cos x \sin y \\ \cos(x + y) &= \cos x \cos y - \sin x \sin y\end{aligned}$$


Divide the first equation by the second, and then cancel $\cos x \cos y$ from the top and bottom:

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$


Do the same for the subtraction formulas:

$$\tan(x - y) = \frac{\tan x - \tan y}{1 + \tan x \tan y}$$


Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$?								




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0								




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	?							




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$							




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$?						




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1						




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	?					




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$					




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$?				




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0				




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	?			




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$			




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$?		




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1		




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	?	




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$?




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0




Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0



Graph of $\sin x$




x	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	2π
$\sin x$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0



The graph of $\sin x$ is 2π -periodic so the rest of the graph can be inferred from the interval $[0, 2\pi]$.


Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$?								




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1								




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	?							




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$							




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$?						




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0						




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	?					




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$					




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$?				




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1				




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	?			




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$			




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$?		




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0		




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	?	




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$?




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	0




Graph of $\cos x$




x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1



Graph of $\cos x$





x	0	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	π	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	2π
$\cos x$	1	$\frac{1}{2}$	0	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	0





The graph of $\cos x$ is 2π -periodic so the rest of the graph can be inferred from the interval $[0, 2\pi]$.



Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$





Graph of $\tan x$




Graph of $\tan x$



Graph of $\tan x$




Graph of $\tan x$




Near $\pm\frac{\pi}{2}$ the graph of $\tan x$ approaches $\pm\infty$.

Graph of $\tan x$



Near $\pm \frac{\pi}{2}$ the graph of $\tan x$ approaches $\pm \infty$. The graph of $\tan x$ is π -periodic so the rest of the graph can be inferred from the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$.


Graphs of the Trigonometric Functions



$$y = \sin x$$

$$y = \cos x$$

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right

Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$

- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right we get the graph of $\sin x$.


Graphs of the Trigonometric Functions




$$y = \sin x$$

$$y = \cos x$$


- $\sin x$ has zeroes at $n\pi$ for all integers n .
- $\cos x$ has zeroes at $\frac{\pi}{2} + n\pi$ for all integers n .
- $-1 \leq \sin x \leq 1$.
- $-1 \leq \cos x \leq 1$.
- If we translate the graph of $\cos x$ by $\frac{\pi}{2}$ units to the right we get the graph of $\sin x$. This is a consequence of $\cos\left(x - \frac{\pi}{2}\right) = \sin x$.




$$y = \tan x$$



$$y = \cot x$$




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis




$$y = \tan x$$




$$y = \cot x$$

If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis, we get the graph of $\cot x$.




$$y = \tan x$$




$$y = \cot x$$


If we move the graph of $\tan x$ by $\frac{\pi}{2}$ units to the left (or right) and reflect across the x axis, we get the graph of $\cot x$. This follows from $\tan\left(x \pm \frac{\pi}{2}\right) = -\cot x$.




$$y = \csc x$$




$$y = \sec x$$




$$y = \csc x$$



$$y = \sec x$$



$$y = \csc x$$



$$y = \sec x$$