

# Calculus I

## Antiderivatives, indefinite integrals and the Evaluation Theorem

Todor Milev

2019

# Outline

## 1 Antiderivatives

# Outline

## 1 Antiderivatives

## 2 Evaluating Definite Integrals

- The Evaluation Theorem (FTC part 2)
- Indefinite Integrals

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Antiderivatives

## Definition (Antiderivative)

A function  $F$  is called an antiderivative of  $f$  on an interval  $I$  if  $F'(x) = f(x)$  for all  $x$  in  $I$ .

## Example

- Let  $f(x) = x^2$ .

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \quad$ , then  $F'(x) = x^2 = f(x)$ .

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) =$  , then  $F'(x) = x^2 = f(x)$ .



## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \frac{1}{3}x^3$ , then  $F'(x) = x^2 = f(x)$ .

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \frac{1}{3}x^3$ , then  $F'(x) = x^2 = f(x)$ .
- Is this the only one?

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \frac{1}{3}x^3$ , then  $F'(x) = x^2 = f(x)$ .
- Is this the only one?
- No. If  $G(x) = \frac{1}{3}x^3 + 1$ , then  $G'(x) = x^2 = f(x)$ .

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \frac{1}{3}x^3$ , then  $F'(x) = x^2 = f(x)$ .
- Is this the only one?
- No. If  $G(x) = \frac{1}{3}x^3 + 1$ , then  $G'(x) = x^2 = f(x)$ .
- $\frac{1}{3}x^3 + 2$  will also work.

## Example

- Let  $f(x) = x^2$ .
- Use the Power Rule to find an antiderivative of  $f$ :
- If  $F(x) = \frac{1}{3}x^3$ , then  $F'(x) = x^2 = f(x)$ .
- Is this the only one?
- No. If  $G(x) = \frac{1}{3}x^3 + 1$ , then  $G'(x) = x^2 = f(x)$ .
- $\frac{1}{3}x^3 + 2$  will also work.
- Any function of the form  $H(x) = \frac{1}{3}x^3 + C$ , where  $C$  is a constant, is an antiderivative of  $f$ .

## Theorem

*If  $F$  is an antiderivative of  $f$  on an interval  $I$ , then an arbitrary antiderivative of  $f$  on  $I$  is of the form*

$$F(x) + C$$

*where  $C$  is an arbitrary constant.*

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) =$  , then  
 $F'(x) = \sin x$ .



## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) =$  , then  
 $F'(x) = \sin x$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  
 $F'(x) = \sin x$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  $F'(x) = \sin x$ .
- Therefore antiderivative is of the form  $G(x) = -\cos x + C$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  $F'(x) = \sin x$ .
- Therefore antiderivative is of the form  $G(x) = -\cos x + C$ .
- If  $F(x) = \frac{x^{n+1}}{n+1}$ , then  $F'(x) = x^n$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  $F'(x) = \sin x$ .
- Therefore antiderivative is of the form  $G(x) = -\cos x + C$ .
- If  $F(x) = \frac{x^{n+1}}{n+1}$ , then  $F'(x) = x^n$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  $F'(x) = \sin x$ .

- Therefore antiderivative is of the form  $G(x) = -\cos x + C$ .

- If  $F(x) = \frac{x^{n+1}}{n+1}$ , then  $F'(x) = x^n$ .

## Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If  $F(x) = -\cos x$ , then  $F'(x) = \sin x$ .
- Therefore antiderivative is of the form  $G(x) = -\cos x + C$ .

- If  $F(x) = \frac{x^{n+1}}{n+1}$ , then  $F'(x) = x^n$ .
- Therefore any antiderivative is of the form  $G(x) = \frac{x^{n+1}}{n+1} + C$ .

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .



## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) =$  , then  $F'(x) = \frac{1}{x}$ .

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) =$  , then  $F'(x) = \frac{1}{x}$ .

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .
- This is valid for any interval on which  $\frac{1}{x}$  is defined.

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .
- This is valid for any interval on which  $\frac{1}{x}$  is defined.
- $\frac{1}{x}$  is defined

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .
- This is valid for any interval on which  $\frac{1}{x}$  is defined.
- $\frac{1}{x}$  is defined everywhere except at 0.

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .
- This is valid for any interval on which  $\frac{1}{x}$  is defined.
- $\frac{1}{x}$  is defined everywhere except at 0.
- The most general answer needs two different constants, one for  $(-\infty, 0)$  and one for  $(0, \infty)$ .

## Example

Find the most general antiderivative of  $f(x) = \frac{1}{x}$ .

- If  $F(x) = \ln |x|$ , then  $F'(x) = \frac{1}{x}$ .
- This is valid for any interval on which  $\frac{1}{x}$  is defined.
- $\frac{1}{x}$  is defined everywhere except at 0.
- The most general answer needs two different constants, one for  $(-\infty, 0)$  and one for  $(0, \infty)$ .

$$G(x) = \begin{cases} \ln |x| + C_1 & \text{if } x > 0 \\ \ln |x| + C_2 & \text{if } x < 0 \end{cases}$$



Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula.  
Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	



Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula.  
Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	
$\sec x \tan x$	



Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose  $F' = f$  and  $G' = g$ .

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln  x $
$e^x$	$e^x$
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	$\sec x$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4 \quad + 2 \quad -$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4 \quad + 2 \quad -$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2x^5 - 2\sqrt{x} + C$$



## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \quad -$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} -$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} -$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}}$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}} + C$$

## Example

Find all functions  $g$  such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$\begin{aligned} g'(x) &= 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}} \\ g(x) &= 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}} + C \\ &= -4 \cos x + \frac{2}{5} x^5 - 2\sqrt{x} + C \end{aligned}$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$



## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) =$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) = \frac{x^{-1/2}}{-\frac{1}{2}}$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) = \frac{x^{-1/2}}{-\frac{1}{2}} + C$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

To find  $C$ , use the fact that  $f(1) = 1$ .

$$f(1) = 1$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

To find  $C$ , use the fact that  $f(1) = 1$ .

$$\begin{aligned} f(1) &= 1 \\ -\frac{2}{\sqrt{1}} + C &= 1 \end{aligned}$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

To find  $C$ , use the fact that  $f(1) = 1$ .

$$f(1) = 1$$

$$-\frac{2}{\sqrt{1}} + C = 1$$

$$C = 3$$

## Example

Find  $f$  if  $f'(x) = \frac{1}{x\sqrt{x}}$  for  $x > 0$ , and  $f(1) = 1$ .

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

To find  $C$ , use the fact that  $f(1) = 1$ .

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

$$f(1) = 1$$

$$-\frac{2}{\sqrt{1}} + C = 1$$

$$C = 3$$

Therefore

$$f(x) = -\frac{2}{\sqrt{x}} + 3.$$



## Theorem (The Evaluation Theorem (FTC part 2))

*If  $f$  is continuous on  $[a, b]$ , then*

$$\int_a^b f(x)dx = F(b) - F(a),$$

*where  $F$  is any antiderivative of  $f$ .*

$\int_a^b f(x)dx$  exists for any continuous (over  $[a, b]$ )  
function  $f$ .

### Theorem (The Evaluation Theorem (FTC part 2))

If  $f$  is continuous on  $[a, b]$ , then

$$\int_a^b f(x)dx = F(b) - F(a),$$

where  $F$  is any antiderivative of  $f$ .

## Theorem

*Let  $f$  be a continuous function on  $[a, b]$ . Then  $f$  is integrable over  $[a, b]$ .*

In other words,  $\int_a^b f(x)dx$  exists for any continuous (over  $[a, b]$ ) function  $f$ .


## Theorem (The Evaluation Theorem (FTC part 2))

*If  $f$  is continuous on  $[a, b]$ , then*

$$\int_a^b f(x)dx = F(b) - F(a),$$


*where  $F$  is any antiderivative of  $f$ .*

## Example



Evaluate the integral  $\int_{-2}^1 x^3 dx$ .


## Example



Evaluate the integral  $\int_{-2}^1 x^3 dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).


## Example



Evaluate the integral  $\int_{-2}^1 x^3 \, dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).
- An antiderivative is  $F(x) = ?$


## Example



Evaluate the integral  $\int_{-2}^1 x^3 dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).
- An antiderivative is  $F(x) = \frac{1}{4}x^4$ .

## Example




Evaluate the integral  $\int_{-2}^1 x^3 dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).
- An antiderivative is  $F(x) = \frac{1}{4}x^4$ .

$$\int_{-2}^1 x^3 dx = F(1) - F(-2)$$



## Example




Evaluate the integral  $\int_{-2}^1 x^3 dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).
- An antiderivative is  $F(x) = \frac{1}{4}x^4$ .

$$\int_{-2}^1 x^3 dx = F(1) - F(-2) = \frac{1}{4}(1)^4 - \frac{1}{4}(-2)^4$$

## Example



Evaluate the integral  $\int_{-2}^1 x^3 dx$ .

- $x^3$  is continuous on  $[-2, 1]$  (in fact, it's continuous everywhere).
- An antiderivative is  $F(x) = \frac{1}{4}x^4$ .

$$\int_{-2}^1 x^3 dx = F(1) - F(-2) = \frac{1}{4}(1)^4 - \frac{1}{4}(-2)^4 = \frac{1}{4} - \frac{16}{4} = -\frac{15}{4}$$

We often use the notation

$$F(x)]_a^b = F(b) - F(a)$$

or

$$[F(x)]_a^b = F(b) - F(a)$$

Therefore we can write


$$\int_a^b f(x)dx = F(x)]_a^b$$

or

$$\int_a^b f(x)dx = [F(x)]_a^b$$

## Example


Find the area under the parabola  $y = x^2$  from 0 to 1.



## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.


- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).



## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.


- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is ?



## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.


- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .



## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.

- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .




$$\int_0^1 x^2 \, dx = \left[ \frac{1}{3}x^3 \right]_0^1$$



## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.

- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .




$$\int_0^1 x^2 \, dx = \left[ \frac{1}{3}x^3 \right]_0^1$$

## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.

- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .




$$\int_0^1 x^2 dx = \left[ \frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3$$

## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.

- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .




$$\int_0^1 x^2 \, dx = \left[ \frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3$$

## Example

Find the area under the parabola  $y = x^2$  from 0 to 1.

- $x^2$  is continuous on  $[0, 1]$  (in fact, it's continuous everywhere).
- An antiderivative of  $x^2$  is  $\frac{1}{3}x^3$ .



$$\int_0^1 x^2 \, dx = \left[ \frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3 = \frac{1}{3}$$

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- **An antiderivative of  $\cos x$  is ?**

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- An antiderivative of  $\cos x$  is  $\sin x$ .



## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- An antiderivative of  $\cos x$  is  $\sin x$ .

$$\int_0^b \cos x \, dx = [\sin x]_0^b$$

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- **An antiderivative of  $\cos x$  is  $\sin x$ .**

$$\int_0^b \cos x \, dx = [\sin x]_0^b$$

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- An antiderivative of  $\cos x$  is  $\sin x$ .

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0)$$

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- An antiderivative of  $\cos x$  is  $\sin x$ .

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0)$$

## Example



Find the area under the cosine curve from 0 to  $b$ , where  $0 \leq b \leq \frac{\pi}{2}$ .

- $\cos x$  is continuous on  $[0, \frac{\pi}{2}]$  (in fact, it's continuous everywhere).
- An antiderivative of  $\cos x$  is  $\sin x$ .

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0) = \sin b$$

# Indefinite Integrals

- The Evaluation Theorem establishes a connection between antiderivatives and definite integrals.
- It says that  $\int_a^b f(x)dx$  equals  $F(b) - F(a)$ , where  $F$  is an antiderivative of  $f$ .
- We need convenient notation for writing antiderivatives.
- This is what the indefinite integral is.

## Definition (Indefinite Integral)

The indefinite integral of  $f$  is another way of saying the antiderivative of  $f$ , and is written  $\int f(x)dx$ . In other words,

$$\int f(x)dx = F(x) \quad \text{means} \quad F'(x) = f(x).$$

## Example

$$\int x^4 dx = ?$$

## Example

$$\int x^4 dx = \frac{x^5}{5}$$



## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left( \frac{x^5}{5} + C \right) = x^4.$$

## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left( \frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.

## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left( \frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of  $\frac{1}{x}$  is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left( \frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of  $\frac{1}{x}$  is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.

## Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left( \frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of  $\frac{1}{x}$  is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.
- $\int \frac{1}{x} dx = \ln|x| + C$ , and this is valid either on  $(-\infty, 0)$  or  $(0, \infty)$ .

## Example

Compute the integral.

$$\int (2x^2 - x - 5) \, dx$$

## Example

Compute the integral.

$$\int (2x^2 - x - 5) dx = 2 \int x^2 dx - \int x dx - 5 \int dx$$



## Example

Compute the integral.

$$\int (2x^2 - x - 5) dx = 2 \int x^2 dx - \int x dx - 5 \int dx$$

## Example

Compute the integral.

$$\int (2x^2 - x - 5) dx = 2 \int x^2 dx - \int x dx - 5 \int dx$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) dx &= 2 \int x^2 dx - \int x dx - 5 \int dx \\ &= 2? - ? - 5?\end{aligned}$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) dx &= 2 \int x^2 dx - \int x dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - ? - 5?\end{aligned}$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) \, dx &= 2 \int x^2 dx - \int x \, dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - ? - 5?\end{aligned}$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) \, dx &= 2 \int x^2 dx - \int x \, dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - \frac{x^2}{2} - 5x + C\end{aligned}$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) \, dx &= 2 \int x^2 dx - \int x dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - \frac{x^2}{2} - 5?\end{aligned}$$

## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) \, dx &= 2 \int x^2 dx - \int x dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - \frac{x^2}{2} - 5x\end{aligned}$$



## Example

Compute the integral.

$$\begin{aligned}\int (2x^2 - x - 5) \, dx &= 2 \int x^2 \, dx - \int x \, dx - 5 \int dx \\ &= 2 \frac{x^3}{3} - \frac{x^2}{2} - 5x + C\end{aligned}$$

## Example

Integrate.

$$\int 5x^{\frac{3}{2}} dx = 5$$

## Example

Integrate.

$$\int 5x^{\frac{3}{2}} dx = 5?$$

## Example

Integrate.

$$\int 5x^{\frac{3}{2}} dx = 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1}$$

## Example

Integrate.

$$\int 5x^{\frac{3}{2}} dx = 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1} + C$$

## Example

Integrate.

$$\begin{aligned}\int 5x^{\frac{3}{2}} dx &= 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1} + C \\ &= 5 \frac{x^{\frac{5}{2}}}{\frac{5}{2}} + C\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int 5x^{\frac{3}{2}} dx &= 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1} + C \\ &= \cancel{5} \frac{x^{\frac{5}{2}}}{\cancel{\frac{5}{2}}} + C \\ &= 2x^{\frac{5}{2}} + C\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int 5x^{\frac{3}{2}} dx &= 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1} + C \\ &= \cancel{5} \frac{x^{\frac{5}{2}}}{\cancel{5}} + C \\ &= 2x^{\frac{5}{2}} + C\end{aligned}$$



## Example

Integrate.

$$\begin{aligned}\int 5x^{\frac{3}{2}} dx &= 5 \frac{x^{\frac{3}{2}+1}}{\frac{3}{2}+1} + C \\ &= 5 \frac{x^{\frac{5}{2}}}{\frac{5}{2}} + C \\ &= 2x^{\frac{5}{2}} + C\end{aligned}$$

## Example

Integrate.

$$\int \frac{1}{\sqrt[3]{x^4}} dx$$

## Example

Integrate.

$$\int \frac{1}{\sqrt[3]{x^4}} dx = \int x^? dx$$

## Example

Integrate.

$$\int \frac{1}{\sqrt[3]{x^4}} dx = \int x^{-\frac{4}{3}} dx$$

## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\ &= ?\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\ &= \frac{x^{-\frac{4}{3}+1}}{-\frac{3}{4}+1}\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\ &= \frac{x^{-\frac{4}{3}+1}}{-\frac{3}{4}+1} + C\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\ &= \frac{x^{-\frac{4}{3}+1}}{-\frac{3}{4}+1} + C \\ &= \frac{x^{-\frac{1}{3}}}{-\frac{1}{3}} + C\end{aligned}$$



## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\&= \frac{x^{-\frac{4}{3}+1}}{-\frac{4}{3}+1} + C \\&= \frac{x^{-\frac{1}{3}}}{-\frac{1}{3}} + C \\&= -3x^{-\frac{1}{3}} + C\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \frac{1}{\sqrt[3]{x^4}} dx &= \int x^{-\frac{4}{3}} dx \\&= \frac{x^{-\frac{4}{3}+1}}{-\frac{4}{3}+1} + C \\&= \frac{x^{-\frac{1}{3}}}{-\frac{1}{3}} + C \\&= -3x^{-\frac{1}{3}} + C\end{aligned}$$

## Example

Integrate.

$$\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx$$

## Example

Integrate.

$$\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx = \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\ &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\ &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}} x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\ &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\ &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\ &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\ &= \int \left( x^3 - 2x^{\frac{3}{2} - \frac{1}{3}} + x^{-\frac{2}{3}} \right) dx\end{aligned}$$



## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\ &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\ &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2} - \frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \textcolor{red}{?} - 2? + ?\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2? \quad + ?\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2\text{?} + \text{?}\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + ?\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + ?\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1}\end{aligned}$$



## Example

Integrate.

$$\begin{aligned}\int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\&= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\&= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\&= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\&= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1} + C\end{aligned}$$

## Example

Integrate.

$$\begin{aligned}
 \int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\
 &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\
 &= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1} + C \\
 &= \frac{x^4}{4} - \frac{2x^{\frac{13}{6}}}{\frac{13}{6}} + \frac{x^{\frac{1}{3}}}{\frac{1}{3}} + C
 \end{aligned}$$

## Example

Integrate.

$$\begin{aligned}
 \int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\
 &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\
 &= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{\frac{-2}{3}+1}}{\frac{-2}{3}+1} + C \\
 &= \frac{x^4}{4} - \frac{2x^{\frac{13}{6}}}{\frac{13}{6}} + \frac{x^{\frac{1}{3}}}{\frac{1}{3}} + C
 \end{aligned}$$

## Example

Integrate.

$$\begin{aligned}
 \int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\
 &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\
 &= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1} + C \\
 &= \frac{x^4}{4} - \frac{2x^{\frac{13}{6}}}{\frac{13}{6}} + \frac{x^{\frac{1}{3}}}{\frac{1}{3}} + C \\
 &= \frac{x^4}{4} - \frac{12x^{\frac{13}{6}}}{13} + 3x^{\frac{1}{3}} + C
 \end{aligned}$$

## Example

Integrate.

$$\begin{aligned}
 \int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\
 &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\
 &= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1} + C \\
 &= \frac{x^4}{4} - \frac{2x^{\frac{13}{6}}}{\frac{13}{6}} + \frac{x^{\frac{1}{3}}}{\frac{1}{3}} + C \\
 &= \frac{x^4}{4} - \frac{12x^{\frac{13}{6}}}{13} + 3x^{\frac{1}{3}} + C
 \end{aligned}$$

## Example

Integrate.

$$\begin{aligned}
 \int \left( x^{\frac{3}{2}} - \frac{1}{x^{\frac{1}{3}}} \right)^2 dx &= \int \left( x^{\frac{3}{2}} - x^{-\frac{1}{3}} \right)^2 dx \\
 &= \int \left( \left( x^{\frac{3}{2}} \right)^2 - 2x^{\frac{3}{2}}x^{-\frac{1}{3}} + \left( x^{-\frac{1}{3}} \right)^2 \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{3}{2}-\frac{1}{3}} + x^{-\frac{2}{3}} \right) dx \\
 &= \int \left( x^3 - 2x^{\frac{7}{6}} + x^{-\frac{2}{3}} \right) dx \\
 &= \frac{x^4}{4} - 2 \frac{x^{\frac{7}{6}+1}}{\frac{7}{6}+1} + \frac{x^{-\frac{2}{3}+1}}{-\frac{2}{3}+1} + C \\
 &= \frac{x^4}{4} - \frac{2x^{\frac{13}{6}}}{\frac{13}{6}} + \frac{x^{\frac{1}{3}}}{\frac{1}{3}} + C \\
 &= \frac{x^4}{4} - \frac{12x^{\frac{13}{6}}}{13} + 3x^{\frac{1}{3}} + C
 \end{aligned}$$

## Example

Find the indefinite integral.

$$\int (8x^3 - 3 \sec^2 x) dx$$

## Example

Find the indefinite integral.

$$\int (8x^3 - 3 \sec^2 x) dx = 8 \int x^3 dx - 3 \int \sec^2 x dx$$



## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8? - 3?\end{aligned}$$

## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3?\end{aligned}$$

## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \text{?}\end{aligned}$$

## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x\end{aligned}$$

## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x + C\end{aligned}$$

## Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x + C \\ &= 2x^4 - 3 \tan x + C\end{aligned}$$

## Example

Find the general indefinite integral.

$$\int \frac{\cos \theta}{\sin^2 \theta} d\theta$$

## Example

Find the general indefinite integral.

$$\int \frac{\cos \theta}{\sin^2 \theta} d\theta = \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta$$



## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \textcolor{red}{?} \quad ? \quad d\theta\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta? \quad d\theta\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \text{?} \quad d\theta\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= ?\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= -\csc \theta\end{aligned}$$

## Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= -\csc \theta + C\end{aligned}$$

## Example

$$\int_0^3 (x^3 - 6x) dx$$



## Example

$$\int_0^3 (x^3 - 6x)dx = \left[ \int (x^3 - 6x)dx \right]_0^3$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\ &= \left[ \int x^3 dx - 6 \int x dx \right]_0^3\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[ \int (x^3 - 6x)dx \right]_0^3 \\ &= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[ ? - 6? \right]_0^3\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\ &= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[ \frac{x^4}{4} - 6x \right]_0^3\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[ \int (x^3 - 6x)dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6? \right]_0^3\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[ \int (x^3 - 6x)dx \right]_0^3 \\ &= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left( \frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left( \frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right)\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left( \frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left( \frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right)\end{aligned}$$



## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left( \frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left( \frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left( \frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left( \frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0\end{aligned}$$

## Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[ \int (x^3 - 6x) dx \right]_0^3 \\&= \left[ \int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[ \frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left( \frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left( \frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0 = -\frac{27}{4}.\end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

## Example

Evaluate: 
$$\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$$

$$= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9$$
$$= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9$$
$$= \left[ ? + ? - ? \right]_1^9$$



## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + ? - ? \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9$$
$$= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9$$
$$= \left[ t^2 + ? - ? \right]_1^9$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - ? \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - ? \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3}t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left( 9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left( 1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \end{aligned}$$



## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left( 9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left( 1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left( 9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left( 1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \\ &= 81 + 18 + \frac{1}{9} - 1 - \frac{2}{3} - 1 \end{aligned}$$

## Example

Evaluate:  $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left( 2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[ \int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[ \int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[ t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[ t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left( 9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left( 1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \\ &= 81 + 18 + \frac{1}{9} - 1 - \frac{2}{3} - 1 = \frac{868}{9}. \end{aligned}$$