#### **Precalculus**

# Equations involving logarithms and exponents

**Todor Miley** 

2019

Equations involving logarithms

- Equations involving logarithms
- Equations involving exponents

- Equations involving logarithms
- Equations involving exponents
- 3 Inverse function problems and exponents

- Equations involving logarithms
- Equations involving exponents
- 3 Inverse function problems and exponents
- Basic exponential inequalities

#### License to use and redistribute

These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
   https://creativecommons.org/licenses/by/3.0/us/and the links therein.

$$\log_3(2x^2+1) = 2$$

Solve the equation.

$$\log_3(2x^2+1) = 2$$

$$3^{\log_3(2x^2+1)} = 3^2$$

Solve the equation.

$$\log_3(2x^2 + 1) = 2 
3^{\log_3(2x^2+1)} = 3^2 
2x^2 + 1 = 9$$

Solve the equation.

$$\begin{array}{rcl} \log_3(2x^2+1) & = & 2 \\ 3^{\log_3(2x^2+1)} & = & 3^2 \\ 2x^2+1 & = & 9 \\ 2x^2 & = & 8 \end{array}$$

Solve the equation.

the equation.  

$$\log_{3}(2x^{2} + 1) = 2$$

$$3^{\log_{3}(2x^{2} + 1)} = 3^{2}$$

$$2x^{2} + 1 = 9$$

$$2x^{2} = 8$$

$$x^{2} = \frac{8}{2} = 4$$

Solve the equation.

the equation.
$$\log_{3}(2x^{2} + 1) = 2$$

$$3^{\log_{3}(2x^{2} + 1)} = 3^{2}$$

$$2x^{2} + 1 = 9$$

$$2x^{2} = 8$$

$$x^{2} = \frac{8}{2} = 4$$

$$x = \pm \sqrt{4} = \pm 2$$

$$\log_3(2x^2+1) = 2$$
 | Exponentiate base 3  $3^{\log_3(2x^2+1)} = 3^2$   $2x^2+1 = 9$   $2x^2 = 8$   $x^2 = \frac{8}{2} = 4$   $x = \pm\sqrt{4} = \pm 2$   $x = 2$  or  $x = -2$  | final answer

The logarithmic property  $\log_a(xy) = \log_a x + \log_a y$  holds only for positive x, y. Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

# Example

$$\log_2(x+2) + \log_2(x-1) = 2$$

The logarithmic property  $\log_a(xy) = \log_a x + \log_a y$  holds only for positive x, y. Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

#### Example

$$\log_2(x+2) + \log_2(x-1) = 2$$

$$\log_2((x+2)(x-1)) = 2$$

$$(x+2)(x-1) = 2^2$$

$$x^2 + x - 2 = 4$$

$$x^2 + x - 6 = 0$$

$$(x-2)(x+3) = 0$$

$$x = 2 \quad \text{or} \quad x = 3$$

$$x = -3 \text{ not a solution (outside of domain)}$$

$$16^{4t} = 8^{t-2}$$

$$16^{4t} = 8^{t-2}$$
  
Find a common base: (?)  $^{4t} = (?)^{t-2}$ 

$$\begin{array}{rcl} & 16^{4t} & = & 8^{t-2} \\ \text{Find a common base:} & \left(2^4\right)^{4t} & = & \left(2^3\right)^{t-2} \end{array}$$

Find a common base: 
$$(2^4)^{4t} = 8^{t-2}$$
  
 $2^{16t} = 2^{3t-6}$ 

Find a common base: 
$$(2^4)^{4t} = 8^{t-2}$$
  
 $2^{16t} = 2^{3t-6}$   
 $16t = 3t-6$ 

Find a common base: 
$$(2^4)^{4t} = 8^{t-2}$$
  
 $2^{16t} = 2^{3t-6}$   
 $16t = 3t-6$   
 $13t = -6$ 

$$2^{1-5x} = 12$$

Solve the equation.

$$2^{1-5x} = 12$$
  
 $\log_2(2^{1-5x}) = \log_2 12$ 

apply log<sub>2</sub>

Solve the equation.

$$2^{1-5x} = 12$$
  
 $\log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12$ 

apply log<sub>2</sub>

$$2^{1-5x} = 12$$
 apply  $\log_2 \log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = ?$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2 \log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4\cdot 3)$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2$   
 $\log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$   
 $1-5x = \log_2 4 + \log_2 3$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2 \log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4\cdot 3)$   
 $1-5x = \log_2 4 + \log_2 3$   
 $1-5x = ? + \log_2 3$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2 \log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$   
 $1-5x = \log_2 4 + \log_2 3$   
 $1-5x = 2 + \log_2 3$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2 \log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4\cdot 3)$   
 $1-5x = \log_2 4 + \log_2 3$   
 $1-5x = 2 + \log_2 3$   
 $5x = 1 - (2 + \log_2 3)$ 

$$2^{1-5x} = 12$$
 | apply  $\log_2$   
 $\log_2(2^{1-5x}) = \log_2 12$   
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$   
 $1-5x = \log_2 4 + \log_2 3$   
 $1-5x = 2 + \log_2 3$   
 $5x = 1 - (2 + \log_2 3)$   
 $x = \frac{-1}{2}$ 

$$\begin{array}{rclcrcl} 2^{1-5x} & = & 12 & & | \ \mathsf{apply} \ \mathsf{log}_2 \\ \mathsf{log}_2(2^{1-5x}) & = & \mathsf{log}_2 \ \mathsf{12} \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{12} = \mathsf{log}_2(4 \cdot 3) \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{4} + \mathsf{log}_2 \ \mathsf{3} \\ & 1-5x & = & 2+\mathsf{log}_2 \ \mathsf{3} \\ & 5x & = & 1-(2+\mathsf{log}_2 \ \mathsf{3}) \\ & x & = & \frac{-1-\mathsf{log}_2 \ \mathsf{3}}{2} \end{array}$$

$$\begin{array}{rclcrcl} 2^{1-5x} & = & 12 & & & | \operatorname{apply} \ \log_2(2^{1-5x}) & = & \log_2 12 & & \\ 1-5x & = & \log_2 12 = \log_2(4\cdot3) & & \\ 1-5x & = & \log_2 4 + \log_2 3 & & \\ 1-5x & = & 2 + \log_2 3 & & \\ 5x & = & 1 - (2 + \log_2 3) & & \\ x & = & \frac{-1 - \log_2 3}{5} & & & \end{array}$$

$$e^{x-3} = 2e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$
 Divide by  $e^{2x-1}$ 
 $\frac{e^{x-3}}{e^{2x-1}} = 2$ 

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

Divide by 
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

Divide by 
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

$$e^{-x-2} = 2$$

Divide by 
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

$$e^{-x-2} = 2$$

Divide by 
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$
 Divide by  $e^{2x-1}$   $\frac{e^{x-3}}{e^{2x-1}} = 2$   $e^{x-3-(2x-1)} = 2$   $e^{-x-2} = 2$  Apply In  $-x-2 = \ln 2$ 

$$e^{x-3} = 2e^{2x-1}$$
 Divide by  $e^{2x-1}$   $\frac{e^{x-3}}{e^{2x-1}} = 2$   $e^{x-3-(2x-1)} = 2$  Apply In  $-x-2 = \ln 2$ 

Solve the equation.

$$e^{x-3} = 2e^{2x-1}$$
 Divide by  $\frac{e^{x-3}}{e^{2x-1}} = 2$   $e^{x-3-(2x-1)} = 2$   $e^{-x-2} = 2$  Apply In  $-x-2 = \ln 2$   $-x = \ln 2 + 2$ 

Divide by 
$$e^{2x-1}$$

Apply In

$$e^{x-3} = 2e^{2x-1}$$
 Divide by  $e^{2x-1}$ 
 $\frac{e^{x-3}}{e^{2x-1}} = 2$ 
 $e^{x-3-(2x-1)} = 2$ 
 $e^{-x-2} = 2$  Apply In
 $-x-2 = \ln 2$ 
 $-x = \ln 2 + 2$ 
 $x = -(\ln 2 + 2)$ 

$$e^{x-3}=2e^{2x-1}$$
 Divide by  $e^{2x-1}$ 
 $\frac{e^{x-3}}{e^{2x-1}}=2$ 
 $e^{x-3-(2x-1)}=2$ 
 $e^{-x-2}=2$  Apply In
 $-x-2=\ln 2$ 
 $-x=\ln 2+2$ 
 $x=-(\ln 2+2)$ 
 $x=-\ln 2-2$  Final answer

$$e^{x-3}=2e^{2x-1}$$
 Divide by  $e^{2x-1}$ 
 $\frac{e^{x-3}}{e^{2x-1}}=2$ 
 $e^{x-3-(2x-1)}=2$ 
 $e^{-x-2}=2$  Apply In
 $-x-2=\ln 2$ 
 $-x=\ln 2+2$ 
 $x=-(\ln 2+2)$ 
 $x=-\ln 2-2$  Final answer
 $x\approx -2.693$  Calculator

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

Solve.

$$3^{2x+5}$$
 - 5.2<sup>-x+1</sup>

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$
$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$
$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} \ = \ 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = \frac{1}{2}$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = \frac{1}{2}$$

$$(\log_2 3)(2x + 5) + x - 1 = \log_2 5$$

$$x( + ) + = \log_2 5$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} \ = \ 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 3$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + ) + = \log_2 5$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} \ = \ 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + = \log_2 5$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 4$$

$$(\log_2 3)(2x+5)+x-1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 3$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1}{1}$$

Common base

$$a = b^{\log_b a}$$

Apply  $\log_2$ 

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1 - 5\log_2 3}{2}$$

#### Common base

$$a = b^{\log_b a}$$

Solve.

Solve. 
$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x\left(2\log_2 3 + 1\right) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1 - 5\log_2 3}{2\log_2 3 + 1}$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$
  
 $x = \frac{\log_2 5 + 1 - 5\log_2 3}{2\log_2 3 + 1}$ 

$$x \approx -1.1038$$

Common base

$$a = b^{\log_b a}$$

Apply log<sub>2</sub>

Calculator

$$e^{5-3x} = 10$$

$$e^{5-3x} = 10$$
 apply In In  $(e^{5-3x}) = \ln 10$ 

$$e^{5-3x}=10$$
 apply In  $\ln(e^{5-3x})=\ln 10$   $5-3x=\ln 10$ 

$$e^{5-3x} = 10$$
 apply In  $\ln(e^{5-3x}) = \ln 10$   
 $5-3x = \ln 10$   
 $3x = 5 - \ln 10$ 

$$e^{5-3x} = 10$$
 apply In  $\ln(e^{5-3x}) = \ln 10$   $5-3x = \ln 10$   $3x = 5 - \ln 10$   $x = \frac{5 - \ln 10}{3}$ 

$$e^{5-3x}=10$$
 apply In  $\ln(e^{5-3x})=\ln 10$   $5-3x=\ln 10$   $3x=5-\ln 10$   $x=\frac{5-\ln 10}{3}$  Calculator:  $x\approx 0.8991$ .

### Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for *t*:

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t) =

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t) = r(t)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for 
$$t$$
:  $c(t) = r(t)$ 

Equations involving logarithms and expo...

2019

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t =$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$ 

$$48 \cdot 2^{t} =$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t = 6 \cdot 4^t$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$   

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t = 6 \cdot 4^t$   
 $8 \cdot 2^t = 4^t$ 

Find a common base:  $2^{?} \cdot 2^{t} = 2^{?}$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t = 6 \cdot 4^t$   
 $8 \cdot 2^t = 4^t$ 

Find a common base:  $2^3 \cdot 2^t = 2^?$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t = 6 \cdot 4^t$   
 $8 \cdot 2^t = 4^t$ 

Find a common base:  $2^3 \cdot 2^t = 2^?$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for *t*: 
$$c(t) = r(t)$$
  
 $48 \cdot 2^{t} = 6 \cdot 4^{t}$   
 $8 \cdot 2^{t} = 4^{t}$ 

Find a common base:  $2^3 \cdot 2^t = 2^{2t}$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$   
 $48 \cdot 2^t = 6 \cdot 4^t$   
 $8 \cdot 2^t = 4^t$   
Find a common base:  $2^3 \cdot 2^t = 2^{2t}$   
 $2^{t+3} = 2^{2t}$ 

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$ 

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$
Find a common base:  $2^{3} \cdot 2^{t} = 2^{2t}$ 

$$2^{t+3} = 2^{2t}$$

$$t+3 = 2t$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for 
$$t$$
:  $c(t) = r(t)$ 

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$
Find a common base:  $2^{3} \cdot 2^{t} = 2^{2t}$ 

$$2^{t+3} = 2^{2t}$$

$$t+3=2t$$

$$t=3$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for 
$$t$$
:  $c(t) = r(t)$ 

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$
Find a common base:  $2^{3} \cdot 2^{t} = 2^{2t}$ 

$$2^{t+3} = 2^{2t}$$

$$t+3=2t$$

$$t=3$$

Therefore the chicken and rabbit populations are equal after 3 years.

$$9^x = 2 \cdot 3^x + 63$$

$$9^x = 2 \cdot 3^x + 63$$
$$9^x - 2 \cdot 3^x - 63 = 0$$

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $2^{x} - 2u - 63 = 0$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $2u - 63 = 0$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $u^{2} - 2u - 63 = 0$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $u^{2} - 2u - 63 = 0$   
 $(?)(?) = 0$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $u^{2} - 2u - 63 = 0$   
 $(u - 9)(u + 7) = 0$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
  
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$   
 $u^{2} - 2u - 63 = 0$   
 $(u - 9)(u + 7) = 0$   
 $u = 9 \text{ or } u = -7$   
 $3^{x} = 9 \text{ or } 3^{x} = -7$   
 $x = ?$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 
 $x = 2$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 
 $x = 2$ 

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 
 $x = 2 \text{ no real solution}$ 

Solve for x.

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 
 $x = 2$  no real solution

Therefore x = 2 is the solution.

#### Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set  $e^{x} = u$ .

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set  $e^x = u$ . Then  $e^{2x} = ?$ .

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set  $e^x = u$ . Then  $e^{2x} = u^2$ .

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2-3u-4=0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
(? ) (? ) = 0

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set  $e^x = u$ . Then  $e^{2x} = u^2$ .

$$u^2 - 3u - 4 = 0$$
  
(u-4)(u+1) = 0

$$u = 4$$

or

u = -1

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u = 4$$
 or  $u = -1$   
 $e^x = 4$  or  $e^x = -1$ 

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u=4$$
 or  $u=-1$   
 $e^x=4$  or  $e^x=-1$   
 $x=\ln 4$  or no real solution

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u=4$$
 or  $u=-1$   
 $e^x=4$  or  $e^x=-1$   
 $x=\ln 4$  or no real solution  
 $x\approx 1.3863$ 

$$4^{x+1} - 2^{x+2} - 3 = 0$$

## Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set u = ?.

### Solve the equation

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set  $u = 2^x$ .

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = ?$ ,

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = ?$ .

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .

$$4u^2 - 4u - 3 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .

$$4u^2 - 4u - 3 = 0$$

$$(?)$$
  $(?)$   $=$  0

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set 
$$u = 2^x$$
. Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .

$$4u^2 - 4u - 3 = 0$$

$$(2u-3)(2u+1) = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right)$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$
or no real solution

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln(?)}{\ln?}$$
or no real solution

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln\left(\frac{3}{2}\right)}{\ln 2} \text{ or no real solution}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln\left(\frac{3}{2}\right)}{\ln 2} \approx 0.58496 \text{ or no real solution}$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = 2$ 

 $3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$ 

## Example (Exponential equation that reduces to quadratic)

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1}$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1}$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$
$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$
$$4 - 2 - 63 \cdot 3^{-1} = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$
$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$
$$u - 2 - 63u^{-1} = 0$$

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(?)(?) = 0$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$   
Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$   
Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$
or  $u = -7$ 

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$
or 
$$u = -7$$
or no real solution

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$   
Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$
or no real solution

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

> 2 terms 
$$\Rightarrow$$
 transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

> 2 terms 
$$\Rightarrow$$
  
transfer one side  
 $3^{2x} = u$   
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$   
Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = ?$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = 2$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = 2$$

$$x = 1$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$  Multiply  $\cdot u$ 

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$


$$2x = 2$$

$$x = 1$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$  Multiply  $\cdot u$ 

Find 
$$f^{-1}(x)$$
 for

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .




y = f(x)

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .

$$\frac{e^x-e^{-x}}{e^x+e^{-x}}=y$$


Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{(u - ?)}{(u + ?)} = y$$

Set  $u = e^x$ 


Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{(u - ?)}{(u + ?)} = y$$

Set 
$$u = e^x$$
  
 $e^{-x} =$ ?


Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right)}{\left(u + \frac{1}{u}\right)} = y$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 


Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right) u}{\left(u + \frac{1}{u}\right) u} = y$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 


Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$
$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

Set 
$$u = e^{x}$$
  
 $e^{-x} = \frac{1}{e^{x}} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$


$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$


$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$


$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$


$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$


$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$\left(e^{x}\right)^{2} = \frac{1 + y}{1 - y}$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$


$$u^{2} = \frac{1 + y}{1 - y}$$

$$\left(e^{x}\right)^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$


$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$


$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



answer
$$f^{-1}(y) = \frac{1}{2} \ln \left( \frac{1+y}{1-y} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$


$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln\left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u})u}{(u + \frac{1}{u})u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$


$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln\left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Find 
$$f^{-1}(x)$$
 for 
$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$
 $f = \tanh = \text{hyperbolic}$ 

tangent function.



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$


$$x = \frac{1}{2}\ln\left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Take In


# Solve the inequality. $2^{-3x-5} < 7$

$$2^{-3x-5}$$
 < 7




# Solve the inequality. $2^{-3x-5}$ < 7

$$2^{-3x-5}$$
 < 7




# Solve the inequality. $2^{-3x-5} < 7$



Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\log_2 2^{-3x-5} \ < \ \log_2 7$$



Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\log_2 2^{-3x-5} < \log_2 7$$



### Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\begin{array}{rcl} \log_2 2^{-3x-5} & < & \log_2 7 \\ -3x-5 & < & \log_2 7 \end{array}$$



#### Solve the inequality.

$$2^{-3x-5}$$
 < 7

$$\begin{array}{rcl} \log_2 2^{-3x-5} & < & \log_2 7 \\ -3x-5 & < & \log_2 7 \\ -3x & < & \log_2 7 + 5 \end{array}$$



#### Solve the inequality.

$$2^{-3x-5} < 7$$


$$\log_{2} 2^{-3x-5} < \log_{2} 7$$

$$-3x-5 < \log_{2} 7$$

$$-3x < \log_{2} 7+5$$

$$x > -\frac{\log_{2} 7+5}{3}$$

Logarithms preserve inequalities: apply log<sub>2</sub>



#### Solve the inequality.

$$2^{-3x-5} < 7$$


$$\log_{2} 2^{-3x-5} < \log_{2} 7$$

$$-3x-5 < \log_{2} 7$$

$$-3x < \log_{2} 7+5$$

$$x > -\frac{\log_{2} 7+5}{3}$$

Logarithms preserve inequalities: apply log<sub>2</sub>



#### Solve the inequality.


$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$\begin{array}{rcl}
-3x - 5 & < \log_2 7 \\
-3x & < \log_2 7 + 5 \\
x & > -\frac{\log_2 7 + 5}{3}
\end{array}$$

$$x \in \left(-\frac{5 + \log_2 7}{3}, \infty\right)$$

Logarithms preserve inequalities: apply log<sub>2</sub>



#### Solve the inequality.


$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$\begin{array}{rcl}
-3x - 5 & < & \log_2 7 \\
-3x & < & \log_2 7 + 5 \\
x & > & -\frac{\log_2 7 + 5}{3}
\end{array}$$

$$x \in \left(-\frac{5 + \log_2 7}{3}, \infty\right)$$

Logarithms preserve inequalities: apply log<sub>2</sub>



# Solve the inequality. $2^{-3x-5} < 7$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$-3x - 5 < \log_2 7$$
  
 $-3x < \log_2 7 + 5$ 

$$x > -\frac{\log_2 7 + 5}{3}$$

Logarithms preserve inequalities: apply log<sub>2</sub>

