

# Calculus II

## Partial fractions

Todor Milev

2019

# Outline

- 1 Integration of Rational Functions
  - Partial fractions

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} =$$

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} =$$

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} \quad .$$

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} \quad .$$

- We can now solve the following integral:

$$\int \frac{x+5}{x^2+x-2} dx =$$

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} .$$

- We can now solve the following integral:

$$\int \frac{x+5}{x^2+x-2} dx = \int \left( \frac{2}{x-1} - \frac{1}{x+2} \right) dx =$$



# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} \quad .$$

- We can now solve the following integral:

$$\int \frac{x+5}{x^2+x-2} dx = \int \left( \frac{2}{x-1} - \frac{1}{x+2} \right) dx = 2 \ln |x-1| - \ln |x+2| + C$$

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} \quad .$$

- We can now solve the following integral:

$$\int \frac{x+5}{x^2+x-2} dx = \int \left( \frac{2}{x-1} - \frac{1}{x+2} \right) dx = 2 \ln |x-1| - \ln |x+2| + C$$

- From (linear substitutions of) basic building blocks we constructed a larger example, which we can therefore solve.

# From building blocks to all rational functions: example

- We know how to solve  $\int \frac{2}{x-1} dx$  and  $\int \frac{1}{x+2} dx$ .
- Consider the difference

$$\frac{2}{x-1} - \frac{1}{x+2} = \frac{2(x+2) - (x-1)}{(x-1)(x+2)} = \frac{x+5}{x^2+x-2} \quad .$$

- We can now solve the following integral:

$$\int \frac{x+5}{x^2+x-2} dx = \int \left( \frac{2}{x-1} - \frac{1}{x+2} \right) dx = 2 \ln |x-1| - \ln |x+2| + C$$

- From (linear substitutions of) basic building blocks we constructed a larger example, which we can therefore solve.
- We now learn how to do the reverse procedure: given a rational function, split it into “partial fractions”.

# Partial fractions definition

## Definition

A partial fraction is rational function of one of the 2 forms below.

- $\frac{A}{(ax+b)^n}, n \geq 1.$
- $\frac{Ax+B}{(ax^2+bx+c)^n},$  where  $b^2 - 4ac < 0$  and  $n \geq 1.$

# Partial fractions definition

## Definition

A partial fraction is rational function of one of the 2 forms below.

- $\frac{A}{(ax+b)^n}, n \geq 1.$
- $\frac{Ax+B}{(ax^2+bx+c)^n},$  where  $b^2 - 4ac < 0$  and  $n \geq 1.$

## Theorem

*Every rational function can be written as a sum of a polynomial and partial fractions.*

# Partial fractions definition

## Definition

A partial fraction is rational function of one of the 2 forms below.

- $\frac{A}{(ax+b)^n}, n \geq 1.$
- $\frac{Ax+B}{(ax^2+bx+c)^n},$  where  $b^2 - 4ac < 0$  and  $n \geq 1.$

## Theorem

*Every rational function can be written as a sum of a polynomial and partial fractions.*

- We already learned how to integrate all partial fractions (using linear substitutions and building blocks I, II and III).

# Partial fractions definition

## Definition

A partial fraction is rational function of one of the 2 forms below.

- $\frac{A}{(ax+b)^n}, n \geq 1.$
- $\frac{Ax+B}{(ax^2+bx+c)^n},$  where  $b^2 - 4ac < 0$  and  $n \geq 1.$

## Theorem

*Every rational function can be written as a sum of a polynomial and partial fractions.*

- We already learned how to integrate all partial fractions (using linear substitutions and building blocks I, II and III).
- Thus, if we can produce the partial fractions whose existence is promised by the theorem, we can integrate all rational functions.

# Review of polynomial notation

- Recall that a rational function is a function of the form

$$f(x) = \frac{P(x)}{Q(x)}$$

where  $P$  and  $Q \neq 0$  are polynomials.



# Review of polynomial notation

- Recall that a rational function is a function of the form

$$f(x) = \frac{P(x)}{Q(x)}$$

where  $P$  and  $Q \neq 0$  are polynomials.

- Recall that the degree of  $P$  is the highest power of  $x$  in  $P$  that has a non-zero coefficient.

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x), R(x)$  such that  $\deg R < \deg Q$  and

$$P(x) = S(x)Q(x) + R(x)$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the **dividend**  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x)$ ,  $R(x)$  such that  $\deg R < \deg Q$  and

$$P(x) = S(x)Q(x) + R(x)$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the **divisor**  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x)$ ,  $R(x)$  such that  $\deg R < \deg Q$  and

$$P(x) = S(x)Q(x) + R(x)$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get **quotient  $S(x)$**  with remainder  $R(x)$  means to find polynomials  $S(x)$ ,  $R(x)$  such that  $\deg R < \deg Q$  and

$$P(x) = \textcolor{red}{S(x)}Q(x) + R(x)$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with **remainder  $R(x)$**  means to find polynomials  $S(x)$ ,  $R(x)$  such that  **$\deg R < \deg Q$**  and

$$P(x) = S(x)Q(x) + R(x)$$



# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x), R(x)$  such that  $\deg R < \deg Q$  and

$$\begin{aligned} P(x) &= S(x)Q(x) + R(x) && | \text{ divide by } Q(x) \\ \frac{P(x)}{Q(x)} &= \frac{S(x)Q(x)}{Q(x)} + \frac{R(x)}{Q(x)} \end{aligned}$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x), R(x)$  such that  $\deg R < \deg Q$  and

$$\begin{aligned} P(x) &= S(x)Q(x) + R(x) && | \text{ divide by } Q(x) \\ \frac{P(x)}{Q(x)} &= \frac{S(x)\cancel{Q(x)}}{\cancel{Q(x)}} + \frac{R(x)}{Q(x)} \\ \frac{P(x)}{Q(x)} &= S(x) + \frac{R(x)}{Q(x)} \end{aligned}$$

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x), R(x)$  such that  $\deg R < \deg Q$  and

$$\begin{aligned} P(x) &= S(x)Q(x) + R(x) && | \text{ divide by } Q(x) \\ \frac{P(x)}{Q(x)} &= \frac{\cancel{S(x)Q(x)}}{\cancel{Q(x)}} + \frac{R(x)}{Q(x)} \\ \frac{P(x)}{Q(x)} &= S(x) + \frac{R(x)}{Q(x)} \end{aligned}$$

- The above transforms  $\frac{P(x)}{Q(x)}$  to a polynomial plus a fraction in which the numerator has degree smaller than the denominator.

# Ensure denominator degree $>$ numerator degree

- To decompose  $\frac{P(x)}{Q(x)}$  in partial fractions we ensure first the degree of the numerator is smaller than the degree of the denominator.
- We recall that to divide the dividend  $P(x)$  by the divisor  $Q(x)$  to get quotient  $S(x)$  with remainder  $R(x)$  means to find polynomials  $S(x)$ ,  $R(x)$  such that  $\deg R < \deg Q$  and

$$\begin{aligned} P(x) &= S(x)Q(x) + R(x) && | \text{ divide by } Q(x) \\ \frac{P(x)}{Q(x)} &= \frac{\cancel{S(x)Q(x)}}{\cancel{Q(x)}} + \frac{R(x)}{Q(x)} \\ \frac{P(x)}{Q(x)} &= S(x) + \frac{R(x)}{Q(x)} \end{aligned}$$

- The above transforms  $\frac{P(x)}{Q(x)}$  to a polynomial plus a fraction in which the numerator has degree smaller than the denominator.
- The polynomials  $Q(x)$  and  $S(x)$  are computed via polynomial long division. We recall the procedure through examples.

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r} x-1 \overline{) x^3 \phantom{+ 0x^2 + 0x} + x} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \phantom{+ 0} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \phantom{+ 0} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \phantom{+ 0} \end{array}$$

---

---

---

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r} x-1 \overline{) x^3 \phantom{+ 0x^2 + 0x} + x} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \\ \hline \phantom{x-1} \phantom{) x^3} \phantom{+ 0x^2} \phantom{+ 0x} \end{array}$$

Divide  $x^3$  by  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r} x^2 \\ x-1 \overline{) x^3 \phantom{+ 0x^2 + 0x} + x} \\ \underline{\phantom{x^3} x^2 \phantom{+ 0x} + x} \phantom{+ 0} \\ \phantom{x^3} \phantom{x^2} \phantom{+ 0x} \underline{\phantom{+ 0} x} \phantom{+ 0} \\ \phantom{x^3} \phantom{x^2} \phantom{+ 0x} \phantom{+ 0} \underline{\phantom{+ 0} 0} \phantom{+ 0} \end{array}$$

Divide  $x^3$  by  $x$



## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 \\
 x-1 \overline{) x^3 \phantom{+ 0x^2 + 0x} + x} \\
 \underline{\phantom{x^3} x^2 - x^2 + x} \phantom{+ 0} \\
 \phantom{x^3} \phantom{x^2} \phantom{+ 0x^2} 2x \phantom{+ 0} \\
 \underline{\phantom{x^3} \phantom{x^2} 2x - 2x + x} \\
 \phantom{x^3} \phantom{x^2} \phantom{+ 0x^2} \phantom{+ 0} x
 \end{array}$$

Multiply  $x^2$  by  $x - 1$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 \phantom{x^3 - } x^2 + x \\
 \phantom{x^3 - } \underline{x^2 - x} \\
 \phantom{x^3 - } \phantom{x^2 + } 2x
 \end{array}$$

Multiply  $x^2$  by  $x - 1$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r} x^2 \\ x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\ \underline{x^3 - x^2} \phantom{+ x} \\ \phantom{x^3 - } x^2 \phantom{+ x} \end{array}$$

Subtract  $x^3 - x^2$  from  $x^3$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 \phantom{+ x}
 \end{array}$$

Subtract  $x^3 - x^2$  from  $x^3$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 \\
 x-1 \overline{) x^3 \phantom{+ 0x^2 + 0x + 0} + \quad x} \\
 \underline{x^3 - x^2 \phantom{+ 0x + 0} \phantom{+ x}} \\
 x^2 \phantom{+ 0x + 0} + \quad x \\
 \underline{\phantom{x^2 + 0x + 0} \phantom{+ x}} \\
 \phantom{x^2 + 0x + 0} \phantom{+ x}
 \end{array}$$

Bring down the  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 0x} \\
 x^2 \phantom{+ 0x} + x \\
 \underline{\phantom{x^2 + 0x} x} \\
 \phantom{x^2 + 0x} 0
 \end{array}$$

Divide  $x^2$  by  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{\phantom{x^3 - x^2} x^2 + x} \\
 \phantom{x^3 - x^2} 0
 \end{array}$$

Divide  $x^2$  by  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{\phantom{x^3 - x^2} x^2 + x} \\
 \phantom{x^3 - x^2} \phantom{x^2 + x} 0
 \end{array}$$

Multiply  $x$  by  $x - 1$



## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x
 \end{array}$$

Multiply  $x$  by  $x - 1$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 \hline
 \phantom{x^2 + } 2x
 \end{array}$$

Subtract  $x^2 - x$  from  $x^2 + x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x
 \end{array}$$

Subtract  $x^2 - x$  from  $x^2 + x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 0x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x
 \end{array}$$

Divide  $2x$  by  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x
 \end{array}$$

Divide  $2x$  by  $x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x
 \end{array}$$

Multiply 2 by  $x - 1$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

Multiply 2 by  $x - 1$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

Subtract  $2x - 2$  from  $2x$



## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

Subtract  $2x - 2$  from  $2x$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

$$\begin{aligned}
 & \int \frac{x^3+x}{x-1} dx \\
 &= \int \left( x^2 + x + 2 + \frac{2}{x-1} \right) dx
 \end{aligned}$$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 \textcolor{red}{x^2} + \textcolor{red}{x} + \textcolor{red}{2} \\
 x-1 \overline{) x^3 \phantom{+ 0x^2} + 0x + 0} \\
 \underline{x^3 - x^2} \phantom{+ 0x + 0} \\
 x^2 + 0x \phantom{+ 0} \\
 \underline{x^2 - x} \phantom{+ 0} \\
 2x \phantom{+ 0} \\
 \underline{2x - 2} \\
 2
 \end{array}$$

$$\begin{aligned}
 & \int \frac{x^3+x}{x-1} dx \\
 &= \int \left( \textcolor{red}{x^2} + \textcolor{red}{x} + \textcolor{red}{2} + \frac{2}{x-1} \right) dx
 \end{aligned}$$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

$$\begin{aligned}
 & \int \frac{x^3+x}{x-1} dx \\
 &= \int \left( x^2 + x + 2 + \frac{2}{x-1} \right) dx
 \end{aligned}$$

## Example

Find  $\int \frac{x^3+x}{x-1} dx$ .

$$\begin{array}{r}
 x^2 + x + 2 \\
 x-1 \overline{) x^3 \phantom{+ 2x^2} + x} \\
 \underline{x^3 - x^2} \phantom{+ 2x} \\
 x^2 + x \\
 \underline{x^2 - x} \\
 2x \\
 \underline{2x - 2} \\
 2
 \end{array}$$

$$\begin{aligned}
 & \int \frac{x^3+x}{x-1} dx \\
 &= \int \left( x^2 + x + 2 + \frac{2}{x-1} \right) dx \\
 &= \frac{x^3}{3} + \frac{x^2}{2} + 2x \\
 & \quad + 2 \ln |x-1| + C
 \end{aligned}$$

- The next step in producing a partial fraction decomposition is to factor the denominator  $Q(x)$ .

- The next step in producing a partial fraction decomposition is to factor the denominator  $Q(x)$ .
- Factoring of  $Q(x)$  can always be done in quadratic and linear terms as asserted in the following.

### Corollary (Corollary to the Fundamental Theorem of Algebra)

*Let  $Q(x)$  be a polynomial (with real coefficients). Then  $Q(x)$  can be factored as a product of terms of the form  $(ax + b)^n$  (powers of linear terms) and product of terms of the form  $(ax^2 + bx + c)^n$  with  $b^2 - 4ac < 0$  (powers of quadratic terms).*

- The next step in producing a partial fraction decomposition is to factor the denominator  $Q(x)$ .
- Factoring of  $Q(x)$  can always be done in quadratic and linear terms as asserted in the following.

### Corollary (Corollary to the Fundamental Theorem of Algebra)

*Let  $Q(x)$  be a polynomial (with real coefficients). Then  $Q(x)$  can be factored as a product of terms of the form  $(ax + b)^n$  (powers of linear terms) and product of terms of the form  $(ax^2 + bx + c)^n$  with  $b^2 - 4ac < 0$  (powers of quadratic terms).*

- The above result is a corollary to the Fundamental Theorem of Algebra.



- The next step in producing a partial fraction decomposition is to factor the denominator  $Q(x)$ .
- Factoring of  $Q(x)$  can always be done in quadratic and linear terms as asserted in the following.

### Corollary (Corollary to the Fundamental Theorem of Algebra)

*Let  $Q(x)$  be a polynomial (with real coefficients). Then  $Q(x)$  can be factored as a product of terms of the form  $(ax + b)^n$  (powers of linear terms) and product of terms of the form  $(ax^2 + bx + c)^n$  with  $b^2 - 4ac < 0$  (powers of quadratic terms).*

- The above result is a corollary to the Fundamental Theorem of Algebra. We state the Fundamental Theorem of algebra without proving it.

### Theorem (The Fundamental Theorem of Algebra)

*Every polynomial has at least one complex root.*

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$  and  
the  $B_j$  and  $C_j$ 's are constants - one pair for each power  $1 \leq j \leq M$ .

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$  and the  $B_j$  and  $C_j$ 's are constants - **one pair for each power  $1 \leq j \leq M$ .**



- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$  and the  $B_j$  and  $C_j$ 's are constants - one pair for each power  $1 \leq j \leq M$ .

- We use  $N$  different constants for each new linear factor of the form  $(ax + b)^N$  and  $2 \times M$  different constants for each factor of the form  $(ax^2 + bx + c)^N$ .

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$  and the  $B_j$  and  $C_j$ 's are constants - one pair for each power  $1 \leq j \leq M$ .

- We use  $N$  different constants for each new linear factor of the form  $(ax + b)^N$  and  $2 \times M$  different constants for each factor of the form  $(ax^2 + bx + c)^N$ .
- Thus the total number of constants used equals the degree of  $Q$ .

- Let  $\frac{R(x)}{Q(x)}$  be a rational function with  $\deg Q > \deg R$ .
- Suppose  $Q(x)$  factors into factors of the form

$$(ax + b)^N \quad \text{and} \quad (ax^2 + bx + c)^M.$$

- Then we can split  $\frac{R(x)}{Q(x)}$  into sum of partial fractions of the form

$$\frac{A_i}{(ax + b)^i}, \text{ with } i \leq N \quad \text{or} \quad \frac{B_j x + C_j}{(ax^2 + bx + c)^j}, \text{ with } j \leq M,$$

where the  $A_i$ 's are constants - one for each power  $1 \leq i \leq N$  and the  $B_j$  and  $C_j$ 's are constants - one pair for each power  $1 \leq j \leq M$ .

- We use  $N$  different constants for each new linear factor of the form  $(ax + b)^N$  and  $2 \times M$  different constants for each factor of the form  $(ax^2 + bx + c)^N$ .
- Thus the total number of constants used equals the degree of  $Q$ .
- The difficulty of finding the constants  $A_i, B_j, C_j$  increases as the number of distinct factors increases, as well as when the exponents of those factors increase.

# $Q(x)$ has distinct linear factors

- Suppose  $Q(x)$  is a product of distinct linear factors:

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_kx + b_k)$$

where no factor is repeated and no factor is a constant multiple of another.

## $Q(x)$ has distinct linear factors

- Suppose  $Q(x)$  is a product of distinct linear factors:

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_kx + b_k)$$

where no factor is repeated and no factor is a constant multiple of another.

- Then there exist constants  $A_1, A_2, \dots, A_k$  such that

$$\frac{R(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_k}{a_kx + b_k}$$

## $Q(x)$ has distinct linear factors

- Suppose  $Q(x)$  is a product of distinct linear factors:

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_kx + b_k)$$

where no factor is repeated and no factor is a constant multiple of another.

- Then there exist constants  $A_1, A_2, \dots, A_k$  such that

$$\frac{R(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_k}{a_kx + b_k}$$

- We show how to find  $A_1, A_2, \dots, A_k$  on examples.

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.



## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$



## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \end{aligned}$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \end{aligned}$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{\textcolor{red}{B}}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{\textcolor{red}{1}}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \end{aligned}$$

Solution:

$$A = \frac{1}{2}, \textcolor{red}{B} = \frac{1}{5}, C = -\frac{1}{10}.$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \end{aligned}$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \\ &= \frac{1}{2} \ln |x| + \frac{1}{10} \ln |2x-1| \\ & \quad - \frac{1}{10} \ln |x+2| + K \end{aligned}$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \\ &= \frac{1}{2} \ln|x| + \frac{1}{10} \ln|2x-1| \\ & \quad - \frac{1}{10} \ln|x+2| + K \end{aligned}$$

## Example

Find  $\int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx$ .

- $\deg(x^2 + 2x - 1) < \deg(2x^3 + 3x^2 - 2x)$ : don't divide.
- Factor denominator:  $2x^3 + 3x^2 - 2x = x(2x - 1)(x + 2)$ .

$$\frac{x^2 + 2x - 1}{x(2x - 1)(x + 2)} = \frac{A}{x} + \frac{B}{2x - 1} + \frac{C}{x + 2}$$

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

$$x^2 + 2x - 1 = (2A + B + 2C)x^2 + (3A + 2B - C)x - 2A$$

$$2A + B + 2C = 1$$

$$3A + 2B - C = 2$$

$$-2A = -1$$

Solution:

$$A = \frac{1}{2}, B = \frac{1}{5}, C = -\frac{1}{10}.$$

$$\begin{aligned} & \int \frac{x^2+2x-1}{2x^3+3x^2-2x} dx \\ &= \int \left( \frac{1}{2} \frac{1}{x} + \frac{1}{5} \frac{1}{2x-1} - \frac{1}{10} \frac{1}{x+2} \right) dx \\ &= \frac{1}{2} \ln |x| + \frac{1}{10} \ln |2x - 1| \\ & \quad - \frac{1}{10} \ln |x + 2| + K \end{aligned}$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$



NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ;

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ;

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$0^2 + 2 \cdot 0 - 1 = A(2 \cdot 0 - 1)(0 + 2)$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \end{aligned}$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 = B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right)$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 &= B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right) \\ \frac{1}{4} &= \frac{5}{4}B \end{aligned}$$



NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 &= B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right) \\ \frac{1}{4} &= \frac{5}{4}B \\ B &= \frac{1}{5} \end{aligned}$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 &= B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right) \\ \frac{1}{4} &= \frac{5}{4}B \\ B &= \frac{1}{5} \end{aligned}$$

$$(-2)^2 + 2(-2) - 1 = C(-2)(2(-2) - 1)$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 &= B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right) \\ \frac{1}{4} &= \frac{5}{4}B \\ B &= \frac{1}{5} \end{aligned}$$

$$\begin{aligned} (-2)^2 + 2(-2) - 1 &= C(-2)(2(-2) - 1) \\ -1 &= 10C \end{aligned}$$

NOTE: There is a quick trick to find  $A$ ,  $B$ , and  $C$ .

$$x^2 + 2x - 1 = A(2x - 1)(x + 2) + Bx(x + 2) + Cx(2x - 1)$$

To find  $A$ , set  $x = 0$ ; to find  $B$ , set  $x = \frac{1}{2}$ ; to find  $C$ , set  $x = -2$ .

$$\begin{aligned} 0^2 + 2 \cdot 0 - 1 &= A(2 \cdot 0 - 1)(0 + 2) \\ -1 &= -2A \\ A &= \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \left(\frac{1}{2}\right)^2 + 2 \cdot \frac{1}{2} - 1 &= B\left(\frac{1}{2}\right)\left(\frac{1}{2} + 2\right) \\ \frac{1}{4} &= \frac{5}{4}B \\ B &= \frac{1}{5} \end{aligned}$$

$$\begin{aligned} (-2)^2 + 2(-2) - 1 &= C(-2)(2(-2) - 1) \\ -1 &= 10C \\ C &= -\frac{1}{10} \end{aligned}$$

## $Q(x)$ has linear factors with higher multiplicity

- Suppose  $Q(x)$  is a product of linear factors, some of which appear with power greater than 1.

# $Q(x)$ has linear factors with higher multiplicity

- Suppose  $Q(x)$  is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power  $r$ , that is,  $(a_1x + b_1)^r$  occurs in the factorization of  $Q(x)$ .

## $Q(x)$ has linear factors with higher multiplicity

- Suppose  $Q(x)$  is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power  $r$ , that is,  $(a_1x + b_1)^r$  occurs in the factorization of  $Q(x)$ .
- Then instead of a single term  $\frac{A}{a_1x+b_1}$  we use

$$\frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \cdots + \frac{A_r}{(a_1x + b_1)^r}$$

## $Q(x)$ has linear factors with higher multiplicity

- Suppose  $Q(x)$  is a product of linear factors, some of which appear with power greater than 1.
- For example suppose the first linear factor has power  $r$ , that is,  $(a_1x + b_1)^r$  occurs in the factorization of  $Q(x)$ .
- Then instead of a single term  $\frac{A}{a_1x+b_1}$  we use

$$\frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \cdots + \frac{A_r}{(a_1x + b_1)^r}$$

- In a similar fashion we add more partial fractions to account for all other terms of the form  $(a_sx + b_s)^t$ .



## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$
- Factor denominator:  $x^3 - x^2 - x + 1 = ?$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1}$
- Factor denominator:  $x^3 - x^2 - x + 1 = (x - 1)^2(x + 1)$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = ?$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$



## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$
$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$
$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$
$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = ?$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$
$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
 
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
 
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1)$



## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
 
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = ?$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:
$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$
$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:  

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$
- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$

- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$

- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$

- Plug-in  $x = 0$ :

$$-2 = A(0-1)(0+1) + B(0+1) + C(0-1)^2$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$

- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$

- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$

- Plug-in  $x = 0$ :

$$-2 = A(0-1)(0+1) + B(0+1) + C(0-1)^2$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}.$

- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$

- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$

- Plug-in  $x = 0$ :

$$-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = ?$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx$$

- Divide:  $\frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{x^3 - x^2 - x + 1} = x + 2 + \frac{-x^2 + 5x - 2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .

- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .



## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

• Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}.$

• Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1).$

• Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

• Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2.$

• Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1.$

• Plug-in  $x = 0$ :

$$-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1.$$

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

= ?

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .



## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## Example

$$\int \frac{x^4 + x^3 - 4x^2 + 4x}{x^3 - x^2 - x + 1} dx = \int \left( x + 2 + \frac{1}{x-1} + \frac{1}{(x-1)^2} - \frac{2}{x+1} \right) dx$$

$$= \frac{x^2}{2} + 2x + \ln|x-1| - \frac{1}{x-1} - 2\ln|x+1| + K$$

- Divide:  $\frac{x^4+x^3-4x^2+4x}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{x^3-x^2-x+1} = x + 2 + \frac{-x^2+5x-2}{(x-1)^2(x+1)}$ .
- Factor denominator:  $x^3 - x^2 - x + 1 = (x-1)^2(x+1)$ .
- Set up the partial fraction decomposition:

$$\frac{-x^2 + 5x - 2}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$-x^2 + 5x - 2 = A(x-1)(x+1) + B(x+1) + C(x-1)^2$$

- Plug-in  $x = -1$ :  $-(-1)^2 + 5(-1) - 2 = C(-1-1)^2 \Rightarrow C = -2$ .
- Plug-in  $x = 1$ :  $-(1)^2 + 1 \cdot 5 - 2 = B(1+1) \Rightarrow B = 1$ .
- Plug-in  $x = 0$ :  
 $-2 = A(0-1)(0+1) + 1 \cdot (0+1) + (-2)(0-1)^2 \Rightarrow A = 1$ .

## $Q(x)$ contains quadratic factors, multiplicity 1

- Suppose  $Q(x)$  contains quadratic factors  $ax^2 + bx + c$  with where  $b^2 - 4ac < 0$  (i.e., the factor is irreducible).

## $Q(x)$ contains quadratic factors, multiplicity 1

- Suppose  $Q(x)$  contains quadratic factors  $ax^2 + bx + c$  with where  $b^2 - 4ac < 0$  (i.e., the factor is irreducible).
- Suppose none of the quadratic factors is repeated.

# $Q(x)$ contains quadratic factors, multiplicity 1

- Suppose  $Q(x)$  contains quadratic factors  $ax^2 + bx + c$  with where  $b^2 - 4ac < 0$  (i.e., the factor is irreducible).
- Suppose none of the quadratic factors is repeated.
- The for each quadratic factor we need to add a partial fraction of the form

$$\frac{Ax + B}{ax^2 + bx + c}.$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .



## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx$$



## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{\textcolor{red}{A}}{\textcolor{red}{x}} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$\textcolor{red}{A} = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \left( \frac{\textcolor{red}{1}}{\textcolor{red}{x}} + \frac{x - 1}{x^2 + 4} \right) dx$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 4}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \end{aligned}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \end{aligned}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \\ &= ? \end{aligned}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \\ &= \ln|x| + \frac{1}{2} \ln(x^2 + 4) - \frac{1}{2} \arctan\left(\frac{x}{2}\right) + K \end{aligned}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \\ &= \ln|x| + \frac{1}{2} \ln(x^2 + 4) - \frac{1}{2} \arctan\left(\frac{x}{2}\right) + K \end{aligned}$$



## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \\ &= \ln|x| + \frac{1}{2} \ln(x^2 + 4) - \frac{1}{2} \arctan\left(\frac{x}{2}\right) + K \end{aligned}$$

## Example

Find  $\int \frac{2x^2 - x + 4}{x^3 + 4x} dx$ .

- $\deg(2x^2 - x + 4) < \deg(x^3 + 4x)$ : don't divide.
- Factor denominator:  $x^3 + 4x = x(x^2 + 4)$ .

$$\frac{2x^2 - x + 4}{x(x^2 + 4)} = \frac{A}{x} + \frac{Bx + C}{(x^2 + 4)}$$

$$2x^2 - x + 4 = A(x^2 + 4) + (Bx + C)x$$

$$2x^2 - x + 4 = (A + B)x^2 + Cx + 4A$$

$$A = 1 \quad C = -1 \quad A + B = 2, \text{ therefore } B = 1$$

$$\begin{aligned} \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx &= \int \left( \frac{1}{x} + \frac{x - 1}{x^2 + 4} \right) dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2 + 4} dx - \int \frac{1}{x^2 + 4} dx \\ &= \ln|x| + \frac{1}{2} \ln(x^2 + 4) - \frac{1}{2} \arctan\left(\frac{x}{2}\right) + K \end{aligned}$$

# $Q(x)$ has quadratic factors with multiplicity $> 1$

- Suppose  $Q(x)$  has the factor  $(ax^2 + bx + c)^r$ , where  $b^2 - 4ac < 0$  and  $r > 1$ .

# $Q(x)$ has quadratic factors with multiplicity $> 1$

- Suppose  $Q(x)$  has the factor  $(ax^2 + bx + c)^r$ , where  $b^2 - 4ac < 0$  and  $r > 1$ .
- Then the partial fraction decomposition should include summands of the form

$$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_rx + B_r}{(ax^2 + bx + c)^r}$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^3}$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2+x+1)(x^2+1)^3}$$

= ?

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2+x+1)(x^2+1)^3}$$

$$= \frac{A}{x} +$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(\textcolor{red}{x} - \textcolor{red}{1})(x^2 + x + 1)(x^2 + 1)^3}$$

$$= \frac{A}{x} + \textcolor{red}{?}$$



## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^3}$$

$$= \frac{A}{x} + \frac{B}{x-1} +$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^3}$$

$$= \frac{A}{x} + \frac{B}{x-1} + ?$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^3}$$
$$= \frac{A}{x} + \frac{B}{x-1} + \frac{Cx + D}{x^2 + x + 1} +$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2+x+1)(x^2+1)^3}$$
$$= \frac{A}{x} + \frac{B}{x-1} + \frac{Cx+D}{x^2+x+1} + ?$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2+x+1)(x^2+1)^3}$$
$$= \frac{A}{x} + \frac{B}{x-1} + \frac{Cx+D}{x^2+x+1} + \frac{Ex+F}{x^2+1} + \frac{Gx+H}{(x^2+1)^2} + \frac{Ix+J}{(x^2+1)^3}.$$

## Example

Write out the form of the partial fraction decomposition of

$$\frac{x^3 + x^2 + 1}{x(x-1)(x^2 + x + 1)(x^2 + 1)^3}$$

$$= \frac{A}{x} + \frac{B}{x-1} + \frac{Cx + D}{x^2 + x + 1} + \frac{Ex + F}{x^2 + 1} + \frac{Gx + H}{(x^2 + 1)^2} + \frac{Ix + J}{(x^2 + 1)^3}.$$

For example of this size it makes sense to use a computer algebra system; one such system easily produces the decomposition:

$$= \frac{-1}{x} + \frac{\frac{1}{8}}{x-1} + \frac{-x-1}{(x^2 + x + 1)} + \frac{\frac{15}{8}x - \frac{1}{8}}{(x^2 + 1)} + \frac{\frac{3}{4}x + \frac{3}{4}}{(x^2 + 1)^2} + \frac{-\frac{x}{2} + \frac{1}{2}}{(x^2 + 1)^3}.$$